


61 20 Fr. Qajor

IN MEMORIAM FLORIAN CAJORI





BRADY'S

BOOKSTOR

BALTIMORE, MI

This ?

ZZZZZZZENIA ZZZZ

Digitized by the Internet Archive in 2008 with funding from Microsoft Corporation

INTRODUCTION

TO

ALGEBRA,

WITH

NOTES AND OBSERVATION

DESIGNED FOR THE

USE OF SCHOOLS AND PLACES OF PUBLIC EDUCATION.

TO WHICH IS ADDED

AN APPENDIX

ON THE

APPLICATION OF ALGEBRA TO GEOMETRY.

9.

By JOHN BONNYCASTLE,
Professor of Mathematics in the Royal Military Academy, Woolwich.

SECOND NEW-YORK, FROM THE ELEVENTH LONDON EDITION.

REVISED, CORRECTED, AND ENLARGED, WITH A VARIETY OF EXAMPLES,

AND MANY OTHER USEFUL ADDITIONS.

By JAMES RYAN, Private Teacher of Mathematics.

Ingenus didicisse fideliter artes Emollit mores, nec sinit esse ferosi Ovid.

NEW-YORK:

PUBLISHED BY EVERT DUYCKINCK AND GEORGE LONG.

1822.

CAJORI

Southern District of New-York, ss.

BE IT REMEMBERED, That on the twenty-eighth day of December, in the

DL 11 ALBIERBLEKED, That on the twenty-eighn day of December, in the forty-sixty near of the independence of the United Strets of America, George Long, of the said district, hath deposited in this office the title of a book, the right whereof he claims as proprietor, in the words following, to wir.

"An Introduction to Algebra, with Notes and Observations; designed for the Use of An Introduction to Algebra, with Notes and Observations; designed for the Use of the American and plant of the Country. By Jehn Bampeantle, Professor of Machematics in the Royal Military Academy, Woolwich. Second New-York from the Elevant London Edition. Revised, corrected, and enlared, with a variety of Experts. Eleventh London Edition. Revised, corrected, and enlarged, with a variety of Examples, and many other useful Additions, by James Ryan, Private Teacher of Mathematics.

 Ingenuas didicisse fideliter artes Emollit mores, nee sinit esse feros.

In conformity to the act of Congress of the United States, entitled, "An act for the encouragement of learning, by securing the copies of maps, charts, and books, to the aithors and proprietors of such copies, during the time therein mentioned. And also to an act, entitled, "a net, supplementary to an act, entitled, An act for the encouragement of learning, by securing the copies of maps, charts, and hooks, to the authors and proprietors of such copies, during the times therein mentioned, and examines. tending the benefits thereof to the arts of designing, engraving, and etching historical and other prints."

JAMES DILL.

Clerk of the Southern District of New-York.

PREFACE.

THE powers of the mind, like those of the body are increased by frequent exertion; application and industry supply the place of genius and invention; and even the creative faculty itself may be strengthened and improved by use and perseverance. Uncultivated nature is uniformly rude and imbecile, it being by imitation alone that we at first acquire knowledge, and the means of extending its bounds. A just and perfect acquaintance with the sumple elements of science, is a necessary step towards our future progress and advancement; and this, assisted by laborious investigation and habitual inquiry, will constantly lead to eminence and perfection.

Books of rudiments, therefore, concisely written, well digested, and methodically arranged, are treasures of inestimable value; and too many attempts cannot be made to render them perfect and complete. When the first principles of any art or science are firmly fixed and rooted in the mind, their application soon becomes easy, pleasant, and obvious; the understanding is delighted and enlarged; we conceive clearly, reason distinctly, and form just and and satisfactory conclusions. But, on the contrary, when the mind, instead of reposing on the stability of truth and received principles, is wandering in doubt and uncertainty, our ideas will necessarily be confused and obscure; and every step we take must be attended with fresh difficulties and endless perplexity.

That the grounds, or fundamental parts, of every science, are dull and unentertaining, is a complaint universally made, and a truth not to be denied; but then, what is obtained with difficulty is usually remembered with ease; and what is purchased with pain is often possessed with pleasure. The seeds of knowledge are sown in every soil, but it is by proper culture alone that they are cherished and brought to maturity. A few years of early and assiduous application, never fails to procure us the reward of our industry; and who is there, who knows the pleasures and advantages which the sciences afford, that would think his time, in this case mis-spent, or his labours useless? Riches and honours are the gifts of fortune, casually bestowed, or hereditarily received, and are frequently abused by their possessors; but the superiority of wisdom and knowledge is a pre eminence of merit, which originates with the man, and is the noblest of all distinctions.

Nature, bountiful and wise in all things, has provided us with an infinite variety of scenes, both for our instruction and entertainment; and, like a kind and indulgent parent, admits all her children to an equal participation of her blessings. But, as the modes, situations and circumstances of life are various, so accident, habit and education, have each their predominating influence, and give to every mind its particular bias. Where examples of excellence are wanting, the attempts to attain it are but few; but eminence excites attention, and produces imitation. To raise the curiosity, and to awaken the listless and dormant powers of younger minds, we have only to point out to them a valuable acquisitiou, and the means of obtaining it; the active principles are immediately put into motion, and the certainty of the conquest is ensured from a determination

to conquer.

But, of all the sciences which serve to call forth this spirit of enterprise and inquiry, there are none more eminently useful than Mathematics. By an early attachment to these elegant and sublime studies, we acquire a habit of reasoning, and an elevation of thought, which fixes the mind, and prepares it for every other pursuit. From a few simple axioms, and evident principles, we proceed

gradually to the most general propositions, and remote analogies: deducing one truth from another, in a chain of argument well connected and logically pursued; which brings us at last, in the most satisfactory manner, to the conclusion, and serves as a general direction in all our in-

quiries after truth.

And it is not only in this respect that mathematical learning is so highly valuable; it is, likewise equally estimable for its practical utility. Almost all the works of art and devices of man, have a dependence upon its principles, and are indebted to it for their origin and perfection. The cultivation of these admirable sciences is, therefore, a thing of the utmost importance, and ought to be considered as a principal part of every liberal and well-regulated plan of education. They are the guide of our youth, the perfection of our reason, and the foundation of every great and noble undertaking.

From these considerations, I have been induced to compose an introductory course of mathematical science; andfrom the kind encouragement which I have hitherto received, am not without hopes of a continuance of the same. candour and approbation. Considerable practice as a teacher, and a long attention to the difficulties and obstructions which retard the progress of learners in general, have enabled me to accommodate myself the more easily to their capacities and understandings. And as an earnest desire of promoting and diffusing useful knowledge is the chief motive for this undertaking, so no pains or attention shall be wanting to make it as complete and perfect as possible.

The subject of the present performance is ALGEBRA: which is one of the most important and useful branches of those sciences, and may be justly considered as the key to all the rest. Geometry delights us by the simplicity of its principles, and the elegance of its demonstrations: Arithmetic is confined in its object, and partial in its application; but Algebra, or the Analytic Art, is general and comprehensive, and may be applied with success in all cases where truth is to be obtained and proper data can be established.

To trace this science to its birth, and to point out the various alterations and improvements it has undergone in

its progress, would far exceed the limits of a preface.* It will be sufficient to observe that the invention is of the highest antiquity, and has challenged the praise and admiration of all ages. Diophantus, a Greek mathematician, of Alexandria in Egypt, who flourished in or about the third century after Christ, appears to have been the first, among the ancients, who applied it to the solution of indeterminate or unlimited problems; but it is to the moderns that we are principally indebted for the most curious refinements of the art, and its great and extensive usefulness in every abstruse and difficult inquiry. Newton, Maclaurin, Sanderson, Simpson, and Emerson, among our own countrymen, and Clairant, Euler, Lagrange and Lacroix, on the continent, are those who have particularly excelled in this respect; and it is to their works that I would refer the young student, as the patterns of elegance and perfection.

The following compendium is formed entirely upon the model of those writers, and is intended as a useful and necessary introduction to them. Almost every subject, which belongs to pure Algebra, is concisely and distinctly treated of; and no pains have been spared to make the whole as easy and intelligible as possible. A great number of elementary books have already been written upon this subject : but there are none, which I have yet seen, but what appear to me to be extremely defective. Besides being totally unfit for the purpose of teaching, they are generally calculated to vitiate the taste, and mislead the judgment. A tedious and inelegant method prevails through the whole, so that the beauty of the science is generally destroyed by the clumsy and awkward manner in which it is treated; and the learner, when he is afterwards introduced to some of our best writers, is obliged, in a great measure, to unlearn and forget every thing which he has been at so much pains in acquiring.

There is a certain taste and elegance in the sciences, as

a Those who are desirous of a knowledge of this kind, may consult the Introduction to my Treatise on Algebra; where they will find a regular historical detail of the rise and progress of the science, from its first rude beginnings to the present times.

well as in every branch of polite literature, which is only to be obtained from the best authors, and a judicious use of their instructions. To direct the student in his choice of books, and to prepare him properly for the advantages he may receive from them, is therefore, the business of every writer who engages in the humble, but useful task of a preliminary tutor.—This information I have been careful to give, in every part of the present performance, where it appeared to be in the least necessary; and, though the nature and confined limits of my plan admitted not of diffuse observations, or a formal enumeration of particulars, it is presumed nothing of real use and importance has been omitted. My principal object was to consult the ease, satisfaction, and accommodation of the learner; and the favourable reception the work has met with from the public, has afforded me the gratification of believing that my labours have not been unsuccessfully emyloyed.

The state of the s

ර ද මුදා ,ලැබ් වේ රෝග් වෙමින් . ර අතර රෝග ලැප න්රේර්

4

gradient gradient

ADVERTISEMENT.

The present performance having passed through a number of editions since the time of its first publication, without any material alterations having been made, either with respect to its original plan, or the manner in which it was executed, I have been induced, from the flattering approbation it has constantly received, to undertake an entire revision of the work; and, by availing myself of the improvements that have been subsequently made in the science, to render it still more deserving the public favour.

In its present state, it may be considered as a copious abridgment of the most practical and useful parts of my larger work, entitled, A Treatise on Algebra, in 2 vols. 8vo. published in 1813; from which, except in certain cases, where a different mode of proceeding appeared to be necessary, it has been chiefly compiled: great care having been taken, at the

same time, to adapt it, as much as possible, to the wants of learners, and the general purposes of instruction, agreeably to the design with which it was first written.

With this view, as well as in compliance with the wishes of several intelligent teachers, I have also been led to subjoin to it, by way of an Appendix, a small tract on the application of Algebra to the solution of Geometrical Problems; which, it is hoped, will prove acceptable to such classes of students as may not have an opportunity of consulting more voluminous and expensive works on this interesting branch of the science.

JOHN BONNYCASTLE.

ROYAI. MILITARY ACADEMY, Woolwich. October 22, 1815.

ADVERTISEMENT

THE SECOND NEW-YORK EDITION.

Ir would be superfluous to advance any thing in commendation of "Bonnycastle's Introduction to Algebra," as the number of European editions, and the increase of demand for it since its publication in this country, are sufficient proofs of its great utility.

But to make it universally useful both to the tutor and scholar, I have given in this edition, the answers that were omitted by the author in the original.

In the course of the work, particularly in Addition, Subtraction, Multiplication, Division, Fractions, Simple Equations, and Quadratics, I have added a great variety of practical examples, as being essentially necessary to exercise young students in the elementary principles.

Several new rules are introduced, those of principal note are the following: Case 12.

Surds, containing two rules for finding any root of a Binomial Surd, the Solution of Cubics by Converging Series, the Solution of Biquadratics by Simpson's and Euler's methods: all these rules are investigated in the plainest manner possible, with notes and remarks, interspersed throughout the work, containing some very useful matter.

There is also given all the Dipohantine Analysis, contained in Bonnycastle's Algebra, Vol. 1. 8vo. 1820., being a methodical abstract of this part of the science, which comprehends most of the methods hitherto known for resolving problems of this kind, and will be found a ready compendium for such readers as may acquire some knowledge of the Analytic Art.

JAMES RYAN.

New-York, Jan. 1, 1822.

P. S. A new and correct Key to the present edition, comprising in an easy and elegant manner, the solutions of the questions, is now ready for the press, and will be speedily published. J. R.

CONTENTS.

							-0
DEFINITIONS		2	*				. 1
Addition		-			٠,		8
Subtraction		•	•	•		•	12
Multiplication		- •	٠,			٠,	14
Division							19
Algebraic Fractions			٠		•	•	27
Involution, or the Raising of Powers						٠	44
Evolution, or the Extraction of Roots		1/1	٠	. "	٠,	•	47
Of Irrational Quantities, or Surds				•		*	54
Of Arithmetical Proportion and Progression						٠	87
Of Geometrical Proportion and Progression	٠,٠		4.				91
Of Equations						٠	99
Of the Resolution of Simple Equations							101
Miscellaneous Questions						٠	119
Of Quadratic Equations							128
Questions producing Quadratic Equations					٠.		139
Of Cubic Equations							148
Of the Solution of Cubic Equations							150
Of the Solution of Cubic Equations by Converging S	erie	es .					156
Of the Resolution of Biquadratic Equations							168
To find the Roots of Equations by Approximation							176
To find the Roots of Exponential Equations							182
Of the Binomial Theorem							184
Of the Indeterminate Analysis							190
Of the Diophantine Analysis							201
Of the Summation and Interpolation of Series							233
Of Logarithms							256
Multiplication by Logarithms							268
Division by Logarithms							271
The Rule of Three by Logarithms							273
Involution, by Logarithms							275
Evolution, by Logarithms	.5						277
A Collection of Miscellaneous Questions							280
Appendix, on the Application of Algebra to Geomet	rv						286

ALGEBRA.

ALGEBRA is the science which treats of a general method of performing calculations, and resolving mathematical problems, by means of the letters of the alphabet.

Its leading rules are the same as those of drithmetic; and the operations to be performed are denoted by the

following characters:

+ plus, or more, the sign of addition; signifying that the quantities between which it is placed are to be added together.

Thus, a+b shows that the number, or quantity, represented by b, is to be added to that represented by a; and is read a plus b.

— minus, or less, the sign of subtraction; signifying that the latter of the two quantities between which it is placed is to be taken from the former.

Thus a-b shows that the quantity represented by b is to be taken from that represented by a; and is read a minus b.

Also, ash represents the difference of the two quantities a and b, when it is not known which of them is the greater.

 \times into, the sign of multiplication; signifying that the quantities between which it is placed are to be multiplied together.

Thus, $a \times b$ shows that the quantity represented by a is to be multiplied by that represented by b; and is read a into b.

The multiplication of simple quantities is also frequently denoted by a point, or by joining the letters together in the form of a word. Thus, $a \times b$, a. b, and ab, all signify the product of a and b: also, $3 \times a$, or 3a, is the product of 3 and a; and is read 3 times a.

by, the sign of division; signifying that the former of the two quantities between which it is placed is to be divided by the latter.

Thus, $a \div b$, shows that the quantity represented by a is to be divided by that represented by b; and is read a by b, or a divided by b.

Division is also frequently denoted by placing one of the two quantities over the other, in the form of a fraction.

Thus, $b \div a$ and $\frac{b}{a}$ both signify the quotient of b divided by a; and $\frac{a-b}{a+c}$ signifies that a-b is to be divided by a+c.

= equal to, the sign of equality; signifying that the quantities between which it is placed are equal to each other.

Thus, x=a+b shows that the quantity denoted by x is equal to the sum of the quantities a and b; and is read x equal to a plus b.

<u> identical</u> to, or the sign of equivalence; signifying that the expressions between which it is placed are of the same value, for all values of the letters of which they are composed.

* Thus, $(x+a) \times (x-a) \pm x^2 - a^2$, whatever numeral values may be given to the quantities represented by x and a.

^{*} Euler calls x-1=x-1 an identical equation; and shows that x is incleaning; or that any number whatever may be substituted for it; which equation is to be expressed according to the present notation; thus, x-1=x-1. In my opinion, Euler's definition of an indentical equation, is preferable to Bonnycastles'; See Euler's Algebra page 239. Vol.

> greater than, the sign of majority; signifying that the former of the two quantities between which it is placed is greater than the latter.

Thus, $a \triangle b$ shows that the quantity represented by a is greater than that represented by b; and is read a greater than b.

< less than, the sign of minority; signifying that the former of the two quantities between which it is placed is less than the latter.

Thus, $a \angle b$ shows that the quantity represented by a is less than that represented by b; and is read a less than b.

: as, or to, and :: so is, the signs of an equality of ratios; signifying that the quantities between which they are placed are proportional.

Thus; a:b::c:d denotes that a has the same ratio to b that c has to d, or that a, b, c, d, are proportionals; and is read, as a is to b so is c to d, or a is to b as c is to d.

√ the radical sign, signifying that the quantity before which it is placed is to have some root of it extracted.

Thus, \sqrt{a} is the square root of a; $\sqrt[3]{a}$ is the cube root of a; and $\sqrt[4]{a}$ is the fourth root of a; &c.

The roots of quantities are also represented by figures placed at the right hand corner of them, in the form of a fraction.

Thus, $a^{\frac{1}{2}}$ is the square root of a; $a^{\frac{1}{3}}$ is the cube root of a; and $a^{\frac{1}{n}}$ is the nth root of a, or a root denoted by any number n

In like manner, a^2 is the square of a; a^3 is the cube of a; and a^m is the mth power of a, or any power denoted by the number m.

co is the sign of infinity, signifying that the quantity

standing before it is of an unlimited value, or greater than any quantity that can be assigned.

The coefficient of a quantity is the number or letter which is prefixed to it.

Thus, in the quantities 3b, $-\frac{a}{3}b$, 3 and $-\frac{a}{3}$ are the coefficients of b; and a is the coefficient of x in the quantity ax.

A quantity without any coefficient prefixed to it is supposed to have 1 or unity; and when a quantity has no sign before it, + is always understood.

Thus, a is the same as +a, or +1a; and -a is the same as -1a.

A term is any part or member of a compound quantity, which is separated from the rest by the signs + or -.

Thus, a and b are the terms of a + b; and 3a, -2b, and +5cd, are the terms of 3a - 2b + 5cd.

In like manner, the terms of a product, fraction, or proportion, are the several parts or quantities of which they are composed.

Thus, a and b are the terms of ab, or of $\frac{a}{b}$; and a, b, c, d, are the terms of the proportion a:b::c:d.

A factor is one of the terms, or multipliers which form

the product of two or more quantities.

A composite number, or quantity, is that which is produced by the multiplication of two or more terms or fac-

tors.

Thus, 6 is a composite number, formed of the factors 2 and 3, or 2×3 ; and 3abc is a composite quantity, the factors of which are 3, a, b, c.

Like quantities, are those which consist of the same letters or combinations of letters; as a and 3a, or 5ab

and 7ab, or 2a2b and 9a2b.

Unlike quantities are those which consist of different letters, or combinations of letters; as a and b, or 3a and a^2 , or $5ab^2$ and $7a^2b$.

Given quantities are such as have known values, and are generally represented by some of the first letters of the alphabet; as a, b, c, d, &c.

Unknown quantities, are such as have no fixed values, and are usually represented by some of the final letters of the alphabet; as x, y, z.

Simple quantities, are those which consist of one term only; as 3a, 5ab, $= 8a^2b$, &c.

Compound quantities, are those which consist of several terms; as 2a+b, or 3a-2c, or a+2b-3c, &c.

Positive, or affirmative quantities, are those which are to be added; as a, or +a, or +3ab, &c.

Negative quantities, are those which are to be subtracted; as -a, or -3ab, or $-7ab^2$, &c.

Like signs, are such as are all positive, or all negative; as + and +, or - and -.

Unlike signs, are when some are positive and others negative; as + and -; or - and +.

A monomial, is a quantity consisting of one term only: as a, 2b, $-3a^2b$, &c.

A binomial, is a quantity consisting of two terms; as a+b, or a-b; the latter of which is, also, sometimes called a residual quantity.

A trinomial, is a quantity consisting of three terms, as a+2b-3c; a quadrinomial of four, as a-2b+3c-d: and a polynomial, or multinomial, is that which has many terms.

The power of a quantity, is its square, cube, biquadrate, &c.; called also its second, third, fourth power, &c.; as a^2 , a^3 , a^4 , &c.

The index, or exponent of a quantity, is the number which denotes its power or root.

Thus, -1 is the index of a-1, 2 is the index of a^2 , and 1 of a or 1/a.

When a quantity appears without any index, or expo-

nent, it is always understood to have unity, or 1.

Thus, a is the same as a^1 , and 2x is the same as $2x^1$;

the 1, in such cases, being usually omitted.

A rational quantity, is that which can be expressed in finite terms, or without any radical sign, or fractional index; as a, or $\frac{2}{3}a$, or 5a, &c.

* An irrational Quantity, or Surd, is that of which the value cannot be accurately expressed in numbers, as the square roots of 2, 3, 5. Surds are commonly expressed by means of the radical sign 1; as 1/2, 1/a, 4/a3, or a fractional index; as $2^{\frac{1}{2}}$, $a^{\frac{2}{3}}$, &c.

A square or cube number, &c. is that which has an ex-

act square or cube root. &c.

Thus, 4 and $\frac{9}{16}a^2$ are square numbers; and 64 and 3 a are cube numbers, &c.

A measure of any quantity, is that by which it can be

divided without leaving a remainder.

Thus, 3 is a measure of 6, 7a is a measure of 35a, and 2ab of 27 a2b2.

Commensurable quantities, are such as can be each divided by the same quantity, without leaving a remainder.

Thus, 6 and 8, $2\sqrt{2}$ and $3\sqrt{2}$, $5a^2b$ and $7ab^2$, are commensurable quantities; the common divisors being 2, $\sqrt{2}$, and ab.

Incommensurable quantities, are such as have no common measure, or divisor, except unity.

Thus, 15 and 16, $\sqrt{2}$ and $\sqrt{3}$, and a+b and a^2+b^2 , are incommensurable quantities.

^{*} Bonnycastle's definition of a surd was erroneous. Dr. Hutton was led into the same error, until his mathematics was published in New-York, revised and corrected by Robert Adrain, LL.D. Professor of Mathematics and Natural Philosophy Columbia College. Afterwards, Dr. Hutton published the seventh edition of his mathematics, in which he adopted this definition, (which I have copied from his edition) without mentioning who was the Author. EDITOR.

A multiple of any quantity, is that which is some exact number of times that quantity.

Thus, 12 is a multiple of 4, 15a is a multiple of 3a, and

20a2b2 of 5ab.

The reciprocal of any quantity, is that quantity inverted, or unity divided by it.

Thus, the reciprocal of a, or $\frac{a}{1}$, is $\frac{1}{a}$; and the reciprocal of $\frac{a}{7}$ is $\frac{b}{2}$.

A function of one or more quantities, is an expression into which those quantities enter, in any manner whatever, either combined, or not, with known quantities.

Thus, a-2x, $ax+3x^2$, 2x-a $(a^2-x^2)^{\frac{1}{2}}$, ax^m , a^x , &c., are functions of x; and $axy + bx^2$, $ay + x (ax^2-by^2)^{\frac{1}{2}}$, &c. are functions of x and y.

A vinculum, is a bar -, or parenthesis (), made

use of to collect several quantities into one.

Thus, $\overline{a+b} \times c$, or (a+b) c, denotes that the compound quantity a+b is to be multiplied by the simple quantity c; and $\sqrt{ab+c^2}$, or $(ab+c^2)^{\frac{1}{2}}$, is the square root of the compound quantity $ab+c^2$.

Practical Examples for computing the numeral Values of various Algebraic Expressions, or Combinations of Letters.

Supposing a=6, b=5, c=4, d=1, and e=0.

- 1. $a^2 + 2ab c + d = 36 + 60 4 + 1 = 93$.
- 2. $2a^3 3a^2b + c^3 = 432 540 + 64 = -44$.
- 3. $a^2 \times \overline{a+b} 2abc = 36 \times 11 240 = 156$.
- 4. $2a\sqrt{b^2-ac}+\sqrt{2ac+c^2}=12\times 1+8=20$.
- 5. $3a\sqrt{2ac+c^2}$, or $3a(2ac+c^2)^{\frac{1}{2}}=18\sqrt{64}=144$.
- 6. $\sqrt{2a^2} \sqrt{2ac + c^2} = \sqrt{72 \sqrt{64}} = \sqrt{72 8} = \sqrt{64}$ =8.
- 7. $\frac{2a+3c}{6d+4e} + \frac{4bc}{\sqrt{2ac+c^2}} = \frac{12+12}{6+0} + \frac{80}{\sqrt{48+16}} = \frac{24}{6} + \frac{80}{8} = 14.$

Required the numeral values of the following quantities: supposing a, b, c, d, e, to be 6, 5, 4, 1, and 0, respectively, as above.

- 1. $2a^2 + 3bc 5d = 127$
- 2. $5a^2b 10ab^2 + 2e = -600$
- 3. $7a^2 + b c \times d + e = 253$
- 4. $5\sqrt{ab+b^2-2ab-e^2}=-7.613371$
- $5. \frac{a}{c} \times d \frac{a-b}{d} + 2a^2 e = \frac{1}{2}$
- 6. $3\sqrt{c+2a\sqrt{2a+b-d}}=14$
- 7. $a\sqrt{a^2+b^2+3bc\sqrt{a^2-b^2}}=245.8589862$
- 3. $3a^{2}b + \sqrt[3]{c^{2}} + \sqrt{2ac+c^{2}} = 542.8844991$ 9. $\frac{2b+c}{3a-c} \frac{\sqrt{5b+3}\sqrt{c+d}}{2a+c} = \frac{1}{4}$

ADDITION.

Addition is the connecting of quantities together by means of their proper signs, and incorporating such as are like, or that can be united, into one sum; the rule for performing which is commonly divided into the three following cases*:

CASE I.

When the Quantities are like, and have like Signs.

RULE.

Add all the coefficients of the several quantities together, and to their sum annex the letter or letters belonging, to each term, prefixing, when necessary, the common sign.

^{*} The term Addition, which is generally used to denote this rule, is too scanty to express the nature of the operations that are to be performed in it; which are sometimes those of addition, and sometimes subtraction, according as the quantities are negative or positive. It should, therefore, be called by some name signifying incorporation, or striking a balance; in which case, the incongruity, here mentioned, would be removed.

/	EXAMPLES.	
3a	— 3ax	2b + 3y
	$\frac{-3ax}{-6ax}$	5b+7y
бa		b+2y
ia .	/ax	
7a	— 2ax	8b+y
12a	7ax	4b+4y
		201145
28a	-19ax	20b + 17y
		-
2ay	$-2by^2$	$a-2x^2$
5ay	$-6by^2$	$a-6x^2$
4ay	/by2	4a-x2
7ay	$-8by^2$	$3a-5x^2$
16ay	$-by^2$	$7a-x^{2}$
34ay	$-18by^{3}$	$16u - 15x^2$
- 1 -		0.1:0
$3ax^2$	7x-4y	$2a+x^2$
2ax2	€ - 8y	$3a + x^2$
$12ax^2$	3x-y	$\alpha + 2x^{g}$
$9ax^2$	x-3y	9a + 3x2
$10ax^2$	4x-y	$4a + x^3$
2	40 47	10.10.5
36ax2	16x - 17y	$19a + 8x^2$

CASE II.

When the Quantities are like, but have unlike signs.

RULE.

Add all the affirmative coefficients into one sum, and those that are negative into another, when there are se-

veral of the same kind; then subtract the least of these sums from the greatest, and to the difference prefix the sign of the greater, annexing the common letter or letters as before.

	EXAMPLES.	
-3a +7a +8a	$ 2a - 3x^{2} -7a + 5x^{2} -3a + x^{2} $	6x + 5ay $ -3x + 2ay $ $ x - 6ay$
-a	$+a-3x^2$	2x+ay
-11a	_7a *	6x+2ay
$-2a^{2}$ $-3a^{2}$	2ay—7 — ay+8	-3ab + 7x + 3ab - 10x
- 8a2	+2ay-9	+3ab-6x
$+10a^2 + 13a^2$	-3ay - 11 + 12ay + 13	-ab-2x +2ab+7x
+130-	712ay 715	7240714
$+10a^{2}$	1+12ay-6	+4ab-4x
-		
$-2a\sqrt{x}$	$-6a^{2}+2b$	$6ax^2 + 5x^{\frac{1}{2}}$
$+ a\sqrt{x}$	$+2a^2-3b$	$-2ax^2-6x^{\frac{1}{2}}$
$-3a\sqrt{x}$	$-5a^2-8b$	$+3ax^2-10x^{\frac{1}{2}}$
$+7a\sqrt{x}$	$+4a^2-2b$	$-7ax^2+3x^{\frac{1}{2}}$
$-4a\sqrt{x}$	$-3a^2+9b$	$+ax^2+11x^{\frac{1}{2}}$
-		. 1
$-a\sqrt{x}$	$-8a^2-2b$	$+ax^{2}+3x^{\frac{1}{2}}$

CASE III.

When the Quantities are unlike; or some like and others unlike.

RULE.

Collect all the like quantities together, by taking their sums or differences, as in the foregoing cases, and set down those that are unlike, one after another, with their proper signs.

	EXAMPLES.	10.0
5 xy 4ax	$2xy - 2x^2$ $3x^2 + xy$	$2ax - 30$ $3x^2 - 2ax$
-xy	$x^2 + xy$	$5x^2 - 3x^{\frac{1}{2}}$
$\frac{-4\alpha x}{4xy}$	$\frac{4x^2 - 3xy}{6x^2 + xy}$	$\frac{3\sqrt{x+10}}{8x^2-20}$
		And the second s
$+ax^{\frac{1}{2}}$ $-ax^2$	$8a^2x^2 - 3ax$ $7a x - 5xy$	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
$+3ax^2$ $-ax^{\frac{1}{2}}$	$9x y - 5ax$ $2a^2x^2 + xy$	$50 + 2a^2x$ $a^2x^2 + 120$
$+2ax^2$ 10	$a^2x^2 + 5xy - ax 96$	$b^2 + 3a^2x^2 - a^2x + 170$
$+3a^2y$	$2\sqrt{x-18y}$	$2a^2 - 3a\sqrt{x}$
$\begin{array}{l} -2xy^2 \\ -3y^2x \end{array}$	$3\sqrt{xy+10x} \\ 2x^2y + 25y$	$x^{2} - 2a^{\frac{1}{2}}x^{\frac{1}{2}}$ $3a^{2} - 13xy$
$-8x^2y$	$15xy - \sqrt{xy}$	$xy + 32a^{3}$
$\frac{+2xy^2}{}$	$-8y + 17x^{\frac{1}{2}}$	$\frac{20 - 65x^2}{2}$
$y - 3y^2x - 8x$	$\langle +2x^2y+2\sqrt{x}$	$\begin{cases} 37a^2 - 3a\sqrt{x - 12x} \\ y - 64x^2 - 2a^{\frac{1}{2}}x^{\frac{1}{2}} + \end{cases}$
	(y+10x-y)) on

302

EXAMPLES FOR PRACTICE.

1. Required the sum of $\frac{1}{2}(\alpha+b)$ and $\frac{1}{2}(\alpha-b)$. Ans. a. 2. Add 5x - 3a + b + 7 and -4a - 3x + 2b - 9 to

gether. 2x - 3a + 3b + 7 and 2x - 3a + 2b - 3agether. 2x - 7a + 3b - 2.

3. Add 2a + 3b - 4c - 9 and 5a - 3b + 2c - 10 together. Ans. 7a - 2c - 19.

4. Add 3a + 2b - 5, a + 5b - c, and 6a - 2c + 3 toether.

Ans. 10a + 7b - 3c - 2.

gether. Ans. 10a + 7b - 3c - 2. 5. Add $x^3 + ax^2 + bx + 2$ and $x^3 + cx^2 + dx - 1$ to-

gether. Ans. $2x^3 + (a+c) x^2 + (b+d) x + 1$. 5. Add $6xy - 12x^2$, $-4x^2 + 3xy$, $4x^2 - 2xy$, and -3xy

5. Add $6xy - 12x^2$, $-4x^2 + 3xy$, $4x^2 - 2xy$, and $-3x + 4x^2$ together. Ans. $4xy - 8x^2$

7. Add $4ax-130+3x\frac{1}{2}$, $5x^2+3ax+9x^2$, $7xy-4x^2+90$, and $\sqrt{x+40-6x^2}$ together. Ans. $7ax+8x^2+7xy$. 8. Add $2a^2-3ab+2b^3-3a^2$, $3b^3-2a^2+a^3-5c^3$, $4c^3$

6. Add $2a^2 - 3ab + 2b^2 - 3a^2 + 3b^2 - 3c^3 + 5ab + 100$, and $20ab + 16a^2 - bc - 80$ together.

• Ans. $13a^2 + 22ab + 3b^3 + a^3 - c^3 + 20 - bc$.

9. Add
$$\frac{5a}{b} - \frac{3c^2}{a} + \frac{7\sqrt{bc}}{x} - 9(\frac{ab+x}{d})$$
 and $\frac{8a}{b} + \frac{7c^2}{a} - 12\frac{\sqrt{bc}}{x}$ $+6(\frac{ab+x}{t})$ together.

Ans.
$$\frac{13a}{b} - \frac{4c}{a} - 5\frac{\sqrt{bc}}{x} - 3(\frac{ab+x}{d}).$$

10. Add $3a^2 + 4bc - e^2 + 10$, $-5a^2 + 6bc + 2e^2 - 15$, and $-4a^2 - 9bc - 10e_2^2 + 21$ together.

Ans. $bc-6a^2-9e^2+16$.

SUBTRACTION.

SUBTRACTION is the taking of one quantity from another; or the method of finding the difference between any two quantities of the same kind; which is performed as follows *:

^{*} This rule being the reverse of addition, the method of operation must be so likewise. It depends upon this principle, that to subtract an affirmative quantity from an affirmative, is the same as to add a negative quantity to as affirmative.

RULE.

Change all the signs (+ and -) of the lower line, or quantities that are to be subtracted, into the contrary signs, or rather conceive them to be so changed, and then collect the terms together, as in the several cases of addition.

ddition.		
	EXAMPLES.	
$5a^2 - 2b$	$x^2 - 2y + 3$	5xy + 8x - 2
$2a^{2}+5b$	$4x^2 + 9y - 5$	3xy - 8x - 7
$\frac{3a^2-7b}{}$	$-3x^2 - 11y + 8$	2xy + 16x + 5
5xy-18	$8y^2 - 2y - 5$	10 - 8x - 3xy
-xy+12	$-y^2 + 3y + 2$	-x+3-xy
6xy-30	$9y^2 - 5y - 7$	7-7x-2xy
$-5x^2y - 8a$	$4\sqrt{ax-2x^2y}$	$0x^2 + \sqrt{x - 4y}$
$+3x^2y-7b$	$3\sqrt{ax-5xy^2}$	$6x^2 - 8x - x_{\frac{1}{2}}$
$-8x^2y - 8a + 7b$	$\sqrt{ax-2x^2y+5xy^2}$	$x^2 + 8x + 2\sqrt{x - 4y}$

EXAMPLES FOR PRACTICE.

- 1. Find the difference of $\frac{1}{2}(a+b)$ and $\frac{1}{2}(a-b)$. Ans. b.
- 2. From 3x 2a b + 7, take 8 3b + a + 4x.

Ans. 2b - x - 3a - 1.

3. From 3a+b+c-2d, take b-8c+2d-8.

Ans. 3a+9c-4d+8.

4. From $13x^2 - 2ax + 9b^2$, take $5x^2 - 7ax - b^2$.

Ans. $8x^2 + 5ax + 10b^2$.

That is, -3a taken from +5a, is the same with +3a added to +5a=+8a; because, -3a+3a=0, and +5a+3a=+8a; hence, if I take 0 from +8a, there remains +8a, but, if the same quantity or equal quantities be added to the subtrahend and minuend, their difference continues still to be the same as if nothing was added to them: therefore -3a taken from +5a=+8a. And in like manner +3a taken from -5a=-3a-5a=-5a=-8a.

5. From $20ax - 5\sqrt{x+3a}$ take $4ax + 5x^{\frac{1}{2}} - a$.

Ans. $16ax - 10\sqrt{x + 4a}$.

- 6. From $5ab+2b^2-c+bc-b$, take $b^2-2ab+bc$.

 Ans. $7ab+b^2-c-b$.
- 7. From $ax^3 bx^2 + cx d$, take $bx^2 + ex 2d$. Ans. $ax^3 - 2bx^2 + (c - e)x + d$.
- 8. From 6a 4b 12c + 13x, take 4x 9a + 4b 5c.

 Ans. 3a + 9x 8b 7c.
 - 9. From $6x^2y 3\sqrt{xy} 6ay$, take $3x^2y + 3(xy)\frac{1}{2}$
- 4ay. Ans. $3x^2y 6\sqrt{xy} 2ay$.
 10. From the sum of $4ax 150 + 4x^{\frac{1}{2}}$, $5x^2 + 3ax + 10x^{\frac{1}{2}}$, and $90 2ax 12\sqrt{x}$; take the sum of $2ax 80 + 7x^2$, $7x^{\frac{1}{2}} 8ax 70$, and $30 4\sqrt{x} 2x^2 + 4a^2x^2$.

 Ans. $11ax + 60 x^{\frac{1}{2}} 4a^2x^2$.

MULTIPLICATION.

MULTIFLICATION, or the finding of the product of two or more quantities, is performed in the same manner as in arithmetic; except that it is usual, in this case, to begin the operation at the left hand, and to proceed towards the right, or contrary to the way of multiplying numbers.

The rule is commonly divided into three cases; in each of which, it is necessary to observe, that like signs, in

multiplying, produce +, and unlike signs, -.

It is likewise to be remarked, that powers, or roots of the same quantity, are multiplied together by adding their indices: thus.

 $\varepsilon \times a^2$, or $a^1 \times a^2 = a^3$; $a^2 \times a^3 = a^5$; $a^{\frac{1}{3}} \times a^{\frac{1}{3}} = a^{\frac{5}{6}}$; and $a^m \times a^n = a^{m \cdot n}$.

The multiplication of compound quantities, is also, sometimes, barely denoted by writing them down, with their proper signs, under a vinculum, without performing the whole operation, as

 $3ab\ (a-b)$, or $2a\sqrt{a^2+b^2}$.

Which method is often preferable to that of executing the entire process, particularly when the product of two or more factors is to be divided by some other quantity, because, in this case, any quantity that is common to both the divisor and dividend, may be more readily suppressed; as will be evident from various instances in the following part of the work*.

CASE I.

When the factors are both simple quantities.

RULE.

Multiply the coefficients of the two terms together, and to the product annex all the letters, or their powers, belonging to each, after the manner of a word; and the result, with the proper sign prefixed, will be the product required †.

^{*} The above rule for the signs may be proved thus: If n, b, be any two quantities, of which a is the greater, and $\mathbf{s} - b$ is to be multiplied by a, it is plain that the product, in this case, must be less than an, because $\mathbf{s} - b$ is less than \mathbf{s} ; and, consequently, when each of the terms of the former are multiplied by a, as above, the result will be

 $⁽B-b) \times a = aB - ab$. For if it were aB + ab, the product would be greater than aB, which is ab-

Also, if B be greater than b, and A greater than a, and it is required to mul-

tiply B - b by A - a, the result will be

For the product of $\mathbf{B} - \mathbf{b}$ by $(\mathbf{A} - \mathbf{a}) = \mathbf{A} \mathbf{B} - \mathbf{a} \mathbf{B} - \mathbf{b} \mathbf{A} + ab$. For the product of $\mathbf{B} - \mathbf{b}$ by \mathbf{A} is \mathbf{A} ($\mathbf{B} - \mathbf{b}$), or $\mathbf{A} \mathbf{B} - \mathbf{A}\mathbf{b}$, and that of $\mathbf{B} - \mathbf{b}$ by $-\mathbf{a}$, which is to be taken from the former, is $-\mathbf{a}(\mathbf{B} - \mathbf{b})$, as has been already shown; whence $\mathbf{B} - \mathbf{b}$ being less than $\mathbf{a}_{\mathbf{B}}$; it is evident that the part which is to be taken away must be less than $\mathbf{a}_{\mathbf{B}}$; and consequently since the first part of this product is $-\mathbf{a}_{\mathbf{B}}$, the second part must be $+\mathbf{a}\mathbf{b}$; for if it were $-\mathbf{a}\mathbf{b}$, a greater part than $\mathbf{a}\mathbf{B}$ would be to be taken from $\mathbf{A}(\mathbf{B} - \mathbf{b})$, which is absurd.

[†] When any number of quantities are to be multiplied together, it is the same thing in whatever order they are placed: thus, if ab is to be multiplied by c, the product is either abc, acb, or bca, &c.; though it is usual, in this case, as well as in addition and subtraction, to put them according to their rank in the alphabet. It may here also be observed, in conformity to the rule given above for the signs, that $(+a) \times (+b)$, or $(-a) \times (-b) = \pm ab$; and $(+a) \times (-b)$, or $(-a) \times (-b) = \pm ab$;

EXAMPLES.

12a	-2a	+5a	$-9x^{2}$
36	+46	- 6.x	- 5bx
36ab	07	-30ax	$+45bx^{3}$
3000	-8ab	-30ax	+430%
7ab	$-6a^2x$	$-2xy^2$	-7axy
- bac	+5x	-xy	+6ay
35a2 bc	$-30a^{2}x^{2}$	$+2x^{2}y^{3}$	-42a2xy
500 00		, 2.c g	120 29
3a2 b	12a2x	-6xyz	$-a^2xy$
2b a 2	-2x y	$+ay^2z$	$+2xy^*$
6a b	$-24a^2x^3y$	$-6axy^3z^2$	$-2a^2x^2y^3$
0000	- 240 x g	- ou.eg 2	- ~u g

CASE II.

When one of the factors is a compound quantity.

BULE.

Multiply every term of the compound factor, considered as a multiplicand, separately, by the multiplier, as in the former case; then these products, placed one after another, with their proper signs, will be the whole product required.

EXAMPLES.

3a-2b	6xy - 8	$a^2 - 2x + 1$
4a	3x	4 <i>x</i>
$12a^3-8ab$	$18x^2y - 24x$	$4a^2x - 8x^2 + 4x$

$ \begin{array}{c} 12x - ab \\ 5a \end{array} $	35x - 7a $ -2x$	$3y^2 + y - 2$ xy
$60ax - 5a^2b$	$\phantom{100000000000000000000000000000000000$	$3xy^3 + xy^2 - 2xy$
$ \begin{array}{r} 13x^2 - a^2b \\ -2a \end{array} $	$25xy + 3a^2$ $13x^2$	$3x^2 - xy - 2y^2$ $5x^2$
$\frac{-26ax^2+2a^3b}{-26ax^2+2a^3b}$	$325x^3y + 3\theta a^2x^2$	$15x^4 - 5x^3y - 10x^2y^2$

CASE III.

When both the factors are compound quantities.

RULE.

Multiply every term of the multiplicand separately, by each term of the multiplier. setting down the products one after another, with their proper signs; then add the several lines of products together, and their sum will be the whole product required.

EXAMPLES.

$ \begin{array}{c} x+y\\x+y \end{array} $	5x + 4y $3x - 2y$	$\begin{array}{c} x^2 + xy - y \\ x - y \end{array}$
$\begin{array}{c} \overline{x^2 + xy} \\ +xy + y^2 \end{array}$	$ \begin{array}{r} 15x^2 + 12xy \\ -10xy - 8y^2 \end{array} $	$ \begin{array}{r} x^3 + x^2 y - xy^2 \\ - x^2 y - xy^2 + y^3 \end{array} $
$\overline{x^2 + 2xy + y^2}$	$15x^2 + 2xy - 8y^2$	$x^3 + -2xy^2 + y^3$
$ \begin{array}{c} x+y\\x-y \end{array} $	$\begin{array}{c} x^2 + y \\ x^2 + y \end{array}$	$\begin{array}{c} x^2 + xy + y^2 \\ x - y \end{array}$
$\overline{x^2 + xy} $ $-xy - y^2$	$\begin{array}{c} x^4 + x^2 y \\ + x^2 y + y^2 \end{array}$	$\begin{array}{c} x^{3} + x^{2}y + xy^{2} \\ -x^{2}y - xy^{2} - y^{3} \end{array}$
$x^{2} * - y^{2}$	$x^4 + 2x^2y + y^2$	*x3 * * * y3

EXAMPLES FOR PRACTICE.

1, Required the product of $x^2 - xy + y^2$ and x + y.

Ans. $x^3 + y^3$.

2. Required the product of $x^3+x^2y+xy^2+y^3$ and x-y.

3. Required the product of $x^3 + xy + y^2$ and $x^2 - xy + xy + x^3$.

4. Required the product of $3x^2-2xy+5$, and $x^2+2xy-3$.

Ans. $3x^4+4x^3y-4x^2y^2-4x^2+16xy-15$.

5. Required the product of $2a^2 - 3ax + 4x^2y - 4x^2y + 16xy - 15$. $5ax - 2x^2$. $fas. <math>10a^4 - 27a^3x + 34a^2x^2 - 18ax^3 - 8x^4$.

6. Required the product of $5x^3 + 4ax^2 - 18ax^3 - 8a^3$. $2x^2 - 3ax + a^2$. Ans. $10x^5 - 7ax^4 - a^2x^3 - 3a^3x^2 + a^5$

7. Required the product of $3x^3 + 2x^2y^2 + 3y^3$ and $2x^3 + 3x^3 + 3$

 $-3x^{2}y^{2} + 5y^{3}.$ Ans. $6x^{6} - 5x^{5}y^{2} - 6x^{4}y^{4} + 21x^{3}y^{3} + x^{2}y^{5} + 15x^{5}.$

8. Required the product of $x^3 - ax^2 + bx - c$ and $x^2 - dx + e$. Ans. $x^5 - ax^4 - dx^4 + (b + ad + e)x^3 - (c + bd + ae)$ $x^2 + (cd + eb)x - ce$.

9. † Required the product of the four following fact-

I. III. IV. $(a+b)(a^2+ab+b^2)(a-b)$ and (a^2-ab+b^2) .

Ans. $a^{6} - b^{6}$.

10. Required the product of $a^{3} + 3a^{2}x + 3ax^{2} + x^{3}$ and $a^{3} - 3a^{2}x + 3ax^{2} - x^{3}$.

Ans. $a^6 - 3a^4x^2 + 3a^2x^4 - x^6$.

11. Required the product of $a^4 + a^2c^2 + c^4$ and $a^2 - c^2$.

Ans. $a^6 - c^6$.

12. Required the product of $a^2 + b^2 + c^2 - ab - ac - bc$ and a + b + c. Ans. $a^3 - 3abc + b^3 + c^3$.

[†] I would advise the learner to perform the calculation of this example several ways, viz. First, by multiplying the product of the factors II. and IV. Secondly, by multiplying the product of the factors II. and IV. Secondly, by multiplying the product of the factors II. and IV. Thirdly, by multiplying the product of the factors I. and IV. by the product of the factors II. and III. The last method is the most concise; See Euler's Algebra page 119. Vol. I.

DIVISION.

Division is the converse of multiplication, and is performed like that of numbers; the rule being usually divided into three cases; in each of which like signs give + in the quotient, and unlike signs -, as in finding their products*.

It is here also to be observed, that powers and roots of the same quantity, are divided by subtracting the index of

the divisor from that of the dividend.

Thus,
$$a^3 \div a^2$$
, or $\frac{a_3^3}{a^3} = a$; $a^{\frac{1}{2}} \div a^{\frac{1}{3}}$, or $\frac{a_{\frac{1}{2}}}{a_{\frac{1}{3}}} = a^{\frac{1}{6}}$; $a^{\frac{3}{4}} \div x^{\frac{2}{3}}$, or $\frac{a^{\frac{3}{4}}}{a^{\frac{2}{3}}} = a^{\frac{1}{2}}$; and $a^m \div a^n$, or $\frac{a^m}{a^n} = a^{m-n}$.

CASE I.

When the divisor and dividend are both simple quantities.

RULE.

Set the dividend over the divisor, in the manner of a raction, and reduce it to it simplest form, by cancelling he letters and figures that are common to each term.

EXAMPLES.

$$6ab \div 2a$$
, or $\frac{6ab}{2a} = 3b$; and $12ax^2 \div 3x$, or $\frac{12ax^2}{3x} = 4ax$;
 $4 \div a$, or $\frac{a}{a} = 1$; and $a \div -a$, or $\frac{a}{-a} = -1$.

* According to the rule here given for the signs, it follows that

$$\frac{+ab}{+b} = +a, \frac{-ab}{-b} = +a, \frac{-ab}{+b} = -a, \frac{+ab}{-b} = -a,$$

s will readily appear by multiplying the quotient by the divisor; the signs of the products being then the same as would take place in the former rule.

Also
$$-2a \div 3a$$
, or $\frac{-2a}{3a} = -\frac{2}{3}$; and $9x^{\frac{1}{2}} \div 3x^{\frac{1}{4}} = 3x^{\frac{1}{4}}$.

1. Divide 16x2 by 8x, and 12a2x2 by -- Sa2x.

Ans.
$$2x$$
, and $-\frac{3x}{2}$.

2. Divide $-15ay^2$ by 3ay and $-18ax^2y$ by -8ax. Ans. -5y, and $\frac{9xy}{4}$

3. Divide
$$-\frac{2}{3}a^{\frac{1}{2}}$$
, by $\frac{1}{5}a^{\frac{1}{2}}$, and $ax^{\frac{1}{3}}$ by $-\frac{3}{5}a^{\frac{1}{2}}x^{\frac{1}{4}}$.

Ans. $-3\frac{1}{3}$, and $-\frac{5}{3}a^{\frac{1}{2}x^{\frac{1}{2}x}}$. 4. Divide $12a^{2}b^{2}$ by $-3a^{2}b$, and $-15ay^{\frac{3}{2}}$ by $-3ay^{\frac{1}{2}}$.

Ans. — 4b, and $5y_0^2$. 5. Divide — $15a^2x^2$ by $5ax^2$, and $21 \ a^2c^2x^{\frac{1}{2}}$ by — 7a

S. Divide $= 15u^2x^2$ by $5ux^2$, and $21u^2-2x^2$ by $= 7u^2$.

Ans. = 3u, and $= 3ux^{\frac{1}{4}}$.

6. Divide — $17x^{\frac{1}{2}}a^{3}c$ by — $5x^{\frac{1}{3}}a^{2}c^{\frac{1}{2}}$ and $24\sqrt{xy}$ by $3\sqrt{xy}$.

Ans. $\frac{17x^{\frac{1}{6}}ac^{\frac{1}{2}}}{5}$, and $3\sqrt{xy}$.

CASE II.

When the divisor is a simple quantity, and the dividend a compound one.

RULE.

Divide each term of the dividend by the divisor, as in the former case; setting down such as will not divide in the simplest form they will admit of.

EXAMPLES

$$(ab+b^2) \div 2b$$
, or $\frac{ab+b^2}{2b} = \frac{1}{2}a + \frac{1}{2}b = \frac{a+b}{2}$
 $(10ab-15ax) \div 5a$, or $\frac{10ab-15ax}{5a} = 2b - 3x$.

$$(30ax - 48x^2) \div 6x$$
, or $\frac{30ax - 48x^2}{6x} = 5a - 8x$

1. Let $3x^3+6x^2+3ax-15x$ be divided by 3x. Ans. $x^2+2x+a-5$.

2. Let 3abc-+12abx-9a2b be divided by 3ab. Ans. c+4x-3a.

3. Let $40a^3b^3 + 60a^2b^2 - 17ab$ be divided by -ab.

Ans. $-40a^2b^2-60ab+17$. 4. Let $15a^2bc-12acx^2+5ad^2$ be divided by -5ac.

4. Let $15a^2bc - 12acx^2 + 5ad^2$ be divided by -5ac.

Ans. $-3ab + \frac{12x^2}{5} - \frac{d^2}{c}$.

5. Let $20ax^3 + 15ax^2 + 10ax + 5a$ be divided by 5a.

Ans. $4x^3 + 3x^2 + 2x + 1$.

6. Let $6bcdz + 4bzd^2 - 2b^2z^2$ be divided by 2bz. Ans. $3cd + 2d^2 - bz$.

7. Let $14a^2 - 7ab + 21ax - 28a$ be divided by 7a.

Ans. 2a - b + 3x - 4.

8. Let $-20ab + 60ab^3 - 12a^2b^2$ be divided by -4ab. Ans. $5 - 15b^2 + 3ab$.

9. Let $15a^2bc - 12acx^2 + 5ad^2$ be divided by -5ac. Ans. $-3ab + \frac{12x^2}{5} - \frac{d^2}{c}$.

CASE III.

When the divisor and dividend are both compound quantities.

RULE.

Set them down in the same manner as in division of numbers, ranging the terms of each of them so, that the higher powers of one of the letters may stand before the lower.

Then divide the first term of the dividend by the first term of the divisor, and set the result in the quotient, with its proper sign, or simply by itself, if it be affirmaive. This being done, multiply the whole divisor by the term thus found; and, having subtracted the result from the dividend, bring down as many terms to the remainder as are requisite for the next operation, which perform as before; and so on, till the work is finished, as in common arithmetic.

EXAMPLES.

$$\begin{array}{c}
x+y)x^{2}+2xy+y^{2}(x+y) \\
x^{2}+xy \\
\hline
xy+y^{2} \\
xy+y^{2}
\end{array}$$

$$\begin{array}{r}
a+x)a^3+5a^2x+5ax^2+x^3(a^2+4ax+x^2) \\
\underline{a^3+a^2x} \\
\underline{4a^2x+5ax^2} \\
\underline{4a^2x+4ax^2} \\
\underline{ax^2+x^3} \\
\underline{ax^2+x^3}
\end{array}$$

$$\begin{array}{r}
x-3)x^3-9x^2+27x-27(x^2-6x+9) \\
x^3-3x^2 \\
-6x^2+27x \\
-6x^2+18x \\
\hline
9x-27 \\
9x-27
\end{array}$$

$$\begin{array}{r}
2x^{3} - 3ax + a^{2})4x^{4} - 9a^{2}x^{2} + 6a^{3}x - a^{4}(2x^{2} + 3ax - a^{2} + 4x^{4} - 6ax^{3} + 2a^{2}x^{2} \\
\underline{6ax^{3} - 11a^{2}x^{2} + 6a^{3}x \\
6ax^{3} - 9a^{2}x^{2} + 3a^{3}x - a^{4} \\
-2a^{2}x^{2} + 3a^{3}x - a^{4}
-2a^{2}x^{2} + 3a^{3}x - a^{4}
\end{array}$$

Note 1. If the divisor be not exactly contained in the dividend, the quantity that remains after the division is finished, must be placed over the divisor, at the end of the quotient, in the form of a fraction: thus*.

$$\begin{array}{r}
 a+x)a^{3}-x^{3}(a^{2}-ax+x^{2}-\frac{2x^{3}}{a+x}) \\
 & -a^{3}+a^{2}x \\
 & -a^{2}x-x^{3} \\
 & -a^{2}x-ax^{2} \\
 & -a^{2}x-ax^{3} \\
 & -a^{2}+x^{3} \\
 & -2x^{3}
\end{array}$$

[•] In the case here given, the operation of division may be considered as rminated, when the highest power of the letter, in the first or leading term the remainder, by which the process is regulated, is less than the power the first term of the divisor; or when the first term of the divisor is not antained in the first term of the remainder; as the succeeding part of the totient, after this, instead of being integral, as it ought to be, would necessity become fractional.

$$\begin{array}{c}
x+y)x^4+y^4(x^3-x^2y+xy^2-y^3+\frac{2y^4}{x+y} \\
\underline{x^4+x^3y} \\
-x^3y+y^4 \\
\underline{-x^3y-x^2y^2} \\
\underline{x^2y^2+y^4} \\
\underline{x^2y^2+xy^3} \\
\underline{-xy^3+y^4} \\
-xy^3-y^4
\end{array}$$

2. The division of quantities may also be sometimes carried on, ad infinitum, like a decimal fraction; in which case, a few of the leading terms of the quotient will generally be sufficient to indicate the rest, without its being necessary to continue the operation; thus,

^{*} Now, it is easy to perceive that the next or 6th term of the quotient will be $-\frac{xs}{as}$, and the seventh term $\frac{x}{as}$, and so on, alternately plus and minus;

And by a process similar to the above, it may be shown that

$$\frac{a}{a-x} = 1 + \frac{x}{a} + \frac{x^2}{a^2} + \frac{x^3}{a^3} + \frac{x^4}{a^4} + \frac{x^5}{a^5} + \&c.$$

Where the law, by which either of these series may be continued at pleasure, is obvious*.

this is called the law of continuation of the series. And the sum of all the terms when infinitely continued is said to be equal to the fraction $\frac{a}{a+x}$.

Thus we say the vulgar fraction $\frac{2}{r}$ when reduced to a decimal is =22222, &c. infinitely continued. The terms in the quotient are found by dividing the remainders by a, the first term of the divisor; thus, the first remainder -x divided by a, gives $-\frac{x}{x}$ the second term in the quotient; and the second re-

mainder $+\frac{x^2}{a}$ divided by a gives $+\frac{x^2}{a^2}$ the third term, &c.

* In this example, if x be less than a, the series is convergent, or the value of the terms continually diminish; but when x is greater than a, it is said to diverge.

To explain this by numbers: suppose a = 3, and x = 2. Then, $1 + \frac{x}{a} + \frac{x^2}{a^2} + \frac{x^3}{a^2}$, &c.

Then,
$$1 + \frac{x}{a} + \frac{x^2}{a^2} + \frac{x^3}{a^2}$$
, &

The corresponding values are, $1 + \frac{2}{3} + \frac{4}{9} + \frac{3}{27}$, &c.

$$1 + \frac{2}{3} + \frac{4}{9} + \frac{8}{27}$$
, &c.

where the fractions or terms of the series grow less and less, and the farther they are extended, the more they converge or approximate to 0, which is supposed to be the last term or limit. But if $\alpha=2$, and $\alpha=3$.

Then,
$$1 + \frac{x}{a} + \frac{x^2}{a^2} + \frac{x^3}{a^3}$$
, &c.

The corresponding values are,
$$1 + \frac{3}{2} + \frac{9}{4} + \frac{27}{8}, &c.$$

In which the terms become larger and larger. This is called a diverging series.

If
$$\alpha = 1$$
, and $\alpha = 1$ in the preceding example:
Then, $\frac{\alpha}{a+x} = 1 - \frac{x}{a} + \frac{x^2}{a^2} - \frac{x^3}{a^3}$, &c. will be $\frac{1}{1+1} = 1 - 1 + 1 - 1$, &c.

Now, because $\frac{1}{1+1} = \frac{1}{2}$, it has been said that 1-1+1-1, &c. infinitely con-

tinued, is =1: a singular conclusion, when it is perceived from the terms themselves, that their sum must necessarily be either 0 or +1, to whatever

EXAMPLES FOR PRACTICE.

1. Let $a^2-2ax+x^2$ be divided by a-x. Ans. a-x.

2. Let $x^3 - 3ax^2 + 3a^2x - a^3$ be divided by x - a.

Ans. $x^2 - 2ax + a^2$.

3. Let $a^3+5a^2x+5ax^2+x^3$ be divided by a+x. $a^2+4ax+x^2$.

4. Let $2y^3 - 19y^2 + 26y - 16$ be divided by y - 8. $Ans. 2y^2 - 3y + 2$.

5. Let x^5+1 be divided by x+1, and x^6-1 by x-1.

Ans. $x^4 - x^3 + x^2 - x + 1$, and $x^5 + x^4 + x^3 + x^2 + x + 1$.

6. Let $48x^3 - 76ax^2 - 64a^2x + 105a^3$ be divided by 2x - 3a.

Ans. $24x^2 - 2ax - 35a^2$.

7. Let $4x^4 - 9x^2 + 6x - 1$ be divided by $2x^2 + 3x - 1$.

Ans.
$$2x^2 - 3x + 1$$
.

- 8. Let $x^4 a^2x^2 + 2a^3x a^4$ be divided by $x^2 ax + a^2$.

 Ans. $x^2 + ax a^2$.
- 9. Let $6x^4 96$ be divided by 3x 6, and $a^5 + x^5$ by a + x.

Ans. $2x^3+4x^2+8x+16$, and $a^4-a^3x+a^2x^2-ax^3+x^4$. 10. Let $32x^5+243$ be divided by 2x+3, and x^6-a^6 by

10. Let $52x^5 + 245$ be divided by 2x + 5, and x = 3 by x - a.

Ans. $16x^4 - 24x^3 + 36x^3 - 54x + 81$, and $x^5 + ax^4 + 36x^5 + 6x^4 + 81$.

Ans. $10x^3 - 24x^3 + 30x^3 - 54x + 51$, and $x^5 + 4x^5 + 4x^5$

11. Let $b^4 - 3y^4$ be divided by b - y, and $a^4 + 4a^2b + 3b$ by a + 2b.

Ans.
$$b^3 + b^2 y + by^2 + y^3 - \frac{2y^4}{b-y}$$
, and $a^3 - 2a^2b + 4ab + 4ab^2 - 8b^3 - 8b^3 + \frac{16b^3 + 24b^4}{a+2b}$.

extent the division is supposed to be continued. The real question, however, results from the fractional parts, which (by the divison) is always $+\frac{1}{2}$ when the sum of the terms is 0, and $-\frac{1}{2}$ when the sum is +1: consequently $\frac{1}{2}$ is the true quotient in the former case, and $1-\frac{1}{2}$ in the other. Editor.

12. Let $x^2 + px + q$ be divided by x + a, and $x^3 - px^2 + q$ qx-r by x-a.

Ans.
$$x+p-a+\frac{q-pa+a^2}{x+a}$$
, and $x^2+(a-p)x-ap+a^2+q+\frac{a^3-a^2p+aq-r}{x-a}$.

13. Let $1-5x+10x^2-10x^3+5x^4-x^5$ be divided by

Ans. $1-3x+3x^2-x^3$. $1 - 2x + x^2$.

14. Let $a^4 + 4b^4$ be divided by $a^2 - 2ab + 2b^2$.

Ans. $a^2 + 2ab + 2b^2$.

15. $a^5 - 5a^4x + 10a^3x^2 - 10a^2x^3 + 5ax^4 - x^5$ be divided by $a^2 - 2ax + x^2$. Ans. $a^3 - 3a^2x + 3ax^2 - x^3$.

16. Let $a^4 + b^4$ be divided by $a^2 + ab\sqrt{2} + b_2$.

Ans. $a^2 - ab\sqrt{2} + b^2$.

OF ALGEBRAIC FRACTIONS.

ALGEBRAIC fractions have the same names and rules of operation as numeral fractions in common arithmetic; and the methods of reducing them, in either of these branches, to their most convenient forms, are as follows:

CASE I.

To find the greatest common measure of the terms of a fraction.

BULE.

1. Arrange the two quantities according to the order of their powers, and divide that which is of the highest dimensions by the other, having first expunged any factor, that may be contained in all the terms of the divisor, without being common to those of the dividend.

2. Divide this divisor by the remainder, simplified, if necessary, as before; and so on, for each successive remainder and its preceding divisor, till nothing remains,

when the divisor last used will be the greatest common measure required; and if such a divisor cannot be found, the terms of the fraction have no common measure.*

Note. If any of the divisors, in the course of the operation, become negative, they may have their signs changed, or be taken affirmatively, without altering the truth of the result; and if the first term of a divisor should not be exactly contained in the first term of the dividend, the several terms of the latter may be multiplied by any number, or quantity, that will render the division complete.

EXAMPLES.

1. Required the greatest common measure of the fraction x^4-1 .

$$\begin{array}{c|c}
 & x^{3}-1 \\
 & x^{5}-x^{3} \\
 & x^{4}-1 \\
 & x^{5}-x \\
 & x^{5}-x \\
 & x^{3}+x \\
 & of x^{2}+1 & x^{4}-1 \\
 & x^{4}+x^{2} \\
 & -x^{2}-1 \\
 & -x^{2}-1
\end{array}$$

plain by inspection, that it is not reducible by any simple divisor; but to know whether it may not, by a compound one, I proceed as above, and find the last remainder to be the simple quantity $7x^2$: whence I conclude that the fraction is already in its lowest terms.

† In finding the greatest common measure of two quantities, either of them may be multiplied, or divided, by any quantity, which is not a divisor of the other, or that contains no factor which is common to them both, without in any

respect changing the result.

It may here, also, be farther added, that the common measure, or divisor, of any number of quantities, may be determined in a similar manner to that given above, by first finding the common measure of two of them, and then of that common measure and a third; and so on to the last.

^{*} If, by proceeding in this manner, no compound divisor can be found, that is, if the last remainder be only a simple quantity, we may conclude the case proposed does not admit of any, but is already in its lowest terms. Thus, for instance, if the fraction proposed were to be $\frac{a^3 + 2a^2 + x + 3a^2 + 4x^3}{a^2 + ax + x^2}$; it is plain by inspection, that it is not reducible by any simple divisor; but to

Whence $x^2 + 1$ is the greatest common measure required. 2. Required the greatest common measure of the frac-

tion
$$\frac{x^{3}-b^{2}x}{x^{2}+2bx+b^{2}}.$$

$$\frac{x^{2}+2bx+b^{2}}{x^{2}+2bx+b^{2}}.$$

$$\frac{x^{3}+2bx^{2}+b^{2}x}{x^{3}+2bx^{2}+b^{2}x}$$

$$\frac{x^{3}+2bx^{2}-2b^{2}x}{x^{2}+2bx+b^{2}(x+b)}$$

$$\frac{x^{2}+bx}{bx+b^{2}}$$

Where x+b is the greatest common measure required.

Where
$$x+b$$
 is the greatest common measure required.
3. Required the greatest common measure of the fraction
$$\frac{3a^2-2a-1}{4a^3-2a^2-3a+1}$$

$$3a^2-2a-1)4a^3-2a^2-3a+1$$

$$3$$

$$12a^3-6a^2-9a+3(4a$$

$$12a^3-8a^2-4a$$

$$2a^2-5a+3)3a^2+2a-1$$

$$2$$

$$6a^2-4a-2(3$$

$$6a^2-15a+9$$

11a-11 or a-1

Where, since $a-1)2a^2-5a+3(2a-3)$, it follows that the last divisor a-1 is the common measure required.

^{*} Here, I divide the remainder -2bx2-2b2x by-2xb, (its greatest simple divisor) and the quotient is x+b; and then I divide the last divisor by Editor. x+b, &c.

In which case the common process has been interrupted in the last step, merely to prevent the work overrunning the page.

4. It is required to find the greatest common measure

 $x^{4}-a^{4}$

Ans. x - a.

5. Required the greatest common measure of the frac $a^4 - x^4$

Ans. a2 - x2.

 $a^3 - a^2 x - a x^2 + x^3$ 6. Required the greatest common measure of the fraction $\frac{x^4 + a^2x^2 + a^4}{x^4 + a^2x^2 + a^4}$ Ans. $x^2 + ax + a^3$.

 $x^4 + ax^3 - a^3x - a^4$

7. Required the greatest common measure of the frac- $7a^2 - 23ab + 6b^2$ Ans. a-3b.

tion $\frac{1}{5a^3-18a^2b+11ab^2-6b^3}$ 3. *Required the greatest common measure of the frac-

tion $\frac{x^3 + ax^2 + bx^2 - 2a^2x + bax - 2ba^2}{a^2x + bax - 2ba^2}$ Ans. x+2a. $x^2 - bx + 2ax - 2ab$

* This fraction can be reduced by Simpson's rule (page 50) thus: Fractions that have in them more than two different letters, and one of the letters rises only to a single dimension, either in the numerator or denominator, it will be best to divide the said numerator or denominator (whichever it is) into two parts so that the said letter may be found in every term of the one part, and be totally excluded out of the other: this being done, let the greatest common divisor of these two parts be found, which will evidently be a divisor to the whole, and by which the division of the quantity is to be tried;

as in the following example, where the fraction given is x3 + ax2 + bx2 - 2a2x + bax - 2ba2

x2-bx-1-2ax-2ab

Here the denominator being the least compounded, and b rising therein to a single dimension only, I divide the same into the parts $x^2 + 2ax$, and -bx2ab; which, by inspection, appear to be equal to $(x+2a) \times x$, and (x+2a) \times -b. Therefore x+2a is a divisor to both the parts, and likewise to the whole, expressed by $(x+2a) \times (x-b)$; so that one of these two factors, if the fraction given can be reduced to lower terms, must also measure the numerator: but the former will be found to succeed, the quotient coming out $x^2 - ax + bx - ab$, exactly: whence the fraction itself is reduced to $x^2-ax+bx-ab$, which is not reducible farther by x-b, since the divi-

sion does not terminate without a remainder, as upon trial will be found. This rule is sometimes of great utility, because it spares great labour, and Editor. is very expeditious in reducing several fractions.

- 9. Required the greatest common measure of the fractor $\frac{x^4 3ax^3 8a^2x^2 + 18a^3x 8a^4}{x^3 ax^2 8a^2x + 6a^3}$. Ans. $x^2 + 2ax 2a^2$.
- 10. Required the greatest common measure of the frac- $5a^5 + 10a^4b + 5a^3b^2$ Ans. a+b. $a^3b+2a^2b^2+2ab^3+b^4$
- 11. Required the greatest common measure of the frac- $6a^5 + 15a^4b - 4a^3c^2 - 10a^2bc^2$ Ans. 3a2-2c2. tion $9a^3b-27a^2bc-6abc^2+18bc^3$

CASE II.

To reduce fractions to their lowest or most simple terms.

RULE.

Divide the terms of the fraction by any number, or quantity, that will divide each of them without leaving a remainder; or find their greatest common measure, as in the last rule, by which divide both the numerator and denominator, and it will give the fraction required.

EXAMPLES.

1. Reduce $\frac{a^2bc}{5a^2b^2}$ and $\frac{x^2}{ax+x^2}$ to their lowest terms.

Here
$$\frac{a^2bc}{5a^2b^2} = \frac{c}{5b}$$
 Ans. And $\frac{x^2}{ax+x^2} = \frac{x}{a+x}$ Ans

Here $\frac{a^2bc}{5a^2b^2} = \frac{c}{5b}$ Ans. And $\frac{x^2}{ax+x^2} = \frac{x}{a+x}$ Ans. 2. It is required to reduce $\frac{cx+x^2}{a^2c+a^2x}$ to its lowest terms.

Here
$$cx+x^2$$
 | a^2c+a^2x
or $c+x$ | $a^2c+a^2x(a^2a^2c+a^2x)$

Whence c+x is the greatest common measure; and c+x) $\frac{cx+x^2}{a^2c+a^2x} = \frac{x}{a^2}$ the fraction required. 3. It is required to reduce $\frac{x^2 - b^2x}{x^2 + 2bx + b^2}$ to its lowest

terms.

$$x^2 + 2bx + b^2$$
) $x^3 - b^2x(x$
 $x^3 + 2bx^2 + b^2x$
 $-2bx^2 - 2b^2x$
or $x + b$
 $x^2 + 2bx + b^2(x + b)$
 $x^2 + bx$
 $bx + b^2$
 $bx + b^2$

Whence x+b is the greatest common measure; and x+b? $\frac{x^3-b^2x}{x^2+2bx+b^2} = \frac{x^2-bx}{x+b}$ the fraction required.

And the same answer would have been found, if $x^3 - b^2x$ had been made the divisor instead of $x^2 + 2bx + b^2$.

4. It is required to reduce $\frac{x^4-a^4}{x^5-a^2x^3}$ to its lowest terms.

Ans.
$$\frac{x^2+a^2}{x^3}$$
.

5. It is required to reduce $\frac{6a^2 + 7ax - 3x^2}{6a^2 + 11ax + 3x^2}$ to its lowest

terms. $Ans. \frac{3a-x}{3a+x}$

6. It is required to reduce $\frac{2x^3-16x-6}{3x^3-24x-9}$ to its lowest terms.

7. It is required to reduce $\frac{9x^{5}+2x^{3}+4x^{2}-x+1}{15x^{4}-2x^{3}+10x^{2}-x+2}$ to its lowest terms. $\frac{3x^{3}+x^{2}+1}{5x^{2}+x+2}.$

8. It is required to reduce $\frac{a^2d^2-c^2d^2-a^2c^2+c^4}{4a^2d-4acd-2ac^2+2c^3}$ to

its lowest terms.

Ans. $\frac{ad^2 + cd^2 - ac^2 - c^3}{4ad - 2c^2}.$

CASE III.

To reduce a mixed quantity to an improper fraction.

RULE.

Multiply the integral part by the denominator of the fraction, and to the product add the numerator, when it is affirmative, or subtract it when negative; then the result, placed over the denominator, will give the improper fraction required.

EXAMPLES.

1. Reduce $3\frac{2}{5}$ and $a - \frac{b}{c}$ to improper fractions.

Here
$$3\frac{2}{5} = \frac{3 \times 5 + 2}{5} = \frac{15 + 2}{5} = \frac{17}{5}$$
 Ans.
And $a = \frac{b}{c} = \frac{a \times c}{c} = \frac{b}{c} = \frac{ac - b}{c}$ Ans.

2. Reduce $x + \frac{a}{x}$ and $x - \frac{a^2 - x^2}{x}$ to improper frac-

tions.

Here
$$x + \frac{a}{x} = \frac{x \times x + a}{x} = \frac{x^2 + a}{x}$$
 Ans.
And $x^* = \frac{a^2 - x^2}{x} = \frac{x^2 - a^2 + x^2}{x} = \frac{2x^2 - a^2}{x}$ Ans.

^{*} $x \times x = x^2$. In adding the numerator $a^2 - x^2$, the sign — affixed to the fraction $\frac{a^2 - x^2}{x}$, denotes that the whole of that fraction is to be subtracted, and consequently that the signs of each term of the numerator must be

- 3. Let $1 \frac{2x}{a}$ be reduced to an improper fraction.

 Ans. $\frac{a-2x}{a}$
- 4. Let $5a \frac{3x b}{a}$ be reduced to an improper fraction.

 Ans. $\frac{5a^2 3x + b}{a}$.
- 5. Let $x = \frac{a+x^2}{2a}$ be reduced to an improper fraction.

 Ans. $\frac{2ax-a-x^2}{2a}$
- 6. Let $5 + \frac{2x-7}{3x}$ be reduced to an improper fraction.

 Ans. $\frac{17x-7}{3x}$
- 7. Let $1 \frac{x-a-1}{a}$ be reduced to an improper fraction.

 Ans. $\frac{2a-x+1}{a}$.
- 8. Let $1+2x-\frac{x-3}{5x}$ be reduced to an improper fraction.

 Ans. $\frac{10x^2+4x+3}{5x}$.

CASE IV.

To reduce an improper fraction to a whole or mixed quantity.

RULE.

Divide the numerator by the denominator, for the integral part, and place the remainder, if any, over the denominator, for the fractional part; then the two, joined together, with the proper sign between them, will give the mixed quantity required.

changed when it is combined with x^2 , hence the improper fraction is $\frac{x^2-a^2+x_2}{x}$ or $\frac{2x^2-a^2}{x}$.

EXAMPLES.

1. Reduce $\frac{27}{5}$ and $\frac{ax+a^2}{x}$ to mixed quantities.

Here $\frac{27}{5} = 27 \div 5 = 5\frac{2}{5} Ans$.

And
$$\frac{ax+a^2}{x} = (ax+a^2) \div x = a + \frac{a^2}{x}$$
 Ans.

- 2. It is required to reduce the fraction $\frac{ax-x^3}{x}$ to a whole quantity.

 Ans. $a=x^2$.
- 3. It is required to reduce the fraction $\frac{ab-2a^2}{ab}$ to a mixed quantity.

 Ans. $1-\frac{2a}{b}$.
- 4. It is required to reduce the fraction $\frac{a^2 + x^2}{a x}$ to a mixed quantity.

 Ans. $a + x + \frac{2x^2}{a x}$.
- 5. It is required to reduce the fraction $\frac{x^3-y^3}{x-y}$ to a vhole quantity.

 Ans. x^2+xy+y^2 .
- 6. It is required to reduce the fraction $\frac{10x^2 5x + 3}{5x}$ o a mixed quantity. Ans. $2x 1 + \frac{3}{5x}$

CASE V.

Po reduce fractions to other equivalent ones, that shall have a common denominator.

BULE.

Multiply each of the numerators, separately, into all the enominators, except its own, for the new numerators, and

all the denominators together for a common denominator*.

EXAMPLES.

1. Reduce $\frac{a}{b}$ and $\frac{b}{c}$ to fractions that shall have a common denominator.

Here $a \times c = ac$ $b \times b = b^2$ the new numerators.

 $b \times c = bc$ the common denominator.

Whence, $\frac{a}{b}$ and $\frac{b}{c} = \frac{ac}{bc}$ and $\frac{b^2}{bc}$, the fractions required.

- 2. Reduce $\frac{2x}{a}$ and $\frac{b}{c}$ to equivalent fractions having a common denominator.

 Ans. $\frac{2cx}{ac}$ and $\frac{ab}{ac}$
- 3. Reduce $\frac{a}{b}$ and $\frac{a+b}{c}$ to equivalent fractions having a common denominator.

 Ans. $\frac{ac^-}{bc}$ and $\frac{ab+b^2}{bc}$.
- 4. Reduce $\frac{3x}{2a}$, $\frac{2b}{3c}$, and d, to equivalent fractions having a common denominator. $\frac{9cx}{6ac}$, $\frac{4ab}{6ac}$ and $\frac{6acd}{6ac}$.
- 5. Reduce $\frac{3}{4}$, $\frac{2x}{3}$ and $a + \frac{4x}{5}$, to fractions having a common denominator.

 Ans. $\frac{45}{60}$, $\frac{40x}{60}$ and $\frac{60a + 48x}{60}$.

which method is often of great use in reducing fractions more readily to a common denominator.

^{*} It may here be remarked, that if the numerator and denominator of a fraction be either both multiplied, or both divided, by the same number or quantity, its value will not be altered: thus

6. Reduce $\frac{a}{2}$, $\frac{3x}{7}$, and $\frac{a+x}{a-x}$, to fractions having a common denominator.

Ans. $\frac{7a^2-7ax}{14a-14x}$, $\frac{6ax-6x^2}{14a-14x}$ & $\frac{14a+14x^4}{14a-14x}$

CASE VI.

To add fractional quantities together.

RULE.

Reduce the fractions, if necessary, to a common denominator; then add all the numerators together, and under their sum put the common denominator, and it will give the sum of the fractions required*.

EXAMPLES.

1. It is required to find the sum of $\frac{x}{x}$ and $\frac{x}{3}$.

Here
$$\begin{bmatrix} x \times 3 = 3x \\ x \times 2 = 2x \end{bmatrix}$$
 the numerators.

And 2×3=6 the common denominator.

Whence $\frac{3x}{6} + \frac{2x}{6} = \frac{5x}{6}$, the sum required.

2. It is required to find the sum of $\frac{a}{b}$, $\frac{c}{d}$, and $\frac{e}{f}$.

Here
$$a \times d \times f = adf$$
 $c \times b \times f = cbf$
 $e \times b \times d = ebd$ the numerators.

^{*} In the adding or subtracting of mixed quantities, it is best to bring the fractional parts only to a common denominator, and then to affix their sum or difference to the sum or difference of the integral parts, interposing the proper sign.

And $b \times d \times f = bdf$ the common denominator. Whence $\frac{adf}{bdf} + \frac{cbf}{bdf} + \frac{ebd}{bdf} = \frac{adf + cbf + ebd}{bdf}$ the sum.

3. It is required to find the sum of $a - \frac{3x^2}{b}$ and $b + \frac{2ax}{c}$.

Here, taking only the fractional parts, We shall have $\begin{cases} 3x^2 \times c = 3cx^2 \\ 2ax \times b = 2abx \end{cases}$ the numerators.

And $b \times c = bc$ the common denominator. Whence $a = \frac{3cx^2}{bc} + b + \frac{2abx}{bc} = a + b + \frac{2abx - 3cx^2}{bc}$ the sum.

4. It is required to find the sum of $\frac{2x}{5}$ and $\frac{5x}{7}$.

Ans. $\frac{39x}{35}$.

5. It is required to find the sum of $\frac{3x}{2a}$ and $\frac{x}{5}$.

Ans. $\frac{15x+2ax}{10a}$.

6. It is required to find the sum of $\frac{x}{2}$, $\frac{x}{3}$, and $\frac{x}{4}$.

Ans. $\frac{13x}{12}$.

7. It is required to find the sum of $\frac{4x}{7}$ and $\frac{x-2}{5}$.

7. It is required to find the sum of $\frac{\pi}{7}$ and $\frac{5}{5}$.

Ans. $\frac{27x - 14}{35}$

8. Required the sum of 2a, $3a + \frac{2x}{5}$, and $a = \frac{8x}{9}$.

Ans. $6a - \frac{22x}{45}$. 9. Required the sum of $2a + \frac{3x}{5}$, $\frac{a}{a-x}$, and $\frac{a-x}{a}$.

Ans. $2a+2+\frac{3a^2x-3ax^2+5x^2}{5a^2-5ax}$

10. Required the sum of
$$5x + \frac{x-2}{3}$$
 and $4x - \frac{2x-3}{5x}$.

Ans. $9x + \frac{5x^2 - 16x + 9}{15x}$.

11. It is required to find the sum of 5x, $\frac{2a}{3x^2}$, and $\frac{a+2x}{4x}$.

Ans. $5x + \frac{8a+3ax+6x^2}{12x^2}$.

CASE VII.

To subtract one fractional quantity from another.

RULE.

Reduce the fractions to a common denominator, if necessary, as in addition; then subtract the less numerator from the greater, and under the difference write the common denominator, and it will give the difference of the fractions required.

EXAMPLES.

1. It is required to find the difference of $\frac{2x}{3}$ and $\frac{3x}{5}$. Here $\frac{2x \times 5 = 10x}{3x \times 3 = 9x}$ the numerators.

And $3 \times 5 = 15$ the common denominator.

Whence $\frac{10x}{15} - \frac{9x}{15} = \frac{x}{15}$, the difference required.

2. It is required to find the difference of $\frac{x-a}{2b}$ and $\frac{2a-4x}{3c}$.

Here
$$(2a-4x) \times 3c = 3cx - 3ac$$
 $\{2a-4x) \times 2b = 4ab - 8bx \}$ the numerators

And $2b \times 3c = 6bc$ the common denominator. 3cx-3ac 4ab-8bx 3cx-3ac-4ab+8bx

the difference required.

3. Required the difference of $\frac{12x}{7}$ and $\frac{3x}{5}$. Ans. $x + \frac{4x}{35}$.

4. Required the difference of 15y and $\frac{1+2y}{c}$.

Ans. $\frac{118y-1}{2}$.

5. Required the difference of $\frac{dx}{h-c}$ and $\frac{dx}{h+c}$.

Ans. $\frac{2axc}{b^2-a^2}$.

6. Required the difference of $x - \frac{x-a}{c}$ and $x + \frac{x}{2h}$.

7. Required the difference of $a + \frac{a-x}{a+x}$ and $a - \frac{a+x}{a-x}$ Ans. $\frac{2a^2+2x^2}{a^2-x^2}$

3. Required the difference of $ax + \frac{2x+7}{9}$ and $x - \frac{2x+7}{9}$ 5x - 6Ans. $ax - \frac{86x - 99}{168}$ 21

9. Required the difference of $2x + \frac{3x - 5}{7}$, and $3x + \frac{3x - 5}{7}$ 11x - 10Ans. $x + \frac{32x+5}{105}$. 15

10. Required the difference of $a + \frac{a-x}{a(a+x)}$ and Ans. $a = \frac{4x}{a^2 - x^2}$

CASE VIII.

To multiply fractional quantities together.

RULE.

Multiply the numerators together for a new numerator, and the denominators for a new denominator; and the former of these, being placed over the latter, will give the product of the fractions, as required *.

EXAMPLES.

1. It is required to find the product of $\frac{x}{6}$ and $\frac{2x}{9}$.

Here
$$\frac{x \times 2x}{6 \times 9} = \frac{2x^2}{54} = \frac{x^2}{27}$$
 the product required.

2. It is required to find the continued product of $\frac{x}{2}$, $\frac{4x}{5}$, and $\frac{10x}{21}$.

Here
$$\frac{x \times 4x \times 10x}{2 \times 5 \times 21} = \frac{40x^3}{210} = \frac{4x^2}{21}$$
 the product.

3. It is required to find the product of $\frac{x}{a}$ and $\frac{a+x}{a-x}$.

Here
$$\frac{x \times (a+x)}{a \times (a-x)} = \frac{x^2 + ax}{a^2 - ax}$$
 the product.

4. It is required to find the product of $\frac{3x}{2}$ and $\frac{5x}{3b}$.

Ans. $\frac{5x^2}{2}$.

When the numerator of one of the fractions to be multiplied, and the denominator of the other, can be divided by some quantity which is common to each of them, the quotients may be used instead of the fractions themselves.

Also, when a fraction is to be multiplied by an integer, it is the same thing whether the numerator be multiplied by it, or the denominator divided by it. Or if an integer is to be multiplied by a fraction, or a fraction by an integer, the integer may be considered as having unity for its denominator, and the two be then multiplied together as usual.

5. It is required to find the product of $\frac{2x}{5}$ and $\frac{3x_2}{2a}$.

Ans. $\frac{3x^2}{5a}$.

6. It is required to find the continued product of $\frac{2x}{3}$, $\frac{4x^2}{7}$, and $\frac{a}{a+x}$.

Ans. $\frac{8ax^3}{21a+21x}$.

7. It is required to find the continued product of $\frac{2x}{a}$, $\frac{3ab}{c}$, and $\frac{5ac}{2b}$.

Ans. 15ax.

8. It is required to find the product of $2a + \frac{bx}{a}$ and $3a - \frac{b}{ax}$.

Ans. $6a^2 + 3bx - \frac{2b}{x} - \frac{b^2}{a}$.

9. It is required to find the continued product of 3x, $\frac{x+1}{2a}$, and $\frac{x-1}{a+b}$.

Ans. $\frac{3x^3-3x}{2a^2+2ab}$.

10 It is required to find the continued product of $\frac{a^2-x^2}{a+b}$, $\frac{a^2-b^2}{ax+x^2}$, and $a+\frac{ax}{a-x}$.

Ans. $\frac{a^3-a^2b}{x}$.

CASE IX.

To divide one fractional quantity by another.

RULE.

Multiply the denominator of the divisor by the numerator of the dividend, for the numerator; and the numerator of the divisor by the denominator of the dividend, for the denominator. Or, which is more convenient in practice, multiply the dividend by the reciprocal of the divisor, and the product will be the quotient required*.

When a fraction is to be divided by an integer, it is the same thing whether the numerator be divided by it, or the denominator multiplied by it.

EXAMPLES.

1. It is required to divide $\frac{x}{3}$ by $\frac{2x}{9}$.

Here
$$\frac{x}{3} \div \frac{2x}{9} = \frac{x}{3} \times \frac{9}{2x} = \frac{9x}{6x} = \frac{3}{2} = 1\frac{1}{2}$$
 .Ans.

2. It is required to divide $\frac{2a}{b}$ by $\frac{4c}{d}$.

Here
$$\frac{2a}{b} \times \frac{d}{4c} = \frac{2ad}{4bc} = \frac{ad}{2bc}$$
 Ans.

3. It is required to divide $\frac{x+a}{x-b}$ by $\frac{x+b}{5x+a}$

Here
$$\frac{x+a}{x-b} \times \frac{5x+a}{x+b} = \frac{5x^2 + 6ax + a^2}{x^2 - b^2}$$
 Ans.

4. It is required to divide
$$\frac{2x^2}{a^3+x^3}$$
 by $\frac{x}{x+a}$.
Here $\frac{2x^2}{a^3+x^3} \times \frac{x+a}{x} = \frac{2x^2(x+a)}{x(a^3+x^3)} = \frac{2x}{x^2-ax+a^2}$.

5. It is required to divide $\frac{7x}{5}$ by $\frac{3}{x}$. Ans. $\frac{7x^2}{15}$.

6. It is required to divide
$$\frac{4x^2}{7}$$
 by ξx . Ans. $\frac{4x}{35}$.

7. It is required to divide $\frac{x+1}{6}$ by $\frac{2x}{3}$. Ans. $\frac{x+1}{4x}$.

3. It is required to divide
$$\frac{x}{1-x}$$
 by $\frac{x}{5}$. Ans. $\frac{5}{1-x}$.

9. It is required to divide $\frac{2ax+x^2}{c^3-x^3}$ by $\frac{x}{c-x}$.

Ans.
$$\frac{2a+x}{c^2+cx+x^2}$$

9. It is required to give $c \circ -x$ $3ns. \frac{2a+x}{c^2+cx+x^2}$ 10. It is required to divide $\frac{x^4-b^4}{x^2-2bx+b^2} \text{ by } \frac{x^2+bx}{x-b}.$ $3ns. \frac{2a+x}{x^2+bx}$ $3ns. \frac{2a+x}{x^2+bx}$ $3ns. \frac{x^2+b^2}{x^2-bx}$

Ans
$$\frac{x^2+b^2}{x}$$

Also, when the two numerators or the two denominators, can be divided by some common quantity, that quantity may be thrown out of each, and the quotients used instead of the fractions first proposed.

INVOLUTION.

Involution is the raising of powers from any proposed root; or the method of finding the square, cube, biquadrate, &c. of any given quantity.

BULE I.

Multiply the index of the quantity by the index of the power to which it is to be raised, and the result will be the power required.

Or multiply the quantity into itself as many times less one as is denoted by the index of the power, and the last

product will be the answer.

Note. When the sign of the root is +, all the powers of it will be +; and when the sign is -, all the even powers will be +, and the odd powers -: as is evident from multiplication*.

EXAMPLES.

a, the root.	1a2 the root.
$a^2 = square.$	a4 = square.
$a^3 = \text{cube}$.	a6=cube.
$a^4 = 4$ th power.	a = 4th power.
$a^5 = 5$ th power.	$a^{10} = 5$ th power.
&c.	&c.
-3a the root.	-2ax2 the root.
$+9a^2 = square.$	$+4a^2x^4$ = square.
$-27a^3$ = cube.	$-8a^3x^6$ = cube.
$+81a^4=4$ th power.	$+16a^4x^3 = 4$ th power.
&c.	&c.

^{*} Any power of the product of two or more quantities is equal to the same power of each of the factors multiplied together. And any power of a fraction is equal to the same power of the numerator divided by the like power of the denominator.

Also, am raised to the nth power is amn; and -- am raised to the nth

power is + amn, according as n is an even or an odd number.

$\frac{x}{a}$ the root. $\frac{x^2}{a^2}$ = square. $\frac{x^3}{a^3}$ = cube. $\frac{x^4}{a^4}$ = 4th power.	$ \frac{-\frac{2ax^{2}}{3b} \text{ the root.}}{+\frac{4a^{2}x^{4}}{9b^{2}} = \text{square.}} $ $ \frac{8a^{3}x^{6}}{27b^{3}} = \text{cube.} $ $ \frac{16a^{4}x^{8}}{81b^{4}} = 4\text{th power.} $ &c.
-a the root.	x+a the root.
$-ax^{2}$ $-ax+a^{2}$	$\begin{array}{c} x^2 + ax \\ + ax + a^2 \end{array}$
$-2ax + a^2$ square.	$\begin{array}{c} x^2 + 2ax + a^2 \text{ square.} \\ x + a \end{array}$
$ \begin{array}{c} -2ax^2 + a^2x \\ -ax^2 + 2a^2x - a^3 \end{array} $	$ \begin{array}{r} x^3 + 2ax^2 + a^2x \\ +ax^2 + 2a^2x + a^3 \end{array} $

x2 -

2:3 .

$x^3 - 3ax^2 + 3a^2x - a^3$ cube. $x^3 + 3ax^2 + 3a^2x + a^3$ cube. EXAMPLES FOR PRACTICE.

- Required the cube or third power, of 2a².
 Ans. 8a⁶.
- 2. Required the biquadrate, or 4th power, of $2a^2x$.
- Ans. $16a^8x^4$.

 3. Required the cube, or third power, of $-\frac{2}{3}x^2y^3$.
 - $Ans. \frac{8}{27}x^6y^3.$
- 4. Required the biquadrate, or 4th power of $\frac{3a^2x}{5b^2}$.

 Ans. $\frac{81a^3x^6}{625b^2}$

5. Required the 4th power of a+x; and the 5th power of a-y.

Ans. $a^3+4a^3x+6a^2x^2+4ax^3+x^4$, and $a^5-5a^4y+10a^3y^2-10a^2y^3+5ay^4-y^5$.

RULE II.

A binomial or residual quantity may also be readily rais-

ed to any power whatever, as follows:

1. Find the terms without the coefficients, by observing that the index of the first, or leading quantity, begins with that of the given power, and decreases continually by 1, in every term to the last; and that in the following quantity, the indices of the terms are 1, 2, 3, 4, &c.

2. To find the coefficients, observe that those of the first and last terms are always 1; and that the coefficient of the second term is the index of the power of the first: and for the rest, if the coefficient of any term be multiplied by the index of the leading quantity in it, and the product be divided by the number of terms to that place, it will give the coefficient of the term next following.

Note. The whole number of terms will be one more than the index of the given power; and when both terms of the root are +, all the terms of the power will be +; but if the second term be -, all the odd terms will be +, and the even terms -; or, which is the same thing, the

terms will be + and - alternately*.

$$(a+b)^m = a^m + ma_{m-1}b + m \cdot \frac{m-1}{2}a^{m-2}b^2 + m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3}a_{m-3}b^3$$
, c

$$(a-b)^m = a^m - ma^{m-1}b + m \cdot \frac{m-1}{2}a^{m-2}b^2 - m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3}a_{m-3}b_3$$
, &c.

which formulæ will, also, equally hold when m is a fraction, as will be

more fully explained hereafter.

^{*} The rule here given, which is the same in the case of integral powers as the binominal theorem of Newton, may be expressed in general terms, as follows;

It may, also, be farther observed, that the sum of the coefficients in every power, is equal to the number 2 raised to that power. Thus 1+3=2, for the first power; 1+2+1=4=22, for the square; 1+3+3+1=3=323, for the cube, or third power; and so on.

EXAMPLES.

1. Let a+x be involved, or raised to the 5th power. Here the terms, without the coefficients, are

 a^5 , a^4x , a^3x^2 , a^2x^3 , ax^4 , x^5 .

And the coefficients, according to the rule, will be

$$1, 5, \frac{5 \times 4}{2}, \frac{10 \times 3}{3}, \frac{10 \times 2}{4}, \frac{5 \times 1}{1};$$

or 1, 5, 10, 10, 5, 1, Whence the entire 5th power of a+x is $a^5+5a^4x+10a^3x^2+10a^2x^3+5ax^4+x^5$.

2. Let a—x be involved, or raised, to the 6th power. Here the terms, without their coefficients, are
a⁵, a⁵x, a⁴x², a³x³, a²x⁴, ax⁵, x⁵.

And the coefficients, found as before, are

1, 6,
$$\frac{6 \times 5}{2}$$
, $\frac{15 \times 4}{3}$, $\frac{20 \times 3}{4}$, $\frac{15 \times 2}{5}$, $\frac{6 \times 1}{6}$;

or 1, 6, 15, 20, 15, 6, 1.

Whence the entire 6th power of a-x is $a^6-6a^5x+15a^4x^2-20a^3x^3+15a^2x^6-6ax^5+x^6$.

- 3. Required the 4th power of a+x, and the 5th power of a-x. Ans. $a^4+4a^3x+6a^2x^2+4ax^3+x^4$, and a^5-
- 4. Required the 6th power of a+b, and the 7th power of a-y.

 Ans. $a^6+6a^5b+15a^4b^2+20a^3b^3+15a^2b^4+6ab^5+b^6$, and $a^7-7a^6y+21a^5y^2-35$. $a^4y^3+35a^3y^4-21a^2y^5+7ay^6-y^7$.

5. Required the 5th power of 2+x, and the cube of a-bx+c. Ans. $32+80x+80x^2+40x^3+10x^3+x^2$, and $a^3+3a^2c+3ac^2+c^2-3a^2bx-6acbx-3c^2bx+8ab^2x^2+3cb^2x^2-b^3x^3$.

EVOLUTION.

Evolution, or the extraction of roots, is the reverse of involution, or the raising powers; being the method of finding the square root, cube root, &c. of any given quantity.

CASE I.

To find any root of a simple quantity.

RULE.

Extract the root of the coefficient for the numeral part, and the root of the quantity subjoined to it for the literal part; then these, joined together, will be the root required.

And if the quantity proposed be a fraction, its root will be found, by taking the root both of its numerator and de-

nominator.

Note. The square root, the fourth root, or any other even root, of an affirmative quantity, may be either + or -. Thus, $\sqrt{a^2 + a}$ or -a, and $\sqrt[4]{b^2 + b}$ or -b, &c. But the cube root, or any other odd root, of a quantity, will have the same sign as the quantity itself. Thus,

 $\sqrt[3]{a^3} = a$; $\sqrt[3]{-a^3} = -a$; and $\sqrt[5]{-a^5} = -a$, &c.*

It may here, also, be farther remarked, that any even root of a negative quantity, is unassignable.

Thus, $\sqrt{-a^2}$ cannot be determined, as there is no quantity, either positive or negative, (+ or —), that, when multiplied by itself, will produce $-a^2$.

EXAMPLES.

1. Find the square root of $9x^2$; and the cube root of $8x^3$.

Here $\sqrt{9x^2} = \sqrt{9} \times \sqrt{x^2} = 3 \times x = 3x$ Ans. And $\sqrt[3]{3x^3} = \sqrt[3]{8} \times \sqrt[3]{x^3} = 2 \times x = 2x$. Ans.

And for the cube root, fifth root, &c. of a negative quantity, it is plain, from the same rule, that

the

 $(-a) \times (-a) \times (-a) = -a^3$; and $(-a^3) \times (+a^2) = -a^5$, And consequently $\sqrt[3]{-a^5} = -a$, and $\sqrt[5]{-a^5} = -a$.

^{*} The reason why + a and - a are each the square root of a_2 is obvious, since, by the rule of multiplication, $(+a) \times (+a)$ and $(-a) \times (-a)$ are both equal to a^2 .

2. It is required to find the square root of $\frac{a^2x^2}{4c^2}$ and $8a^3x^3$

the cube root of $-\frac{8a^3x^3}{27c^3}$.

Here $\sqrt{\frac{a^2x^2}{4c^2}} = \frac{\sqrt{a^2x^2}}{\sqrt{4c^2}} = \frac{ax}{2c}$; and $\sqrt[3]{-\frac{8a^3x^3}{27c^3}} = -\frac{2ax}{3c}$. 3. It is required to find the square root of $4a^2x^6$.

- 3. It is required to find the square root of $4a^2x^5$.

 Ans. $2ax^3$.
- 4. It is required to find the cube root of $-125a^3x^6$.

 Ans. $-5ax^2$.
- 5. It is required to find the 4th root of $256a^3x^2$.

 Ans. $4ax^2$.
- 6. It is required to find the square root of $\frac{4a^4}{9x^2y^2}$.

Ans. $\frac{za^2}{3xy}$.

7. It is required to find the cube root of $\frac{8a^3}{125x^6}$.

Ans. $\frac{2a}{5x^2}$.

8. It is required to find the 5th root of $-\frac{32a^5x^{10}}{243}$:

 $.9ns. \frac{-2ax^2}{3}.$

CASE II.

To extract the square root of a compound quantity.

RULE.

1. Range the terms, of which the quantity is composed, according to the dimensions of some letter in them, beginning with the highest, and set the root of the first term in the quotient.

Subtract the square of the root, thus found from the first term, and bring down the two next terms to the remainder for a dividend. 3. Divide the dividend, thus found, by double that part of the root already determined, and set the result both in

the quotient and divisor.

4. Multiply the divisor, so increased, by the term of the root last placed in the quotient, and subtract the product from the dividend; and so on, as in common arithmetic.

EXAMPLES.

1. Extract the square root of $x^4 - 4x^3 + 6x^2 - 4x + 1$: $x^4 - 4x^3 + 6x^2 - 4x + 1(x^2 - 2x + 1)$

$$2x^{2} - 2x) - 4x^{3} + 6x^{2}$$

$$-4x^{3} + 4x^{2}$$

$$2x^{2} - 4x + 1)2x^{2} - 4x + 1$$

$$2x^{2} - 4x + 1$$

Ans. $x^2 - 2x + 1$, the root required.

2. Extract the square root of $4a^4 + 12a^3x + 13a^2x^2 + 6ax^3 + x^4$.

$$4a^{4} + 12a^{3}x + 13a^{2}x^{2} + 6ax^{3} + x^{4}(2a^{2} + 3ax + x^{2} + 4a^{4})$$

$$4a^{2} + 3ax(12a^{3}x + 13a^{2}x^{2} \\ 12a^{3}x + 9a^{2}x^{2}$$

$$4a^{2} + 6ax + x^{2})4a^{2}x^{2} + 6ax^{3} + x^{4}$$

$$4a^{2}x^{2} + 6ax^{3} + x^{4}$$

Note. When the quantity to be extracted has no exact root, the operation may be carried on as far as is thought necessary, or till the regularity of the terms shows the law by which the series would be continued.

EXAMPLE.

1. It is required to extract the square root of 1+x.

$$1+x(1+\frac{x}{2}-\frac{x^2}{8}+\frac{x^3}{16}-\frac{5x^4}{128}, &c.$$

$$\frac{1}{2+\frac{x}{2}})_x$$

$$x+\frac{x^2}{4}$$

$$2+x-\frac{x^2}{8})-\frac{x^2}{4}$$

$$-\frac{x^2}{4}-\frac{x^3}{8}+\frac{x^4}{64}$$

$$2+x-\frac{x^2}{4}+\frac{x^3}{16})\frac{x^3}{8}-\frac{x^4}{64}$$

$$\frac{x^3}{8}+\frac{x^4}{16}-\frac{x^5}{64}+\frac{x^5}{256}$$

$$-\frac{5x^4}{64}+\frac{x^5}{64}-\frac{x^6}{256}$$

Here, if the numerators and denominators of the two last terms be each multiplied by 3, which will not alter their values, the root will become

$$1 + \frac{x}{2} - \frac{x^2}{2.4} + \frac{3x^3}{2.4.6} - \frac{3.5x^4}{2.4.6.8} + \frac{3.5.7x^5}{2.4.6.8.10}$$
, &c. where the law of the series is manifest.

EXAMPLES FOR PRACTICE.

2. It is required to find the square root of $a^4 + 4a^2x +$ $6a^2x^2 + 4ax^3 + x^4$. Ans. $a^2 + 2ax + x^2$.

3. It is required to find the square root of $x^4 - 2x^3 +$ $\frac{3}{9}x^3 - \frac{1}{2}x + \frac{1}{16}$ Ans. $x^2 - x + 1$.

4. It is required to find the square root of $4x^6 - 4x^4 +$ $12x^3 + x^2 - 6x + 9$. $-Ans. 2x^3 - x + 3.$

5. Required the square root of $x^6 + 4x^5 + 10x^4 + 20x^3$ Ans. $x^3 + 2x^2 + 3x + 4$. $+25x^2+24x+16$.

6. It is required to extract the square root of $a^2 + b$.

Ans. $a + \frac{b}{2a} - \frac{b^2}{8a^3} + \frac{b^3}{16a^5} - \frac{6b^4}{128a^7}$, &c.

7. It is required to extract the square root of 2, or of Ans. $1+\frac{1}{2}-\frac{1}{2}+\frac{1}{16}-\frac{1}{22}+\frac{1}{64}$, &c. 1+1.

CASE III.

To find any root of a compound quantity.

RULE.

Find the root of the first term, which place in the quotient; and having subtracted its corresponding power from that term, bring down the second term for a dividend.

Divide this by twice the part of the root above determined, for the square root; by three times the square of it, for the cube root, and so on; and the quetient will be the next term of the root.

Involve the whole of the root, thus found, to its proper power, which subtract from the given quantity, and divide the first term of the remainder by the same divisor as before; and proceed in this manner till the whole is finished*.

* As this rule, in high powers, is often found to be very laborious, it may be proper to observe, that the roots of various compound quantities

may sometimes be easily discovered, as follows:

Extract the roots of all the simple terms, and connect them together by the signs + or -, as may be judged most suitable for the purpose ; then involve the compound root, thus found, to its proper power, and if it be the same with the given quantity, it is the root required. But if it befound to differ only in some of the signs, change them from + to -, or from - to +, till its power agrees with the given one throughout.

Thus, in the third example next following, the root is 2a - 3x, which is the difference of the roots of the first, and last terms; and in the fourth example, the root is a+b+c, which is the sum of the roots of the first, fourth, and sixth terms. The same may also be observed of the sixth example,

where the root is found from the first and last terms.

EXAMPLES.

1. Required the square root of $a^4 - 2a^3x + 3a^2x^2 - 2ax^3 + x^4$.

$$a^{4} - 2a^{3}x + 3a^{2}x^{2} - 2ax^{3} + x^{4}(a^{2} - ax + x^{2})$$

$$a^{4} - 2a^{3}x$$

$$a^{4} - 2a^{3}x + a^{2}x^{2}$$

$$2a^{2})2a^{2}x^{3}$$

$$a^{4} - 2a^{3}x + 3a^{2}x^{2} - 2ax^{3} + x^{4}$$

2. Required the cube root of $x^6 + 6x^5 - 40x^3 + 96x - 64$.

$$\begin{array}{r}
 x^{6} + 6x^{5} - 40x^{3} + 96x - 64(x^{2} + 2x - 4) \\
 x^{6} \\
 \hline
 3x^{4})6x^{5} \\
 \hline
 x^{6} + 6x^{5} + 12x^{4} + 8x^{3} \\
 \hline
 3x^{4}) - 12x^{4} \\
 \hline
 x^{6} + 6x^{5} - 40x^{3} + 96x - 64
 \end{array}$$

3. Required the square root of $4a^2 - 12ax + 9x^2$.

Ans 2a-3x. 4. Required the square root of $a^2+2ab+2ac+3^2+2bc+c^2$. Ans. a+b+c.

5. Required the cube root of $x^6 - 6x^5 + 15x^4 - 20x^3 + 15x^2 - 6x + 1$.

Ans x - 1.

6. Required the 4th root of $16a^4 - 96a^3x + {}^{\circ}16x^2x^3 - 216ax^3 + 81x^4$. Ans. 2a - 3x.

54 IRRATIONAL QUANTITIES, OR SURDS.

7. Required the 5th root of $32x^5-30x^3+80x^3-40x^2+10x-1$. Ans. 2x-1.

OF IRRATIONAL QUANTITIES, or SURDS.

IRRATIONAL quantities, or surds, are such as have no exact root, being usually expressed by means of the radical sign, or by fractional indices; in which latter case, the numerator shows the power the quantity is to be raised to, and the denominator its root.

Thus, $\sqrt{2}$, or $2^{\frac{1}{2}}$, denotes the square root of 2; and $\sqrt[3]{a^2}$, or $a^{\frac{2}{3}}$, is the square of the cube root of a, &c. *

CASE I.

To reduce a rational quantity to the form of a surd.

RULE.

Raise the quantity to a power corresponding with that denoted by the index of the surd; and over this new quantity place the radical sign, or proper index, and it will be of the form required.

EXAMPLES.

Let 3 be reduced to the form of the square root.
 Here 3×3=3²=9; whence

9 Ans.

^{*} A quantity of the kind here mentioned, as for instance \(\sqrt{2} \), is called an irrational number, or a surd, because no number, either whole or fractional, can be found, which when multiplied by itself, will produce 2. But its approximate value may be determined to any degree of exactness, by the common rule for extracting the square root, being 1 and certain non periodic detignals, which giver terminate.

2. Reduce 2x2 to the form of the cube root.

Here $(2x^2)^3 = 8x^6$; whence $\sqrt[3]{8x^6}$, or $(8x^6)^{\frac{1}{3}}$.

3. Let 5 be reduced to the form of the square root.

Ans. 1/(25)

4. Let-3x be reduced to the form of the cube root. Ans. $\frac{3}{4} - (27x^3)$.

5. Let -2a be reduced to the form of the fourth root. Ans. - 4/ (16 a4).

6. Let a2 be reduced to the form of the fifth root, and $\sqrt{a} + \sqrt{b}$, $\frac{\sqrt{a}}{2a}$ and $\frac{a}{b\sqrt{a}}$ to the form of the square root.

Ans. $\sqrt[5]{a^{10}}$, $\sqrt{(a+2\sqrt{ab+b})}$, $\sqrt{(\frac{1}{4}a)}$, and $\sqrt{\frac{a}{b^2}}$.

Note Any rational quantity may be reduced by the above rule, to the form of the surd to which it is joined, and their product be then placed under the same index, or radical sign.

EXAMPLES.

Thus $2\sqrt{2}=\sqrt{4}\times\sqrt{2}=\sqrt{4}\times2=\sqrt{8}$

And 23/4=3/8×3/4=3/8×4=3/32

Also $3\sqrt{a} = \sqrt{9} \times \sqrt{a} = \sqrt{9} \times a = \sqrt{9}a$

And $\frac{1}{3}\sqrt{4a} = \frac{3}{1} \times \frac{3}{4} \times \frac{3}{4a} = \frac{3}{1} \times 4a = \frac{3}{1} \times \frac{a}{1} \times \frac{a}$

1. Let 5/6 be reduced to a simple radical form.

Ans. (150).

2. Let \frac{1}{5} \sqrt{5a} be reduced to a simple radical form.

Ans $\sqrt{\left(\frac{a}{5}\right)}$

3. Let $\frac{2a}{3}\sqrt[3]{\frac{9}{4a^2}}$ be reduced to a simple radical form.

Ans. $\sqrt[3]{\frac{2a}{2}}$.

CASE II.

To reduce quantities of different indices, to others that shall have a given index.

RULE.

Divide the indices of the proposed quantities by the

given index, and the quotients will be the new indices for those quantities.

Then, over the said quantities, with their new indices, place the given index, and they will be the equivalent quantities required.

EXAMPLES.

1. Reduce $3^{\frac{1}{2}}$ and $2^{\frac{1}{3}}$ to quantities that shall have the index $\frac{1}{6}$.

Here
$$\frac{1}{2} \div \frac{1}{6} = \frac{1}{2} \times \frac{6}{1} = \frac{6}{2} = 3$$
, the 1st index:
And $\frac{1}{3} \div \frac{1}{6} = \frac{1}{3} \times \frac{6}{1} = \frac{6}{3} = 2$, the 2d index.

Whence $(3^3)^{\frac{1}{6}}$ and $(2^2)^{\frac{1}{6}}$, or $27^{\frac{1}{6}}$ and $4^{\frac{1}{6}}$, are the quantities required.

- 2. Reduce $5^{\frac{1}{3}}$ and $6^{\frac{1}{3}}$ to quantities that shall have the common index $\frac{1}{6}$.

 Ans. $12^{\frac{1}{5^{\circ}}}$ and $36^{\frac{1}{6^{\circ}}}$.
- 3. Reduce $2^{\frac{1}{2}}$ and $4^{\frac{7}{4}}$ to quantities that shall have the common index $\frac{1}{8}$ Ans. $16^{\frac{1}{8}}$ and $16^{\frac{1}{4}}$.
- 4. Reduce a^2 and $a^{\frac{1}{2}}$ to quantities that shall have the common index $\frac{1}{4}$ Ans. Ans.
- 5. Reduce $a^{\frac{1}{2}}$ and $b^{\frac{2}{3}}$ to quantities that shall have the common index $\frac{1}{6}$.

 Ans. $(a^3)^{\frac{1}{6}}$ and $(b^4)^{\frac{1}{6}}$.

Note. Surds may also be brought to a common index, by reducing the indices of the quantities to a common denominator, and then involving each of them to the power denoted by its numerator.

EXAMPLES.

1. Reduce $3^{\frac{1}{2}}$ and $4^{\frac{1}{3}}$ to quantities having a common index.

Here
$$3^{\frac{1}{2}} = 3^{\frac{2}{6}} = \overline{3^3}|_{6}^{6} = \overline{21}|_{7}^{7}$$

And $4^{\frac{1}{3}} = 4^{\frac{2}{6}} = 4^{\frac{2}{3}}|_{7}^{7} = \overline{16}|_{6}^{7}$
Whence $\overline{27}|_{6}^{7}$ and $\overline{16}|_{6}^{1}$. Ans

2. Reduce $4^{\frac{1}{3}}$ and $5^{\frac{1}{4}}$ to quantities that shall have a common index.

Ans. $256^{\frac{1}{12}}$ and $125^{\frac{1}{12}}$.

3. Reduce $a^{\frac{1}{2}}$ and $a^{\frac{1}{3}}$ to quantities that shall have a common index.

Ans. a^3 and a^2 $\frac{1}{6}$.

4. Reduce $a^{\frac{1}{3}}$ and $b^{\frac{1}{4}}$ to quantities that shall have a common index.

Ans. $a^{\frac{1}{12}}$ and $b^{\frac{1}{3}}$.

5. Reduce $a^{\frac{1}{n}}$ and $b^{\frac{1}{m}}$ to quantities that shall have a common index.

Ans. $a^m \frac{1}{m}$ and $\overline{b^n} \frac{1}{mn}$.

CASE III.

To reduce surds to their most simple forms.

BULE.

Resolve the given number, or quantity, into two factors, one of which shall be the greatest power contained in it, and set the root of this power before the remaining part, with the proper radical sign between them.*

3, is a square.

^{*} When the given surd contains no factor that is an exact power of the kind required, it is already in its most simple form. Thus, V 15 cannot be reduced lower, because neither of its factors, 5, nor

EXAMPLES.

1. Let $\sqrt{48}$ be reduced to its most simple form. Here $\sqrt{48} = \sqrt{16 \times 3} = 4\sqrt{3}$ Ans.

2. Let 3/108 be reduced to its most simple form.

Here $\sqrt[3]{108} = \sqrt[3]{27} \times 4 = 3\sqrt[3]{4}$ Ans.

Note 1. When any number, or quantity, is prefixed to the surd, that quantity must be multiplied by the root of the factor above mentioned, and the product be then joined to the other part, as before.

EXAMPLES.

1. Let $2\sqrt{32}$ be reduced to its most simple form. Here $2\sqrt{32}=2\sqrt{16\times2}=8\sqrt{2}$ Ans.

2. Let $5\sqrt[3]{24}$ be reduced to its most simple form. Here $5\sqrt[3]{24}=5\sqrt[3]{8\times 3}=10\sqrt[3]{3}$ Ans.

Note 2. A fractional surd may also be reduced to a more convenient form, by multiplying both the numerator and denominator by such a number, or quantity, as will make the denominator a complete power of the kind required; and then joining its root, with 1 put over it, as a numerator, to the other part of the surd.*

EXAMPLES.

1. Let $\sqrt{\frac{2}{7}}$ be reduced to its most simple form.

^{*} The utility of reducing surds to their most simple forms, in order to have the answer in decirals, will be readily perceived from considering the first question above given, where it is found that $\sqrt{2} = \frac{1}{2} \sqrt{14}$; in which case it is only necessary to extract the square root of the whole number 14, (or to find it in some of the tables that have been calculated for this purpose) and then divide it by 7; whereas, otherwise, we must have first divided the numerator by the denominator, and then have found the root of the quotient, for the surd part; or else have determined the root both of the numerator and denominator, and then divided the one by the other; which are each of them troublesome processes when performed by the common rules; and in the next example, for the cube root, the labour would be much greater.

ere
$$\sqrt{\frac{2}{7}} = \sqrt{\frac{14}{49}} = \sqrt{\left(\frac{1}{49} \times 14\right)} = \frac{1}{7} \sqrt{14}$$
 Ans.

2. Let $3\sqrt[3]{\frac{2}{\epsilon}}$ be reduced to its most simple form.

ere $3\sqrt[3]{\frac{2}{5}} = 3\sqrt[3]{\frac{50}{105}} = 3\sqrt[3]{\left(\frac{1}{105} \times 50\right)} = \frac{3}{5}\sqrt[3]{50}$ Ans.

EXAMPLES FOR PRACTICE.

3. Let 125 be reduced to its most simple form. Ans. 5/5.

4. Let √294 be reduced to its most simple form.

Ans. 7 16.

- 5. Let 3/56 be reduced to its most simple form.
- 6. Let 3/192 be reduced to its most simple form. 7. Let 7 80 be reduced to its most simple form.
- Ans. 28./5. 8. Let 93/81 be reduced to its most simple form.
- Ans. 273/3.
- 9. Let $\frac{3}{191}\sqrt{\frac{5}{6}}$ be reduced to its most simple form. Ans. 212 /30.
- 10. Let $\frac{4}{7} \sqrt[3]{\frac{3}{16}}$ be reduced to its most simple form. Ans. 13/12.

11. Let $\sqrt{98a^2x}$ be reduced to its most simple form. Ans. 7a /2x. 12. Let $\sqrt{x^3 - a^2 x^2}$ be reduced to its most simple form.

Ans. $x \checkmark x - a^2$.

CASE IV.

To add surd quantities together.

RULE.

When the surds are of the same kind, reduce them to

their simplest forms, as in the last case; then, if the surd part be the same in them all, annex it to the sum of the rational parts, and it will give the whole sum required.

But if the quantities have different indices, or the surd

added together by the signs + and -.

And \/48=\/16\times3=4\/3

EXAMPLES.

1. It is required to find the sum of $\sqrt{27}$ and $\sqrt{48}$. Here $\sqrt{27} = \sqrt{9 \times 3} = 3\sqrt{3}$

Whence 7/3 the sum.

2. It is required to find the sum of 3/500 and 3/108. Here 3/500=3/125×4=53/4 And 3/108=3/27×4=33/4

Whence 83/4 the sum.

3. It is required to find the sum of 4\sqrt{147} and \sqrt{75}.

Here
$$4\sqrt{147}=4\sqrt{49\times3}=28\sqrt{3}$$

And $3\sqrt{75}=3\sqrt{25\times3}=15\sqrt{3}$

Whence 43/3 the sum.

4. It is required to find the sum of $3\sqrt{\frac{2}{5}}$ and $2\sqrt{\frac{1}{10}}$.

Here $3\sqrt{\frac{2}{5}} = 3\sqrt{\frac{10}{25}} = \frac{3}{5}\sqrt{10}$.

And $2\sqrt{\frac{1}{10}} = 2\sqrt{\frac{10}{100}} = \frac{2}{10}\sqrt{10}$.

Whence $\frac{4}{5}$ 10 the sum

11

EXAMPLES FOR PRACTICE.

- 5. It is required to find the sum of $\sqrt{72}$ and $\sqrt{128}$. Ans. $14\sqrt{(2)}$.
- 6. It is required to find the sum of 180 and 405.
 Ans. 15 4 (5).
- 7. It is required to find the sum of 33/40 and 3/135.
- 8. It is required to find the sum of $4\sqrt[3]{54}$ and $5\sqrt[3]{128}$.

 Ans. $32\sqrt[3]{2}$.
- 9. It is required to find the sum of 9 \(243 \) and 10 \(363. \)
 Ans. 191\((3). \)
- 10. It is required to find the sum of $3\sqrt{\frac{2}{3}}$ and $7\sqrt{\frac{27}{50}}$.
- Ans. $3\frac{1}{10}\sqrt{6}$. 11. It is required to find the sum of $12\sqrt[3]{\frac{1}{4}}$ and $3\sqrt[3]{\frac{1}{32}}$.
- Ans. $6\frac{2}{3}\sqrt[3]{2}$.

 12. It is required to find the sum of $\frac{1}{2}\sqrt{a^2b}$ and
- 12. It is required to find the sum of $\frac{1}{2}$ $\sqrt[3]{a^2b}$ and Ans. $\left(\frac{a}{2} + \frac{2x^2}{3}\right)\sqrt{b}$.

CASE V.

To find the difference of surd quantities.

RULE.

When the surds are of the same kind, prepare the quantities as in the last rule; then the difference of the rational parts annexed to the common surd, will give the whole difference required.

But if the quantities have different indices, or the surd part be not the same in each of them, they can only be subtracted by means of the sign —.

It is required to find the difference of √448 and √112.

Here
$$\sqrt{448} = \sqrt{64 \times 7} = 8\sqrt{7}$$

And $\sqrt{112} = \sqrt{16 \times 7} = 4\sqrt{7}$

Whence 4/7 the difference.

2. It is required to find the difference of $\sqrt[3]{192}$ and $\sqrt[3]{24}$.

Here $\sqrt[3]{192} = \sqrt[3]{64 \times 3} = 4\sqrt[3]{3}$ And $\sqrt[3]{24} = \sqrt[3]{8 \times 3} = 2\sqrt[3]{3}$

Whence 23/3 the difference.

3. It is required to find the difference of $5\sqrt{20}$ and $3\sqrt{45}$.

Here
$$5\sqrt{20}=5\sqrt{4\times5}=10\sqrt{5}$$

And $3\sqrt{45}=3\sqrt{9\times5}=9\sqrt{5}$

Whence \$\square\$5 the difference.

4. It is required to find the difference of $\frac{3}{4}\sqrt{\frac{2}{3}}$, and

$$\frac{2}{5}\sqrt{\frac{1}{6}}$$
.

Here
$$\frac{3}{4}\sqrt{\frac{2}{3}} = \frac{3}{4}\sqrt{\frac{6}{9}} = \frac{3}{12}\sqrt{6} = \frac{1}{4}\sqrt{6}$$

And $\frac{2}{5}\sqrt{\frac{1}{6}} = \frac{2}{5}\sqrt{\frac{6}{36}} = \frac{2}{30}\sqrt{6} = \frac{1}{15}\sqrt{6}$

Whence $\frac{11}{60}$ 6 the difference,

or answer required.

EXAMPLES FOR PRACTICE.

1. It is required to find the difference of $2\sqrt{50}$ and $\sqrt{18}$.

Ans. 7/(2).

63

- 2. It is required to find the difference of 3/320 and 3/40. Ans. 23/(5).
 - 3. It is required to find the difference of $\sqrt{\frac{3}{5}}$ and $\sqrt{\frac{5}{27}}$.
 - Ans. $\frac{4_3}{4_3}\sqrt{(15)}$.

 4. It is required to find the difference of $2\sqrt{\frac{1}{2}}$ and $\sqrt{8}$.

 Ans. $\sqrt{(2)}$.
 - 5. It is required to find the difference of $3\sqrt[3]{\frac{1}{3}}$ and $\sqrt[3]{72}$. Ans. $\sqrt[3]{9}$.
 - 6. It is required to find the difference of $\sqrt[3]{\frac{3}{3}}$ and $\sqrt[3]{\frac{3}{32}}$.
- Ans. $\frac{1}{2}\sqrt[3]{(18)}$.

 7. It is required to find the difference of $\sqrt{80a^4x}$ and $\sqrt{20a^2x^3}$.

 Ans. $(4a^2-2ax)\sqrt{(5x)}$.

8. It is required to find the difference of $(3\sqrt[3]{a})^3 b$ and $(3a - 2a^2)\sqrt[3]{(b)}$.

Note. The two last answers may be written thus, $(2ax-4a^2)\sqrt{(5x)}$, and $(2a^2-8a)^3\sqrt{(b)}$.

CASE VI.

To multiply surd quantities together.

RULE.

When the surds are of the same kind, find the product of the rational parts, and the product of the surds, and the two joined together, with their common radical sign between them, will give the whole product required; which may be reduced to its most simple form by Case III.

But if the surds are of different kinds, they must be reduced to a common index, and then multiplied together as usual.

It is also to be observed, as before mentioned, that the product of different powers, or roots, of the same quantity, is found by adding their indices.

EXAMPLES.

1. It is required to find the product of 3 18 and 2 16. Here 3 1/8

Multiplied 2 V 6

Gives $6\sqrt{48}=6\sqrt{16}\times 3=24./3$ Ans. 2. It is required to find the product of $\frac{1}{2}\sqrt[3]{\frac{2}{3}}$ and $\frac{3}{4}\sqrt[3]{\frac{5}{6}}$.

Here Multiplied $\frac{3}{4}\sqrt[3]{\frac{5}{6}}$

 $\frac{3}{6}\sqrt[3]{\frac{10}{12}} = \frac{3}{6}\sqrt[3]{\frac{5}{9}} = \frac{3}{6}\sqrt[3]{\frac{15}{97}} = \frac{1}{8}\sqrt[3]{15}.$

3. It is required to find the product of $2^{\frac{1}{2}}$ and $3^{\frac{1}{3}}$.

Here $2^{\frac{1}{2}}=2^{\frac{3}{6}}=(2^3)^{\frac{1}{6}}=8^{\frac{1}{6}}$ And $3^{\frac{1}{3}} = 3^{\frac{2}{6}} = (3^2)^{\frac{1}{6}} = 9^{\frac{1}{6}}$

Whence (72) & Ans.

4. It is required to find the product of 5 √ a and 3 √ a.

Here $5\sqrt{a}=5a^{\frac{1}{2}}=5a^{\frac{3}{6}}$ And $3\sqrt[3]{a} = 3a^{\frac{1}{3}} = 3a^{\frac{2}{6}}$

Whence $15a^{\frac{5}{6}} = 15(a^{\frac{5}{6}})^{\frac{1}{6}}$ or $15\sqrt[6]{a^{\frac{5}{6}}}$ Ans.

EXAMPLES FOR PRACTICE.

- 5. It is required to find the product of 5/8 and 3/5. Ans. 30,/(10).
- 6. It is required to find the product of 3/18 and 53/4. Ans. 103/(9).

7. Required the product of $\frac{1}{4}\sqrt{6}$ and $\frac{2}{15}\sqrt{9}$.

Ans. $\frac{1}{10}\sqrt{(6)}$.

8. Required the product of $\frac{1}{2}\sqrt{18}$ and $5\sqrt{20}$.

Ans. 15./(10).

9. Required the product of 2/3 and 13½ 15.

Ans 27/(15)

10. Required the product of $72\frac{1}{4}a^{\frac{2}{3}}$ and $120\frac{1}{2}a^{\frac{1}{4}}$.

Ans. $8706\frac{1}{8}a^{\frac{1}{12}}$.

11. Required the product of $4+2\sqrt{2}$ and $2-\sqrt{2}$. Ans. 4.

12. Required the product of $(a+b)^{\frac{1}{n}}$ and $(a+b)^{\frac{1}{m}}$.

Ans. $(a+b)^{\frac{m+n}{m}}$.

CASE VII.

To divide one surd quantity by another.

BULE.

When the surds are of the same kind, find the quotient of the rational parts, and the quotient of the surds, and the two joined together, with their common radical sign between them, will give the whole quotient required.

But if the surds are of different kinds, they must be reduced to a common index, and then be divided as before.

It is also to be observed, that the quotient of different powers or roots of the same quantity, is found by subtracting their indices.

EXAMPLES.

1. It is required to divide 8 \(\square 108 \) by 2 \(\square 6 \).

Here
$$\frac{8\sqrt{108}}{2\sqrt{6}} = 4\sqrt{18} = 4\sqrt{9\times2} = 12\sqrt{2}$$
 Ans.

66 IRRATIONAL QUANTITIES, OR SURDS.

2. It is required to divide 83/512 by 4 3/2.

Here
$$\frac{8\sqrt[3]{512}}{4\sqrt[3]{2}} = 2\sqrt[3]{256} = 2\sqrt[3]{64} \times 4 = 8\sqrt[3]{4}$$
 Ans.

3. It is required to divide $\frac{1}{2}\sqrt{5}$ by $\frac{1}{3}\sqrt{2}$,

Here
$$\frac{\frac{1}{2}\sqrt{5}}{\frac{1}{3}\sqrt{2}} = \frac{3}{2}\sqrt{\frac{5}{2}} = \frac{3}{2}\sqrt{\frac{10}{4}} = \frac{3}{4}\sqrt{10}$$
 Ans.

4. It is required to divide \$\sqrt{7}\$ by \$\sqrt{7}\$.

Here
$$\frac{\sqrt{7}}{\sqrt[3]{7}} = \frac{7^{\frac{1}{2}}}{7^{\frac{1}{3}}} = \frac{7^{\frac{3}{6}}}{7^{\frac{2}{6}}} = 7^{\frac{3}{6}} = 7^{\frac{1}{6}}$$
 Ans.

5. It is required to divide 6 154 by 3 12.

6. It is required to divide 43/72 by 2 3/18.

Ans. 23/4.

7. It is required to divide $5\frac{3}{4}\sqrt{\frac{1}{135}} \text{ by } \frac{2}{3}\sqrt{\frac{1}{5}}$.

Ans. $\frac{2}{3}\frac{3}{4}\sqrt{3}$

3. It is required to divide $4\frac{5}{7}\sqrt{\frac{2}{3}}$ by $2\frac{2}{5}\sqrt{\frac{3}{4}}$.

Ans. $\frac{55}{4} \frac{5}{2} \sqrt{2}$. 9. It is required to divide $4\frac{1}{9} \sqrt[3]{a}$ by $2\frac{2}{3} \sqrt[3]{ab}$.

Ans.
$$\frac{27}{16} \left(\frac{a}{kz}\right)^{\frac{1}{6}}$$
.

10. It is required to divide $32\frac{2}{5}\sqrt{a}$ by $13\frac{3}{4}\sqrt[3]{a}$.

Ans. $\frac{648}{275}a^{\frac{3}{6}}$.

11. It is required to divide $9\frac{3}{8}a^{\frac{1}{n}}$ by $4\frac{9}{11}a^{\frac{1}{m}}$.

Ans. $\frac{825}{424}a^{\frac{m-n}{m}}$.

12. It is required to divide $\sqrt{20 + \sqrt{12}}$ by $\sqrt{5 + \sqrt{3}}$.

Ans. $\sqrt{4}$.

Note. Since the division of surds is performed by sub-

tracting their indices, it is evident that the denominator of any fraction may be taken into the numerator, or the numerator into the denominator, by changing the sign of its index.

Also, since $\frac{a^m}{a^m} = 1$, or $= a^{m-m} = a^o$, it follows, that the expression a^o is a symbol equivalent to unity, and consequently, that it may always be replaced by 1 whenever it

EXAMPLES.

1. Thus $\frac{1}{a} = \frac{a^{-1}}{1}$, or a^{-1} ; and $\frac{1}{a^n} = \frac{a^{-n}}{1}$, or a^{-n} .

occurs.*

2. Also,
$$\frac{b}{a^3} = \frac{ba^{-2}}{1}$$
, or ba^2 ; and $\frac{a^{-n}}{b^{-m}} = \frac{1}{a^n b^{-m}}$, or $\frac{b^m}{a^n}$.

3. Let $\frac{1}{a^2}$ be expressed with a negative index.

Ans. α-2.

4. Let $a^{-\frac{1}{2}}$ be expressed with a positive index.

$$\in$$
 Ans. $\frac{1}{a^{\frac{1}{2}}}$.

5. Let $\frac{1}{a+x}$ be expressed with a negative index.

Ans.
$$(a+x)^{-1}$$
.

6. Let $a(a^2-x^2)^{-\frac{1}{3}}$ be expressed with a positive index.

Ans.
$$\frac{1}{a(a^2-x^2)^{\frac{1}{3}}}$$
.

To what is above said, we may also farther observe,

a+0=a, and a-0=a.

2. Also, if nought be multiplied or divided by any quantity, both the product and quotient will be nought; because any number of times 0, or any part of 0, is 0; that is,

$$0 \times a$$
, or $a \times 0 = 0$, and $\frac{0}{a} = 0$.

^{1.} That 0 added to or subtracted from any quantity, makes it neither greater nor less; that is,

From this it likewise follows, that nought divided by nought, is a finite quantity, of some kind or other.

CASE VIII.

To involve, or raise surd quantities to any power.

RULE.

When the surd is a simple quantity, multiply its index by 2 for the square, by 3 for the cube, &c and it will give the power of the surd part, which being annexed to the proper power of the rational part, will give the whole power required. And if it be a compound quantity, multiply it by itself the proper number of times, according to the usual rule.*

For since $0 \times \alpha = 0$, or $0 = 0 \times \alpha$, it is evident, that $\frac{0}{0} = \alpha$.

4. Farther, if any finite quantity be divided by 0 the quotient will be infinite.

For let $\frac{a}{a}$ = q, then, if b remains the same, it is plain, the less a is, the greater will be the quotient q; whence, if a be indefinitely small, q will be indefinitely great: and consequently, when a is 0, the quotient q will be infinite: that is,

 $\frac{b}{0}$, or $\frac{1}{0}$ = ∞ .

Which properties are of frequent occurrence in some of the higher parts of the science, and should be carefully remembered.

Since, therefore, $\frac{1}{a+b}$ is the same as $(a+b)^{-1}$. Let us suppose, in the general formula, n=-1; and we shall have for the coefficients n=-1; $\frac{n-1}{2}=-1$; $\frac{n-3}{3}=-1$; $\frac{n-3}{4}=-1$, &c. and for the powers of a we

have $a = a \cdot \frac{1}{2} = \frac{1}{4}$; $a \cdot \frac{1}{2} = a \cdot \frac{1}{2} = \frac{1}{4^2}$; $a \cdot \frac{1}{2} = \frac{1}{4^3}$; $a \cdot \frac{1}{2} = \frac{1}{4^3}$ &c.:

so that $(a+b)^{-1} = \frac{1}{a+b} = \frac{1}{a} - \frac{b}{a^2} + \frac{b^3}{a^3} - \frac{b^3}{a^4} + \frac{b^4}{a^5} - \frac{b^5}{a^6}$, &c. which is the same series that is found by division. For more on this subject see the Binomial Theorem, (further on) or Euler's Algebra.

* When any quantity that is affected with the sign of the square root is to be raised to the second power, or squared, it is done by suppressing the sign of Thus.

 \sqrt{a})2, or $\sqrt{a} \times \sqrt{a} = a$; and $(\sqrt{a+b})$ 2, or $\sqrt{a+b} \times \sqrt{a+b} = a+b$.

EXAMPLES.

1. It is required to find the square of $\frac{2}{3}a^{\frac{1}{3}}$.

Here
$$\left(\frac{2}{3}a^{\frac{1}{3}}\right)^2 = \frac{4}{9}a^{\frac{1}{3}} \times 2 = \frac{4}{9}a^{\frac{2}{3}} = \frac{4}{9}\sqrt[3]{a^2}$$
 Ans.

2. It is required to find the cube of $\frac{2}{3}\sqrt{3}$.

Here
$$\frac{8}{27} \times 3^{\frac{3}{2}} = \frac{8}{27} \sqrt{27} = \frac{8}{27} \sqrt{9 \times 3} = \frac{8}{9} \sqrt{3}$$
 Ans.

3. It is required to find the square of 33/3.

Ans. $9\sqrt[3]{9}$. 4. It is required to find the cube of $17\sqrt{21}$.

- 4. It is required to find the cube of $17\sqrt{21}$.

 Ans. $103173\sqrt{(21)}$.
- 5. It is required to find the 4th power of $\frac{1}{6}\sqrt{6}$.
- 6. It is required to find the square of $3+2\sqrt{5}$.
 - Ans. $29+12\sqrt{5}$.
- 7. It is required to find the cube of $\sqrt{x+3}\sqrt{y}$.
- Ans. $x\sqrt{x+27y}\sqrt{x+9x}\sqrt{y+27y}\sqrt{y}$. 3. It is required to find the 4th power of $\sqrt{3}-\sqrt{2}$.

Ans. 49-20 √6.

CASE 1X.

To find the roots of surd quantities.

RULE.

When the surd is a simple quantity, multiply its index $y \frac{1}{2}$ for the square root, by $\frac{1}{3}$ for the cube root, &c and will give the root of the surd part; which being annexed to the root of the rational part, will give the whole root

70 IRRATIONAL QUANTITIES, OR SURDS.

· required. And if it be a compound quantity, find its root by the usual rule.*

EXAMPLES.

1. It is required to find the square root of 93/3. Here $(9\sqrt[3]{3})^{\frac{1}{2}} = 9^{\frac{1}{2}} \times 3^{\frac{1}{3}} \times \frac{1}{2} = 9^{\frac{1}{2}} \times 3^{\frac{1}{6}} = 3\sqrt[6]{3}$ Ans.

2. It is required to find the cube root of $\frac{1}{2}\sqrt{2}$.

Here
$$\left(\frac{1}{8}\sqrt{2}\right)^{\frac{1}{3}} = \left(\frac{1}{8}\right)^{\frac{1}{3}} \times (2^{\frac{1}{2} \times \frac{1}{3}}) = \frac{1}{2}(2^{\frac{1}{3}}) = \frac{1}{2}(\sqrt{2})$$
 Ans.

3. It is required to find the square root of 103. Ans. 10 / (10).

4. It is required to find the cube root of $\frac{8}{6\pi}a^4$.

Ans. 2α3/α.

5. It is required to find the 4th root of $\frac{16}{21}a^{\frac{2}{3}}$.

Ans. 2 a 6.

6. It is required to find the cube root of \(\frac{a}{2} \sqrt{\frac{a}{2}}. \)

Ans. $\sqrt{\frac{a}{3}}$, or $\frac{1}{3}\sqrt{(3a)}$. 7. It is required to find the square root of $x^2 - 4x \sqrt{c}$

Ans. x-2,/a. -1-4a. 8. It is required to find the square root of $a+2\sqrt{ab+b}$

Ans. 1/a+1/b.

Also, the nth root of the mth root of any number a, or the mth root of th

nth root of a, is a

^{*} The nth root of the mth power of any number a, or the mth power of the *n*th root of a, is $a = \frac{m}{n}$.

From which last expression, it appears that, that the square root of the square root of a is the 4th root of a; and that the cube root of the square root of a, or the square root of the cube root of a, is the 6th root of a; and so o for the fourth, fifth, or any other numerical root of this kind-

CASE X.

To transform a binomial, or a residual surd, into a general surd.

RULE.

Involve the given binomial, or residual, to a power corresponding with that denoted by the surd; then set the radical sign of the same root over it, and it will be the general surd required.

EXAMPLES.

- 1. It is required to reduce $2+\sqrt{3}$ to a general surd. Here $(2+\sqrt{3})^2 = 4+3+4\sqrt{3} = 7+4\sqrt{3}$; therefore $2+\sqrt{3}=\sqrt{7+4\sqrt{3}}$, the answer.
 - 2. It is required to reduce $\sqrt{2+\sqrt{3}}$ to a general surd. Here $(\sqrt{2+\sqrt{3}})^2=2+3+2\sqrt{6}=5+2\sqrt{6}$;
- therefore $\sqrt{2+\sqrt{3}} = \sqrt{5+2\sqrt{6}}$, the answer.
- 3. It is required to reduce $\sqrt[3]{2} + \sqrt[3]{4}$ to a general surd. Here $(\sqrt[3]{2} + \sqrt[3]{4})^3 = 6 + 6\sqrt[3]{2} + 6\sqrt[3]{4}$; therefore $\sqrt[3]{2} + \sqrt[3]{4} = \sqrt[3]{6(1+\sqrt{2}+\sqrt{4})}$, the answer.
 - 4. It is required to reduce $3 \sqrt{5}$ to a general surd.

 Ans. $\sqrt{(14-6\sqrt{5})}$.
 - 5. It is required to reduce $\sqrt{2}-2\sqrt{6}$ to a general surd. Ans. $\sqrt{(26-4\sqrt{12})}$.
 - 6. It is required to reduce $4 \sqrt{7}$ to a general surd.

 Ans. $\sqrt{(23 8\sqrt{7})}$.
 - 7. It is required to reduce 23/3 339 to a general surd. Ans. $\sqrt[3]{(16239 - 10833 - 219)}$.

CASE XI.

To extract the square root of a binomial, or residual surd.

RULE.*

Substitute the numbers, or parts, of which the given surd is composed, in the place of the letters, in one of the two following formulæ, according as it is a binomial or a residual, and it will give the root required.

$$\sqrt{a+\sqrt{b}} = \sqrt{\frac{1}{2}a + \frac{1}{2}\sqrt{a^2 - b}} + \sqrt{\frac{1}{2}a - \frac{1}{2}\sqrt{a^2 - b}}
\sqrt{a-\sqrt{b}} = \sqrt{\frac{1}{2}a + \frac{1}{2}\sqrt{a^2 - b}} - \sqrt{\frac{1}{2}a - \frac{1}{2}\sqrt{a^2 - b}}$$

And if the second part of the binominal, or residual, in this case, be an imaginary surd, the same theorems will still hold, by only changing -b into +b, as below.

$$\sqrt{(a+\sqrt{-b})} = \sqrt{(\frac{1}{2}a + \frac{1}{2}\sqrt{(a^2+b)})} + \sqrt{(\frac{1}{2}a - \frac{1}{2}\sqrt{(a^2+b)})}
\sqrt{(a+\sqrt{-b})} = \sqrt{(\frac{1}{2} + \frac{1}{2}\sqrt{(a^2+b)})} - \sqrt{(\frac{1}{2}a - \frac{1}{2}\sqrt{(a^2+b)})}$$

* Prop. 1. From the lemma, page 16. vol. ii. Bonnycastle's Algebra; it appears, that if two numbers a and b be prime to each other, the fraction

 $\frac{dn}{dm}$ can never be a whole number. Prop. 2. From the same principles it likewise follows, that no whole num-

ber can bave a vulgar fraction for its root. Prop. 3. It may, also, be farther proved, that the square, cube, &c. root

of a whole number, cannot be partly rational and partly irrational. For, if possible, let, in the first place, $\sqrt{x-x} \pm \sqrt{y}$, for the square root. Then, by squaring both sides, $x=x^2 \pm 2x \sqrt{y+y}$, and by transposition, $+2x\sqrt{y}=a-x^2-y$.

Whence, by division, $\pm \sqrt{y} = \frac{a - x^2 - y}{2x} = a$ rational quantity; which

is contrary to what has been proved in the last proposition. Prop. 4. In any equation of the form $x \pm \sqrt{y-a} \sqrt{\pm b}$, the rational parts, on the opposite sides, are equal to each other, and also their irrational parts.

For, if x be not equal to a, let it be equal to $a \pm z$. Then $a \pm z \pm \sqrt{y} =$ $a \pm \sqrt{b}$, or $+\sqrt{b} = \pm z \pm \sqrt{y}$. That is, the \sqrt{b} is partly rational and partly irrational, which, by the last proposition, has been shown to be impossible

Whence, x=a, and consequently $\sqrt{y}=\sqrt{b}$. Prop. 5. It may here also be shown, that the square root of a binomial or residual surd, may sometimes be made equal to the sum, or difference of a whole number and a surd, or of two surds. Thus, let $\sqrt{(a+\sqrt{b})}=x+y$;

Where it is to be observed, that the only cases that are useful in this extraction, are when a is rational, and $a^2 - b$ in the first of these formulæ, or $a^2 + b$ in the latter, is a complete square.

EXAMPLES.

1. It is required to find the square root of 11+172, or 111+6,/2.

Here,

$$\sqrt{\frac{1}{2}a + \frac{1}{2}\sqrt{a^2 - b}} = \sqrt{\frac{11}{2} + \frac{1}{2}\sqrt{121 - 72}} = \sqrt{\frac{11}{2} + \frac{7}{2}} = 3;$$
and
$$\sqrt{\frac{1}{2}a - \frac{1}{2}\sqrt{a^2 - b}} = \sqrt{\frac{11}{2} - \frac{1}{2}\sqrt{121 - 72}} = \sqrt{\frac{11}{2} - \frac{7}{2}} = \sqrt{2}$$

Whence $\sqrt{(11+6\sqrt{2})}=3+\sqrt{2}$, the answer required.

2. It is required to find the square root of 3-2/2.

$$\sqrt{\frac{1}{2}a + \frac{1}{2}\sqrt{a^2 - b}} = \sqrt{\frac{3}{2} + \frac{1}{2}\sqrt{9 - 8}} = \sqrt{\frac{3}{2} + \frac{1}{2}} = \sqrt{2}; \text{ and}$$

$$= \sqrt{\frac{1}{2}a - \frac{1}{2}\sqrt{a^2 - b}} = -\sqrt{\frac{3}{2} - \frac{1}{2}\sqrt{9 - 8}} = -\sqrt{\frac{3}{2} - \frac{1}{2}}$$

Whence $\sqrt{(3-2\sqrt{2})} = \sqrt{2-1}$, the answer required. 3. It is required to find the square root of $6\pm2\sqrt{5}$.

Ans. 15±1. 4. It is required to find the square root of 23±8./7.

Ans. 4±1/7.

then, $\sqrt{(a-\sqrt{b})}=x-y$. And, by squaring each of these equations, we shall have $a+\sqrt{b}=x^2+2xy+y^2$, and $a-\sqrt{b}=x^2-2xy$

 $+y^2$; whence, by addition, $2a=2x^2+2y^2$, or $a=x^2+y^2$. And, by multiplying the like sides of the same two equations, there will arise

 $(a+\sqrt{b})\times(a-\sqrt{b})=x^2-y^3$, or $\sqrt{(a^2-b)}=x^2-y^2$ Wherefore, by adding and subtracting,

Wherefore, by adding and subtracting, $a + \sqrt{(a^2 - b)} = 2x^2$, and $a - \sqrt{(a^2 - b)} = 2y^2$; or $x = \sqrt{(\frac{1}{2}a + \frac{1}{2}\sqrt{(a^2 - b)})}$, and $y = \sqrt{(\frac{1}{2}a - \frac{1}{2}\sqrt{(a^2 - b)})}$. Whence, $\sqrt{(a + \sqrt{b})} = \sqrt{(\frac{1}{2}a + \frac{1}{2}\sqrt{(a^2 - b)})} + \sqrt{(\frac{1}{2}a - \frac{1}{2}\sqrt{(a^2 - b)})}$, and $\sqrt{(a - \sqrt{b})} = \sqrt{(\frac{1}{2}a + \frac{1}{2}\sqrt{(a^2 - b)})} = \sqrt{(\frac{1}{2}a - \frac{1}{2}\sqrt{(a^2 - b)})}$. And, if b be negative instead of affirmative, the two formulæ, in that case, $\sqrt{(\frac{1}{2}a - \frac{1}{2}\sqrt{(a^2 - b)})} = \sqrt{(\frac{1}{2}a - \frac{1}{$

will become $\sqrt{(a+\sqrt{-a+b})} = \sqrt{(\frac{1}{2}a+\frac{1}{2}\sqrt{(a2+b)})} + \sqrt{(\frac{1}{2}a+\frac{1}{2})}$ and $\sqrt{(a-\sqrt{-b})} = \sqrt{(\frac{1}{2}a+\frac{1}{2}\sqrt{(a2+b)})} + \sqrt{(\frac{1}{2}a+\frac{1}{2}\sqrt{(a2+b)})}$; hence the above rule is evident. Q. E. D.

74 IRRATIONAL QUANTITIES, or SURDS.

5. It is required to find the square root of 36 ± 10 \checkmark 11.

Ans. $5 \pm \checkmark$ (11).

3. It is required to find the square root of $33 \pm 12 \checkmark 6$.

Ans. $3 \pm 2 \sqrt{6}$.

7. It is required to find the square root of $1 + 4\sqrt{-3}$, or $1 + \sqrt{-48}$. Ans. $2 + \sqrt{-3}$.

8. It is required to find the square root of $3\pm 4\sqrt{-1}$, or $3\pm \sqrt{-16}$. Ans. $2\pm \sqrt{-1}$.

9. It is required to find the square root of $-1+\sqrt{-8}$.

Ans. $1+\sqrt{-2}$.

10. It is required to find the square root of $a^2 + 2x\sqrt{(a^2-x^2)}$.

Ans. $x + \sqrt{(ax-x^2)}$.

11. It is required to find the square root of $6+2\sqrt{2}$ $-\sqrt{(12)}-\sqrt{(24)}$. Ans. $1+\sqrt{2}-\sqrt{3}$.

FOR TRINOMIAL, QUADRINOMIAL SURDS, &c.

RULE. Divide half the product of any two radicals by a third, gives the square of one radical part of the root; this repeated with different quantities, will give the squares of all the parts of the root, to be connected by + and -. But if any quantity occur oftener than once, it must be taken but once.

For if x+y+z be any trinomial surd, its square will be $x^2+y^2+z^2+2xy+2xz+2yz$; then if half the product of any two rectangles as $2xy\times 2xz$ (or $2x^2yz$) be divided by some third 2yz, the quotient $\frac{2x^2yz}{2yz}=x^2$, must

needs be the square of one of the parts; and the like for the rest.

EXAMPLE 1.

To extract the square root of $10 + \sqrt{(24)} + \sqrt{(40)} + \sqrt{(60)}$. Here $\frac{\sqrt{(24)} \times \sqrt{(40)}}{2\sqrt{(60)}} = 2$, and $\frac{\sqrt{(24)} \times \sqrt{(60)}}{2\sqrt{(40)}} = \sqrt{(25)} = 5$. And the root is $\sqrt{2} + \sqrt{3} + \sqrt{5}$.

EXAMPLE 2.

It is required to find the square root of $12 + \sqrt{(32)} - \sqrt{(48)} + \sqrt{(80)} - \sqrt{(24)} + \sqrt{(40)} - \sqrt{(60)}$.

Here $\frac{\sqrt{(32 \times 48)}}{2\sqrt{(80)}} = \frac{\sqrt{(24)}}{\sqrt{5}}$, this produces nothing.

Again, $\frac{\sqrt{(32 \times 48)}}{2\sqrt{(24)}} = \sqrt{(16)} = 4$. And $\frac{\sqrt{(40 \times 60)}}{2\sqrt{(24)}} = \sqrt{(25)} = 5$; and $\frac{\sqrt{(32 \times 40)}}{2\sqrt{(60)}} = \sqrt{4} = 2$; and $\frac{\sqrt{(48 \times 24)}}{2\sqrt{(32)}} = \sqrt{9} = 3$; and $\frac{\sqrt{(32 \times 80)}}{2\sqrt{(40)}} = \sqrt{(16)} = 4$, &c. therefore the parts of the root are $\sqrt{4}, \sqrt{5}, \sqrt{3}, \sqrt{2}, \sqrt{4}$, &c. and the root $2 + \sqrt{2} - \sqrt{3} + \sqrt{5}$; for being squared it produces the surd quantity given.

CASE XII.

To extract any root (c) of a binomial surd.

RULE I*.

Let the quantity be $A \pm B$, whereof A is the greater part and c the exponent of the root required. Seek the

^{*} Let the sum or difference of two quantities x and y be raised to a power whose exponent is c, and let the 1st, 3d, 5th, 7th, &c. terms of that power, collected into one sum, be called A, and the rest of the terms, in the even blaces, call B; the difference of the squares of x and y raised to the same power c.

For the terms in the c power of x + y writing for their coefficients, respectively, 1, c, d, c, &c. are x + cx y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y2 + ex y3 + &c. x + y + dx y3 + &c.

least number n whose power n^c is divisible by $A^2 - B^2$, the quotient being Q compute $\sqrt[6]{(A+B\times\sqrt{Q})}$ in the nearest integer number, which suppose to be r. Divide A / Q or by its greatest divisor, and let the quotient be s, and let

 $\frac{r+r}{q_0}=t$, the nearest integer. Then the root = $\frac{t \ s \pm \sqrt{(t^2 s^2 - n)}}{\frac{2c}{c}/Q}$, if the c root of $A \pm B$ can be extracted.

Let one, or both of the quantities x, y, b e a quadratic surd, that is, let x+y, the e root of the proposed binomial A+B belong to one of these forms, $p+l\sqrt{q}$, $l\sqrt{q}$, $l\sqrt{p}+l\sqrt{q}$, and it follows: 1. If $x+y=p+l\sqrt{q}$, e being any whole number, A, the sum of the odd terms, will be a rational number; and B, the sum of the terms in the even places, each of which involves an odd power of y, will be a rational number multiplied into the quadratic surd \sqrt{g} . 2. Let c, the exponent of the root sought, be an odd number, as we may

always suppose it, because if it is even, it may be halved by the extraction of the square root, till it becomes odd; and let $x + y = k \checkmark p + q$. Then A

will involve the surd \(\nu p \), and B will be rational.

3. But if both members of the root are irrational, $(x+y=k \vee p + l \vee q)$ A and B are both irrational, the one involving \sqrt{p} , and the other the surd \sqrt{q} . And in all these cases, it is easily seen that when x is greater than y, A will be greater than B. From this composition of the binomial $A + B_0$ we are led to its resolution, as in the foregoing rule, by these steps.

When A is rational, and $A_2 - B_2$ is a perfect c power.

1. By the theorem, $A2-B2=(x^2-y^2)^2$ accurately; and therefore extracting the cross of A2-B2 it will be x^2-y^2 , call this root n.

2. Extract in the nearest integer the e root of A+B, it will be (nearly)

x+y, which put =r. 3. Divide x^2-y^2 (=n) by x+y (=r) the quotient is (nearly) x-y; and the sum of the divisor and quotient is (more nearly) 2x; that is, if an integer value of x is to be found, it will be the nearest to

4.
$$x - (x - y) = y$$
; or, $\left(\frac{r + \frac{n}{2}}{2}\right)^2 - n = y$; whence $y = \sqrt{\left(\frac{r + \frac{n}{2}}{2}\right)^2 - n}$, and therefore putting $t = \frac{r + \frac{n}{2}}{2}$, the root sought $= t$

$$\sqrt{(\frac{1}{2}-n)}$$
, and therefore putting $t=\frac{1}{2}$, the root sought $-t$.

 $+\sqrt{(t^2-n)}$; the same expression as in the rule, when $Q=1$, $s=1$, that such on $A \ge R^2$ is a perfect convex, and the greater member A is re-

16 when A2 - B2 is a perfect c power, and the greater member A is rational.

EXAMPLE.

What is the cube root of $\sqrt{968 + 25}$.

We have $A^2 - B^2 = 343 = 7 \times 7 \times 7$. $Q \times 7^3 = n^3$, whence n = 7, and Q = 1. Then $\sqrt[3]{A + B} \times \sqrt{Q} = \sqrt[3]{56} + r = 4$. A $\sqrt{Q} = \sqrt{968} = 22 \sqrt{2}$, and the radical part $\sqrt{2} = s$, and $\frac{r + \frac{n}{r}}{2s} = \frac{4 + \frac{7}{4}}{2\sqrt{2}} = t = 2$, in the

II.

When A is irrational, and Q=1. By the same process, $x=\frac{r+\frac{n}{r}}{2}$

(= T) and $y = \sqrt{(T^2 - n)}$. But seeing A is supposed irrational, and c an odd number, x will be irrational likewise; and they will both involve the same irreducible surd \sqrt{p} , or s, which is found by dividing A by its greatest rational divisor. Write, therefore for x or T, its value $t \times s$, and $x + y = t + \sqrt{(2 s^2 - n)}$.

III.

If the c root of A2-B2 cannot be taken, multiply A2-B2 by a mimber Q_s such that the product may be the (least) perfect c power $n^c = A2$ Q_s — B2 Q_s). And now (instead of A+B) extract the croot of $(A+B) \times \sqrt{Q_s}$, which found as above, will be $ts + \sqrt{(t2 \cdot s2 - n)}$; and consequently the croot of A+B will be $ts + \sqrt{(t2 \cdot s2 - n)}$, divided by the c root of $\sqrt{Q_s}$; that is,

In the operation, it is required to find a number Q, such, that $(A2 - B2) \times Q$ may be a perfect c power; this will be the case, if Q be taken equal to $(A2 - B2)^{c-1}$; but to find a less number which will answer this condition, let A2 - B2 be divisible by $a_1 a_2 \dots (m)$; $b_1 b_2 \dots (n)$; $d_2 b_3 \dots d_n$

 $d_1, \ldots, (r)$; &c. in succession, that is, let A2 - B2 = ab b d &c. also, m+x + n+y + r+z let Q = ab b d &c. $(A2 - B2) \times Q = a \times b d$ &c. which is a perfect cth power, if x, y, z, &c. be so assumed that m+x, n+y, r+z, are respectively equal to c_1 or some multiple of c_2 . Thus to find a number which multiplied by 180 will produce a perfect cube, divide 180 as

often as possible by $\overset{\cdot}{2}$, $\overset{\cdot}{3}$, $\overset{\cdot}{5}$, &c. and it appears that $\overset{\cdot}{2}$. $\overset{\cdot}{2}$. $\overset{\cdot}{3}$. $\overset{\cdot}{3}$. $\overset{\cdot}{3}$. $\overset{\cdot}{3}$. Herefore, it be multiplied by $\overset{\cdot}{2}$. $\overset{\cdot}{3}$. $\overset{\cdot}{5}$. $\overset{\cdot}{5}$, it becomes $\overset{\cdot}{2}$. $\overset{\cdot}{3}$. $\overset{\cdot}{5}$, or $(\overset{\cdot}{2}$. $\overset{\cdot}{3}$. $\overset{\cdot}{5}$); a perfect cube.

If A and B be divided by their greatest common measure, either integer or quadratic surd, in all cases where the cth root can be obtained by this method, Q will either be unity, or some power of 2, less than 2°.

nearest integer. And $ts=2\sqrt{2}$, $\sqrt{(t^2 s^2 - n)} = \sqrt{(8-7)}$ = 1. $\sqrt[9]{(Q=1)}$. And the root is $\frac{2\sqrt{2+1}}{1} = 2\sqrt{2}$ + 1, whose cube, upon trial, 1 find to be $\sqrt{968+25}$.

Let the surd, that is to have its root extracted, be of the form $\sqrt[3]{(a+\checkmark b)}$, or $\sqrt[3]{(a-\checkmark b)}$. Then if a^2-b

that the same rule gives its root x-y. See Universal Arithmetic, p. 139. Dr. Waring's Med. Alg. p 287, or Maclaurin's Alg. p. 124.

* Thus, let $\sqrt{(\alpha + \sqrt{b})} = x + \sqrt{y}$; and we shall have, by involution,

 $a + \checkmark b = (x + \checkmark y)$. An equation which, by expanding the right hand member, and comparing the rational and irrational parts, gives

$$a = x + \frac{n(n-1)}{2} x - y + \frac{n(n-1)(n-2)(n-3)}{2 \cdot 3 \cdot 4} x - y + &c.$$

$$\checkmark b = n x^{n-1} \checkmark y + \frac{n (n-1) (n-2)}{2.3} x^{n-3} y \checkmark y + \dots + &c.$$

Or which is the same thing, under a different form,

$$a = \frac{1}{2} \left\{ (x + \sqrt{y}) + (x - \sqrt{y}) \right\}$$

$$\sqrt{b} = \frac{1}{2} \left\{ (x + \sqrt{y}) - (x - \sqrt{y}) \right\}$$

Whence by squaring each of these equations and subtracting the latter from the former, we shall have

Or, by rejecting the terms that destroy each, and then multiplying by $\frac{1}{4}$,

$$a - b = (x - y)$$
, or $x - y = (a - b)^{\frac{1}{n}}$

Where supposing a-b to be a complete power of the nth degree, let 2 (a-b)n be put =c. Then since x-y=c, and consequently y=x-c, if this value be substituted for y, in the equation $x+\frac{n(n-1)}{2}$ $x+y+\frac{n(n-1)}{2}$

$$\frac{n(n-1)(n-2)(n-3)}{2\cdot 3\cdot 4\cdot 1}x^{n-4}\frac{2}{y+4c}$$
 &c. = a, we shall obtain an equa-

tion in which the value of x, as before mentioned, is rational, when the extraction required is possible.

be a perfect integral cube, and some whole number, can be found, that, when substituted for n, will make

$$n^3 - 3(\sqrt[3]{a^2 - b}) n = 2a$$

the roots of the two expressions, in this case will be

$$\sqrt[3]{(a+\sqrt{b})} = \frac{1}{2}n + \frac{1}{2}\sqrt{(n^3-4\sqrt[3]{a^3-b})}$$

$$\sqrt[3]{(a-\sqrt{b})} = \frac{1}{2}n - \frac{1}{2}\sqrt{(n^2-4\sqrt[3]{a^2-b})}$$

And if the second part of the binomial, or residual, be an

And if the second part of the binomial, or residual, be an imaginary surd, and $a^2 + b$ be a perfect integral cube, the axtraction may be effected, by finding the integral value of n in the following equation as before.

$$n^3 - 3(3/a^3 + b) n = 2a$$
.

In which last case, the roots of the two expressions will be,

$$\sqrt[3]{(a+\sqrt{-b})} = \frac{1}{2}n + \frac{1}{2}\sqrt{(n^2 - 4\sqrt[3]{a^2 + b})}$$

$$\sqrt[3]{(a-\sqrt{-b})} = \frac{1}{2}n - \frac{1}{2}\sqrt{(n^2 - 4\sqrt[3]{a^2 + b})}$$

each of which formulæ may be obtained, by barely changing the sign of b in the former.

EXAMPLE.

It is required to find the cube root of $10 \pm 6 \sqrt{3}$, or $10 \pm \sqrt{(108)}$.

Here
$$a = 10$$
, and $b = 108$; whence $\sqrt[3]{(a^2 - b)} = \sqrt[3]{100 - 108} = -2$, and $n^3 - 3(\sqrt[3]{a^2 - b})n = 20$,

or $n^3 + 6n = 20$ where it readily appears, from inspection, that n = 2. Whence $\sqrt[3]{(10 + \sqrt{108})} = \frac{2}{2} + \frac{1}{2} \sqrt{(4 - 4 \times - 2)} = \frac{1}{2} + \frac{1}{2} \sqrt{(12)} = 1 + \sqrt{3}$, and $\sqrt[3]{(10 - \sqrt{108})} = \frac{2}{2} - \frac{1}{2} \sqrt{(4 - 4 \times - 2)} = 1 - \frac{1}{2} \sqrt{12} = 1 - \sqrt{3}$.

EXAMPLES FOR PRACTICE.

3. Required the cube root of 68 - 1/4374.

Ans.
$$\frac{4-\sqrt{6}}{\sqrt[3]{2}}$$
.

2. Required the cube root of $11 + 5 \sqrt{7}$.

Ans.
$$\frac{\sqrt{7+1}}{\sqrt[3]{2}}$$
.

30 IRRATIONAL QUANTITIES, OR SURDS.

3. Required the cube root of $2\sqrt{7} + 3\sqrt{3}$.

Ans. $\frac{\sqrt{7} + \sqrt{3}}{2}$

- 4. Required the fifth root of $29\sqrt{6+41}\sqrt{3}$.

 Ans. $\frac{\sqrt{6}+\sqrt{3}}{2\sqrt{9}}$.
- 5. Required the cube root of 45 ± 29 12.

Ans $3 + \sqrt{2}$, and $3 - \sqrt{2}$.

6. Required the cube root of $9 \pm 4 \sqrt{5}$, or $9 \pm \sqrt{80}$.

Ans $\frac{3}{2} + \frac{1}{2} \sqrt{5}$, and $\frac{3}{4} - \frac{1}{2} \sqrt{-5}$.

7. Required the cube root of $20 \pm 68 \sqrt{-7}$.

- Ans. $5 + \sqrt{-7}$, and $5 \sqrt{-7}$. 8. It is required to find the cube root of $35 \pm 69 \sqrt{-6}$.
- Ans. $5+\sqrt{-6}$, and $5-\sqrt{-6}$. 9. It is required to find the cube root of $81 \pm \sqrt{-2700^*}$. Ans. $-3+2\sqrt{-3}$, and $-3-2\sqrt{-3}$.

where the root of $3+\sqrt{-\frac{1}{2}\frac{9}{7}}$, being now more easily found to be $-1+2\sqrt{-\frac{1}{5}}$, $-1+\frac{2}{3}\sqrt{-3}$, we shall have by multiplying by 3, (which is the cube root of 27), $-3+2\sqrt{-3}$, as above.

Thus in the example above given, $81 + \sqrt{-2700} = 27 \times (3 + \sqrt{-2700})$

Also, this is useful, in Cardaus' rule for cubic equations; thus, $\sqrt[3]{(31+\sqrt{(-2700)})} + \sqrt[3]{(81-\sqrt{(-2700)})} = -3 \times 2 = -6$, or $= \frac{-3}{2} \times 2 = -3$

or $\frac{n}{2} \times 2 = 9$, the imaginary parts vanishing, by the contrariety of their signs See De Moivre's appendix to Sanderson's Algebra, Universal Arithmetic, o Maclaurin's Algebra.

^{*} Whenever it can be done, the operation, in cases of this k...d, ought to be abridged, by dividing the given binomial by the gree's tube that it contains and then finding the root of the quotient; which being multiplied by the root of the cube, by which the binomial was divided, will give the root required

CASE XIII.

To find such a multiplier, or multipliers, as will make any binomial surd rational.

RULE.*

1. When one or both of the terms are any even roots, multiply the given biominal, or risidual, by the same ex-

THEOREM I.

Generally, if you multiply a = b by $a = -m + a = 2m \frac{m}{b} + a = 2m \frac$ n-3m 2mb , &c. continued till the terms be in number equal to m, the n n (n-m) n-2m m n-3m 2m m-4m product shall be a-b: for, (a+a) b+a b+a $(a - m) \times (a - b)$. &c. b

> $n \quad n-m \quad m \quad n-2 \quad m \quad 2 \quad m \quad n-3 \quad m \quad 3 \quad m$ a + a b + a b $n-m \ m \ n-2m \ 2m \ n-3m \ 3m$ b &c. -b b n α

THEOREM II.

n-m n-2mm n-3m2m n-4m3mb + a b - a b &c. multiplied by (a + b)gives a + b, which is demonstrated as the other. Here the sign of b is positive when m is an odd number. When any binomial surd is proposed, suppose the index of each number equal to m, and let n be the least integer n-m n-2mm n-3mnumber that is measured by m; then shall a + a b + a&c. give a compound surd, which multiplied into the proposed surd

a = b, will give a rational product. Thus to find the surd which multiplied by $\sqrt[3]{a-\sqrt[3]{b}}$, will give a rational quantity. Here $m=\frac{1}{a}$, and the

The demonstration of this rule is evident from the following theorems.

pression, with the sign of one of its terms changed; and repeat the operation in the same way, as long as there are surds, when the last result will be rational.

least number which is measured by $\frac{1}{3}$, is unit; let n=1, then shall a=1, b=1, b

Let $a \stackrel{t}{=} b$ be multiplied by $a \stackrel{n-m}{=} a \stackrel{n-2 \ m \ l}{=} a \stackrel{n-1}{=} b \stackrel{n-3 \ m}{=} a \stackrel{(2l)}{=} b \stackrel{n-4 \ m}{=} a \stackrel{n}{=} b \stackrel{n}{=} a \stackrel{n}{=}$

taken the least integer that shall give $\frac{n}{m}$ also an integer.

aken the least integer that shall give m also an integer. $n-m \quad n-2m \quad l \quad n-3m \quad (2l) \quad n-4m \quad (3l) \quad o \quad \frac{n}{m}$ Dem. $a \quad +a \quad b \quad +a \quad b \quad +a \quad b$, &c. $a \quad bm-1 \times l$

$$\begin{array}{c} \times (a \pm b). \\ \hline & n + a - b + a - b \\ & \pm a - b + a - b \\ & \pm a - b - a - b - a \\ \hline & n + a - b - a \\ \hline & n + a - b - a \\ & n + a$$

The sign of $b\overline{m}$ is positive only when \overline{m} is an odd number, and the binomial l

proposed is a + b. If any binomial surd is proposed whose two numbers have different indices, let these be m and l, and n equal to the least integer number that is measured by m and by m and m and m and m and m and m by m and m

shall give a compound surd, which multiplied by the proposed $a^{""} \pm b^{"}$, shall give a rational product.

Thus $\sqrt[2]{a} - \sqrt[3]{b}$ being given, suppose $m = 2, l = \frac{1}{3}$, and $\overline{l} = \frac{3}{2}$, therefore n = 3, and a = 4 b + a = 5 b + a = 6 b +

2. When the terms of the biominal surd are odd roots. the rule becomes more complicated; but for the sum or difference of two cube roots, which is one of the most useful cases, the multiplier will be a trinomial surd, consisting of the squares of the two given terms and their product, with its sign changed.

EXAMPLES.

1. To find a multiplier that shall render 5 + \square 3 rational.

Given surd
$$5 + \sqrt{3}$$

Multiplier $5 - \sqrt{3}$

Product 25-3=22, as required.

2. To find a multiplier that shall make \$\sqrt{5} + \sqrt{3} rational.

Given surd
$$\sqrt{5}+\sqrt{3}$$

Multiplier $\sqrt{5}-\sqrt{3}$

Product 5-3=2, as required.

3. To find multipliers that shall make \$15 + \$1/3 raional.

Given surd
$$4/5+4/3$$

1st multiplier $4/5-4/3$
1st product $\sqrt{5}-\sqrt{3}$
2d multiplier $\sqrt{5}+\sqrt{3}$

2d multiplier 15+13

2d product 5-3=2, as required.

4. To find a multiplier that shall make 3/7+3/3 raional.

By these Theorems any binomial surd whatsoever being given, you may nd a surd, which multiplied by it shall give a rational product. See Macurin's Algebra, page 112.

34 IRRATIONAL QUANTITIES, or SURDS.

Given surd
$$\sqrt[3]{7} + \sqrt[3]{3}$$

Multiplier $\sqrt[3]{7^2} - \sqrt[3]{(7 \times 3)} + \sqrt[3]{3^2}$
 $- \sqrt[3]{(3 \times 7^2)} - \sqrt[3]{(7 \times 3^2)} + \sqrt[3]{(7 \times 3^2)$

Product 7+3=10, as was required.

5. To find a multiplier that shall make $\sqrt{5} - \sqrt{x}$ rational. Ans. $\sqrt{5} + \sqrt{x}$.

6. To find a multiplier that shall make $\sqrt{a} + \sqrt{b}$ rational.

Ans $\sqrt{a} - \sqrt{b}$.

7. To find multipliers that shall make $a + \sqrt{b}$ rational.

Ans. $a - \sqrt{b}$.

8. It is required to find a multiplier that shall make

1- $\frac{3}{2}a$ rational. Ans. $1+\frac{3}{2}a+\frac{3}{4}a^2$. 9. It is required to find a multiplier that shall make $\frac{3}{3}-\frac{1}{2}\frac{3}{2}$ rational. Ans. $\frac{3}{9}+\frac{1}{2}\frac{3}{6}+\frac{1}{2}\frac{3}{4}$.

 $\sqrt[3]{3} - \frac{1}{2}\sqrt[3]{2}$ rational. Ans. $\sqrt[3]{9} + \frac{1}{2}\sqrt[3]{6} + \frac{1}{4}\sqrt[3]{4}$.

10. It is required to find a multiplier that shall make $\sqrt[4]{(a^3)} + \sqrt[4]{(b^3)}$, or $a_4^3 + b_4^3$ rational.

Ans. $\sqrt[4]{a^9} - \sqrt[4]{(a^6b^3)} + \sqrt[4]{(a^3b^6)} - \sqrt[4]{b^9}$.

CASE XIV.

To reduce a fraction, whose denominator is either a simple or a compound surd, to another that shall have a rational denominator.

RULE.

1. When any simple fraction is of the form $\frac{b}{\sqrt{a}}$, mu tiply each of its terms by \sqrt{a} , and the resulting fraction will be $\frac{b\sqrt{a}}{a}$.

Or when it is of the form $\frac{b}{\sqrt[3]{a}}$, multiply them by $\sqrt[3]{a^2}$, and the result will be $\frac{b\sqrt[3]{a^2}}{a}$.

And for the general form $\frac{b}{\sqrt[n]{a}}$, multiply by $\sqrt[n]{a^{n-1}}$, and the result will be $\frac{b\sqrt[n]{a^{n-1}}}{a}$.

2. If it be a compound surd, find such a multiplier, by the last rule, as will make the denominator rational; and multiply both the numerator and denominator by it, and the result will be the fraction required.

EXAMPLES.

1. Reduce the fractions $\frac{2}{\sqrt{3}}$ and $\frac{3}{\frac{1}{2}\sqrt[4]{5}}$, to others that shall have rational denominators.

Here
$$\frac{2}{\sqrt{3}} = \frac{2}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{2\sqrt{3}}{3}$$
; and $\frac{3}{\frac{1}{2}\sqrt{5}} = \frac{3}{\frac{1}{2}\sqrt{5}} \times \frac{4\sqrt{5}^3}{\sqrt{5}^3} = \frac{64\sqrt{5}^3}{\frac{1}{2}\times5} = \frac{64\sqrt{5}^3}{5} = \frac{64\sqrt{125}}{5}$ the answer required.

. Reduce $\frac{3}{\sqrt{5-\sqrt{2}}}$ to a fraction, whose denominator shall be rational.

Here
$$\frac{3}{\sqrt{5-\sqrt{2}}} \times \frac{\sqrt{5+\sqrt{2}}}{\sqrt{5+\sqrt{2}}} = \frac{3\sqrt{5+3}\sqrt{2}}{5-2} = \frac{3\sqrt{5+3}\sqrt{2}}{3} = \frac{\sqrt{5+\sqrt{2}}}{1} = \sqrt{5+\sqrt{2}}$$
 the answer required.

3. Reduce $\frac{\sqrt{2}}{3-\sqrt{2}}$ to a fraction, whose denominator shall be rational.

Here
$$\frac{\sqrt{2}}{3-\sqrt{2}} = \frac{\sqrt{2} \times (3+\sqrt{2})}{(3-\sqrt{2}) \times (3+\sqrt{2})} = \frac{3\sqrt{2+2}}{9-2} = \frac{2+3\sqrt{2}}{7}$$

= $\frac{2}{7} + \frac{3}{7}\sqrt{2}$ the answer required.

86 IRRATIONAL QUANTITIES, OR SURDS.

4. Reduce $\frac{\sqrt{6}}{\sqrt{7+\sqrt{3}}}$ to a fraction, that shall have a rational denominator.

Ans.
$$\sqrt{(42)} - \sqrt{(18)}$$
.

5. Reduce $\frac{x}{3+\sqrt{x}}$ to a fraction that shall have a rational denominator.

Ans.
$$\frac{3x-x\sqrt{x}}{9-x}$$
.

- 6. Reduce $\frac{a-\sqrt{b}}{a+\sqrt{b}}$ to a fraction, the denominator of which shall be rational.

 Ans. $\frac{a^2+b-2a\sqrt{b}}{a^2-b}$.
- 7. Reduce $\frac{10}{\sqrt[3]{7-\sqrt[3]{5}}}$ to a fraction that shall have a rational denominator.

Ans.
$$5 \times (\sqrt[3]{(49)} + \sqrt[3]{(35)} + \sqrt[3]{(25)}$$
.

8. Reduce $\frac{\sqrt[3]{3}}{\sqrt[3]{9}+\sqrt[3]{10}}$ to a fraction that shall have a rational denominator.

Ans.
$$\frac{3\sqrt[3]{9+3\sqrt[3]{(10)+\sqrt[3]{(300)}}}}{19}.$$

9. Reduce $\frac{4}{\sqrt{4+\sqrt[4]{5}}}$ to a fraction that shall have a rational denominator.

Ans.
$$\sqrt{(10)} - 2\sqrt{2} + (2-\sqrt{5}) \times \sqrt{5}$$
.

оF

ARITHMETICAL PROPORTION

AND PROGRESSION.

ARITHMETICAL PROPORTION, is the relation which two quantities of the same kind, have to two others, when the difference of the first pair is equal to that of the second.

Hence, three quantities are said to be in arithmetical proportion, when the difference of the first and second is equal to the difference of the second and third.

Thus, 2, 4, 6, and a, a+b, a+2b, are quantities in arith-

metical proportion.

And four quantities are said to be in arithmetical proportion, when the difference of the first and second is equal to the difference of the third and fourth.

Thus, 3, 7, 12, 16, and a, a+b, c, c+b, are quantities

in arithmetical proportion.

Arithmetical Progression is when a series of quantities increase or decrease by the same common difference.

Thus, 1, 3, 5, 7, 9, &c. and a, a+d, a+2d, a+3d, &c. are increasing series in arithmetical progression, the common differences of which are 2 and d.

And 15, 12, 9, 6, &c. and a, a—d, a—2d, a—3d, &c. are decreasing series in arithmetical progression, the common differences of which are 3 and d.

The most useful properties of arithmetical proportion and progression are contained in the following theorems:

1. If four quantities are in arithmetical proportion, the sum of the two extremes will be equal to the sum of the two means.

Thus, if the proportionals be 2, 5, 7, 10, or a, b, c, d; then will 2+10=5+7, and a+d=b+c.

2. And if three quantities be in arithmetical propor-

tion, the sum of the two extremes will be double the mean.

Thus, if the proportionals be 3, 6, 9, or a, b, c, then will $3+9=2\times6=12$, and a+c=2b.

3. Hence an arithmetical mean between any two quantities is equal to half the sum of those quantities.

Thus, an arithmetical mean between 2 and 4 is $=\frac{2+4}{2}$

=3; and between 5 and 6 it is $=\frac{5+6}{2}=5\frac{1}{2}$.

And an arithmetical mean between a and b is $\frac{a+b}{2}$.*

4. In any continued arithmetical progression, the sum of the two extremes is equal to the sum of any two terms that are equally distant from them, or to double the middle term, when the number of terms is odd.

Thus, if the series be 2, 4, 6, 8, 10, then will 2+10=

 $4+8=2\times 6=12.$

And, if the series be a, a+d, a+2d, a+3d, a+4d, then will $a+(a+4d)=(a+d)+(a+3d)=2\times(a+2d)$.

5. The last term of any increasing arithmetical series is equal to the first term plus the product of the common difference by the number of terms less one; and if the series be decreasing, it will be equal to the first term minus that product.

Thus, the nth term of the series a, a+d, a+2d, a+3d,

a+4d, &c. is a+(n-1)d.

Thus, $\frac{2a+b}{3}$ and $\frac{a+2b}{3}$ = two arithmetical means between a and b, a

being the less extreme and b the greater.

And $\frac{na+b}{n+1}$, $\frac{(n-1)a+2b}{n+1}$, $\frac{(n-2)a+3b}{n+1}$, &c. to $\frac{a+nb}{n+1}$ = any number (n)

of arithmetical means between a and b; where $\frac{b-a}{n+1}$ is the common difference; which being added to a, gives the first of these means; and her again to this last, gives the second; and so on.

^{*} If two, or more, arithmetical means between any two quantities be required, they may be expressed as below.

And the nth term of the series a, a-d, a-2d, a-3d, a - 4d, &c. is a - (n - 1)d.

6. The sum of any series of quantities in arithmetical progression is equal to the sum of the two extremes raultiplied by half the number of terms.

Thus, the sum of 2, 4, 6, 8, 10, 12, is $= (2+12) \times$

 $\frac{6}{9} = 14 \times 3 = 42.$

And if the series be a+(a+d)+(a+2d)+(a+3d)+(a+4d) &c. . . . +l, and its sum be denoted by S, we shall have $S=(a+l)\times_{0}^{n}$, where l is the last term, and n

the number of terms.

Or, the sum of any increasing arithmetical series may be found, without considering the last term, by adding the product of the common difference by the number of terms less one to twice the first term, and then multiplying the result by half the number of terms.

And, if the series be decreasing, its sum will be found by subtracting the above product from twice the first term, and then multiplying the result by half the number of

terms, as before.

Thus, if the series be a+(a+d)+(a+2d)+(a+3d)+(a+4d), &c. continued to n terms, we shall have

$$S = \left\{ 2\alpha + (n-1)d \right\} \times \frac{n}{2}.$$

And if the series be a+(a-d)+(a-2d)+(a-3d)+(a-4d), &c. to n terms, we shall have

$$S = \left\{ 2a - (n-1)d \right\} \times \frac{n}{2}(*).$$

^(*) The sum of any number of terms (n) of the series of natural numbers 1, 2, 3, 4, 6, 6, 7, &c. is = $\frac{n(n+1)}{2}$

Thus, 1+2+3+4+5, &c. continued to 100 terms, is $=\frac{100\times101}{2}=50$ 101 × = 5050.

EXAMPLES.

1. The first term of an increasing arithmetical series is 3, the common difference 2, and the number of terms 20; required the sum of the series.

First, $3+2(20-1)=3+2\times19=3+38=41$, the last

term.

And $(3+41) \times \frac{20}{2} = 44 \times \frac{20}{2} = 44 \times 10 = 440$, the sum required.

Or, $\{2 \times 3 + (20 - 1) \times 2\} \times \frac{20}{2} = (6 + 19 \times 2) \times 10 = (6 +$

 $+38) \times 10 = 44 \times 10 = 440$, as before.

2. The first term of a decreasing arithmetical series is 100, the common difference 3, and the number of terms 34; required the sum of the series.

First, $100-3(34-1)=100-3\times33=100-99=1$, the

last term.

And $(100+1) \times \frac{34}{2} = 101 \times \frac{34}{2} = 101 \times 17 = 1717$, the sum required.

Or,
$$\{2 \times 100 - (34 - 1) \times 3\} \times \frac{34}{2} = (200 - 33 \times 3) \times 17$$

 $=(200-99)\times17=101\times17=1717$, as before.

3. Required the sum of the natural numbers, 1, 2, 3, 4, 5, 6, &c. continued to 1000 terms. Ans. 500500.

4. Required the sum of the odd numbers 1, 3, 5, 7, 9, &c. continued to 101 terms.

Thus, 1+3+5+7+9, &c. continued to 50 terms, is =502=2500. And if any three of the quantities a, d, n, S, be given, the fourth may be found from the equation

 $S = \left\{ 2a \pm (n-1)d \right\} \times \frac{n}{2}, \text{ or } (a+l) \times \frac{n}{2}$

Where the upper sign + is to be used when the series is increasing, and the lower sign - when it is decreasing; also the last term $l = a \pm (n-1)d$, as above.

Also the sum of any number of terms (n) of the series of odd numbers $1, 3, 5, 7, 9, 11, &c, is = n^2$.

- 5. How many strokes do the clocks of Venice, which go on to 24 o'clock, strike in a day?
- 6. Required the 365th term of the series of even numbers 2, 4, 6, 8, 10, 12, &c. Ans. 730.
 - 7. The first term of a decreasing arithmetical series is
- 10, the common difference, and the number of terms
- 21; required the sum of the series. Ans. 140.
- 8. One hundred stones being placed on the ground, in a straight line, at the distance of a yard from each other; how far will a person travel, who shall bring them one by one, to a basket, placed at the distance of a yard from the Ans. 5 miles and 1300 yards. first stone?

OF

GEOMETRICAL PROPORTION

AND

PROGRESSION.

* GEOMETRICAL PROPORTION, is the relation which two quantities of the same kind have to two others, when the

extremes.

^{*} If there be taken any four proportionals, a, b, c, d, which it has been usual to express by means of points: thus, a: b :: c : d,

this relation will be denoted by the equation $\frac{a}{b} = \frac{c}{d}$; where the equal ratios are represented by fractions, the numerators of which are the antecedents, and the denominators the consequents. Hence, if each of the two numbers of this equation be multiplied by dd, there will arise ad = bc. From which it appears, as in the common rule, that the product of the two extremes of any four proportionals is equal to that of the means. And if the third c, in this case, be the same as the second, or c = b, the proportion is said to be continued, and we have ad =b2, or, b _ vad; where it is evident, that the product of the extremes of three proportionals is equal to the square of the mean: or, that the mean is equal to the square root of the product of thet we

antecedents, or leading terms of each pair, are the same parts of their consequents, or the consequent of the antecedents.

Also, if each number of the equation $ad \Longrightarrow bc$ be successively divided by d, dc, ac, &c. the results will give

suits will give
$$\frac{a}{b} = \frac{c}{d}$$

$$\frac{a}{c} = \frac{b}{d}$$

$$\frac{b}{a} = \frac{d}{c}$$
or the
$$\frac{a:b::c:d}{a:c::b:d}$$

$$b:a:d:c$$

$$b:a:d:c$$
&c.

So that, by following this method, we can easily obtain all the transformations of the terms of the proportion, that can be made to agree with the equations ad =bc.

In like manner, from the same equality $\frac{a}{b}=\frac{c}{d}$, there will result, by multi-

plication, the following equivalent forms: $\frac{ma}{mb} = \frac{nc}{nd}$; $\frac{ma}{nb} = \frac{mc}{nd}$

Which, being converted into proportions, become ma: mb::nc:nd, and ma: nb:: mc: nd. And, by taking any like powers, or roots, of the differ ent sides of the same equation, we have $\frac{a^m}{b^m} = \frac{c^m}{a^m}$. Or, putting the term in the form of a proportion, $a^m:b^m:c^m:d^m$. In which cases m and may be any whole or fractional numbers whatever.

Again, if there be taken the several equations.

be taken the several equations
$$\frac{a}{b} = \frac{c}{d}$$
 which correspond
$$\frac{e}{f} = \frac{g}{h}$$
 with
$$\frac{e}{h} = \frac{g}{m}$$
 the proportions
$$\begin{cases} a:b::c:d\\ e:f::g:h\\ i:k::l:m \end{cases}$$
 &c.

we shall have, by multiplying their like terms, $\frac{a \times e \times i &c.}{b \times f \times k &c.} = \frac{c \times g \times l &c.}{d \times h \times m &c.}$ Or, by putting the expression into the form of a proportion, as $a \in c$. : $b \in c$ &c.:: cgl&c.:dhm&c. Also, taking $\frac{a}{b} = \frac{c}{d}$, as before, we shall have, multiplication, $\frac{ma}{nb} = \frac{mc}{nd}$; and by augmenting or diminishing each side the equation by 1; $\frac{ma}{nb} \pm 1 = \frac{mc}{nd} \pm 1$; or $\frac{ma \pm nb}{nb} = \frac{mc \pm nd}{nd}$; which,

ing expressed in the form of a proportion, gives ma + nb : nb :: mc + n nd; or $ma \pm nb : mc \pm nd : : nb : nd$.

And if the above mentioned equation $\frac{a}{b} = \frac{c}{a}$, be put by a similar mu

And if two quantities only are to be compared together, the part or parts, which the antecedent is of its consequent,

plication of its terms, under the form $\frac{pa}{qb} = \frac{pc}{qd}$ and then augmented or dimiaished by 1, as in the last case, there will arise $pa \pm qb : pc \pm qd :: qb : qd$. Whence, dividing each of the antecedents of these two analogies by their consequents, the result will give $\frac{ma \pm nb}{mc \pm nd} - \frac{nb}{nd} = \frac{b}{a}$; and $\frac{pa \pm qb}{pc \pm qd} - \frac{qb}{qd}$ $=\frac{b}{2}$. And, consequently, as the two right hand members of these expres-

sions are each $=\frac{b}{d}$, we shall have $\frac{ma \pm nb}{mc \pm nd} = \frac{pa \pm qb}{pc \pm qd}$.

Or, by converting the corresponding terms of this equation into a proportion $ma \pm nb : mc \pm nd :: pa \pm qb : pc \pm qd$. Also, because the common equation $\frac{a}{b} = \frac{c}{d}$ gives $\frac{a}{c} = \frac{b}{d}$, if the latter be put under the equivalent forms

 $\frac{ma}{nc} = \frac{mb}{nd}$, and $\frac{pa}{qc} = \frac{pd}{qd}$, we shall obtain, by a similar process, $ma \pm nc$: $pa \pm qc$:: $mb \pm nd$: $pb \pm qd$; which two analogies may be considered as

general formulæ for changing the terms of the proportion a:b::c:d, without altering its nature. Thus, by supposing m, n, p, q, to be each=1, and taking the antecedents with the superior signs, and the consequents with the inferior, we have a+b:a-b:c+d:c-d, and a+c:a-c:b+d:b-d; which forms, together with several of those already given, are the usual transformations of the common analogy pointed out above.

In like manner, by taking m, n and p each =1, and q=0, there will arise $a \pm b : a :: c \pm d : c$, and $a \pm c : a :: b \pm d : b$; each of which proportions may be verified by making the product of the extremes equal to that of the

means, and observing that ad=bc. Lastly, taking any number of equations of the form before used, for expressing proportions, as $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = \frac{g}{h} = \&c.$; which, according to the

common method are called a series of equal ratios, and are usually denoted by a:b::c:d::e:f::g:h::&c. we shall necesssarily have from the

fractions being all equal to each other $\frac{a}{b} = q$, $\frac{c}{d} = q$, $\frac{e}{f} = q$, $\frac{g}{h} = q$, And by multiplying q by each of the denominators, a=bq, c=dq, e=fq, g=hq, &c.

Whence, equating the sum of all the terms on the left hand side of these equations, with those on the right, we have a+c+e+g+&c.=(b+d+f+h+&c.)q. And consequently, by division, and the properties of proportionals before shown.

 $\frac{a+c+e+g+\&c.}{b+d+f+h+\&c.} = \frac{a}{b} = \frac{a+c}{b+d} = \frac{a+c+e}{b+d+f} = \&c.$

which results show, that, in a series of equal ratios, the sum of any number of the antecedents is to that of their consequents, as one, or more of the antecedents, is to one, or the same number of consequents. Q. E. D.

or the consequent of the antecedent, is called the ratio; observing, in both cases, always to follow the same method.

Hence, three quantities are said to be in geometrical proportion, when the first is the same part, or multiple, of the second, as the second is of the third.

Thus, 3, 6, 12, and a, ar, ar2, are quantities in geome-

trical proportion.

And four quantities are said to be in geometrical proportion, when the first is the same part, or multiple, of the second, as the third is of the fourth.

Thus, 2, 8, 3, 12, and a, ar, b, br, are geometrical pro-

portionals.

Direct proportion, is when the same relation subsists between the first of four terms and the second, as between the third and fourth.

Thus, 3, 6, 5, 10, and a, ar, b, br, are in direct proportion.

Inverse, or reciprocal proportion, is when the first and

second of four quantities are directly proportional to the reciprocals of the third and fourth:

Thus, 2, 6, 9, 3, and a, ar, br, b, are inversely propor-

tional; because 2, 6, $\frac{1}{9}$, $\frac{1}{3}$, and a, ar, $\frac{1}{br}$, $\frac{1}{b}$ are directly

proportional.

GEOMETRICAL PROGRESSION is when a series of quantities have the same constant ratio; or which increase, or ecrease, by a common muliplier, or divisor.

Thus, 2, 4, 8, 16, 32, 64, &c. and a, ar, ar2, ar3, ar4,

&c. are series in geometrical progression.

The most useful properties of geometrical proportion and progression are contained in the following theorems:

1. If three quantities be in geometrical proportion, the product of the two extremes will be equal to the square of the mean.

Thus if the proportionals be 2, 4, 8, or a, b, c, then

will $2 \times 8 = 4^2$, and $a \times c = b^2$.

2. Hence, a geometrical mean proportional, between any two quantities, is equal to the square root of their product.

Thus, a geometric mean between 4 and 9 is = $\sqrt{36}$ =6. And a geometric mean between a and b is = $\sqrt{ab^*}$.

3. If four quantities be in geometrical proportion, the product of the two extremes will be equal to that of the means.

Thus, if the proportionals be 2, 4, 6, 12, or a, b, c, d; then will $2 \times 12 = 4 \times 6$, and $a \times d = b \times c$.

4. Hence, the product of the means of four proportional quantities, divided by either of the extremes, will give the other extreme; and the product of the extremes, divided by either of the means, will give the other mean.

Thus, if the proportionals be 3, 9, 5, 15, or a, b, c, d;

then will $\frac{9\times5}{3}$ =15, and $\frac{3\times15}{5}$ =9: also, $\frac{b\times c}{a}$ = d, and $a\times d$

$$\frac{a\times d}{c} = b.$$

5. Also, if any two products be equal to each other, either of the terms of one of them, will be to either of the terms of the other, as the remaining term of the last is to the remaining term of the first.

Thus, if ad=bc, or $2\times15=6\times5$, then will any of the following forms of these quantities be proportional:

Directly, a:b::c:d, or 2:6::5:15. Invertedly, b:a::d:c, or 6:2::15:5.

Alternately, a:c::b:d, or 2:5::6:15.

Conjunctly, a: a+b:: c: c+d, or 2:8:: 5:20.

 $\sqrt[4]{a3b}$, $\sqrt[4]{a2b}$ and $\sqrt[4]{ab}$ = three geometrical means between a and b. And generally,

$$\frac{1}{(a^n b)^{n+1}}, (a^{n-1}b^2)^{n+1}, (a^{n-2}b^3)^{n+1} = \text{any number } (n) \text{ of geome-}$$

trical means between a and b. Where $(\frac{b}{a})^{\frac{1}{n+1}}$ is the ratio: so that if a

be multiplied by this, it will give the first of these means; and this last being again multiplied by the same, will give the second; and so on.

^{*} If two, or more, geometrical means between any two quantities be required, they may be expressed as below:

 $[\]sqrt[3]{a2b}$ and $\sqrt[3]{ab2}$ = two geometrical means between a and b.

Disjunctly, a:b - a:: c:d - c, or 2:4::5:10. Mixedly, b+a:b - a::d+c:d - c, or 3:4::20:10.

In all of which cases, the product of the two extremes is equal to that of the two means.

6 In any continued geometrical series, the product of the two extremes is equal to the product of any two means that are equally distant from them; or to the square of the mean, when the number of terms is odd.

Thus, if the series be 2, 4, 8, 16, 32; then will

 $2 \times 32 = 4 \times 16 = 8^{2}$

7. In any geometrical series, the last term is equal to the product arising from multiplying the first term by such a power of the ratio as is denoted by the number of terms less one.

Thus, in the series 2, 6, 18, 54, 162, we shall have $2 \times 3^4 = 2 \times 81 = 162$.

And in the series a, ar, ar^2 , ar^3 , ar^4 , &c. continued to n terms, the last term will be

 $l = \sigma r^{n \cdot 1}$.

2 - 1

8. The sum of any series of quantities in geometrical progession, either increasing or decreasing, is found by multiplying the last term by the ratio, and then dividing the difference of this product and the first term by the difference between the ratio and unity.

Thus, in the series 2, 4, 8, 16, 32, 64, 128, 256, 512, we shall have $\frac{512 \times 2 - 2}{2} = 1024 - 2 = 1022$, the sum of

we shall have

the terms .

Or the same rule, without considering the last term,

may be expressed thus :

Find such a power of the ratio as is denoted by the number of terms of the series; then divide the difference between this power and unity, by the difference between the ratio and unity, and the result, multiplied by the first term, will be the sum of the series.

Thus, in the series $a+ar+ar^2+ar^3+ar^4$, &c. to ar^{n-1} we shall have

$$S = a \left(\frac{r^n - 1}{r - 1} \right)$$

Where it is to be observed, that if the ratio, or common multiplier, r, in this last series, be a proper fraction, and consequently the series a decreasing one, we shall have, in that case,

$$a+ar+ar^2+ar^3+ar^4$$
, &c. ad infinitum = $\frac{a}{1-r}$.

 Three quantities are said to be in harmonical proportion, when the first is to the third, as the difference between the first and second is to the difference between the second and third.

Thus, a, b, c, are harmonically proportional, when a:c:a-b:b-c, or a:c:b-a:c-b.

And c is a third harmonical proportion to a and b, when ab

$$c = \frac{ab}{2a - b}.$$

10. Four quantities are in harmonical proportion, when the first is to the fourth, as the difference between the first and second is to the difference between the third and fourth.

Thus, a, b, c, d are in harmonical proportion, when a:d::a-b:c-d, or a:d::b-a:d-c. And d is a fourth harmonical proportional to a, b, c, when d=

 $\frac{ac}{2a-b}$, in each of which cases it is obvious, that twice the first term must be greater than the second, or otherwise the proportionality will not subsist.

11. Any number of quantities a, b, c, d, e, &c. are in harmonical progression, if a:c::a-b:b-c; b:d::b-c:c-d; c:e::c-d:d-e; &c.

12. The reciprocals of quantities in harmonical progression, are in arithmetical progression.

Thus, if a, b, c, d, e, &c. are in harmonical progression, $\frac{1}{a}$, $\frac{1}{b}$, $\frac{1}{c}$, $\frac{1}{d}$, $\frac{1}{e}$, &c. will be in arithmetical progression.

13. An harmonical mean between any two quantities, is equal to twice their product divided by their sum.

Thus, $\frac{2ab}{a+b}$ = an harmonical mean between a and b^* .

EXAMPLES.

1. The first term of a geometrical series is 1, the ratio 2, and the number of terms 10; what is the sum of the series.

Here $1 \times 2^9 = 1 \times 512 = 512$, the last term.

And
$$\frac{512 \times 2 - 1}{2 - 1} = \frac{1024 - 1}{1} = 1023$$
, the sum required.

2. The first term of a geometrical series is $\frac{1}{2}$, the ratio $\frac{1}{3}$, and the number of terms 5; required the sum of the series.

Here
$$\frac{1}{2} \times (\frac{1}{3})^4 = \frac{1}{2} \times \frac{1}{81} = \frac{1}{162}$$
, the last term.

- And $\frac{\frac{1}{2} \frac{1}{162} \times \frac{1}{3}}{1 \frac{1}{3}} = \frac{\frac{1}{2} \frac{1}{486}}{\frac{2}{3}} = \frac{121}{243} \times \frac{3}{2} = \frac{121}{162}$, the sum.

 3. Required the sum of 1, 2, 4, 8, 16, 32, &c. continued to 20 terms.

 Ans. 1048575.
- 4. Required the sum of 1, $\frac{1}{2}$, $\frac{1}{4}$, $\frac{1}{8}$, $\frac{1}{16}$, $\frac{1}{32}$, &c. continued to 8 terms.

 Ans. $1\frac{127}{128}$

\$0

D01

A Life

teire

then

25 93

bons.

- 5. Required the sum of 1, $\frac{1}{3}$, $\frac{1}{9}$, $\frac{1}{27}$, $\frac{1}{81}$, &c. continued to 10 terms.

 Ans. 1 $\frac{9841}{19683}$
- 6. A person being asked to dispose of a fine horse, said he would sell him on condition of having a farthing for

^{*} In addition to what is here said, it may be observed, that the ratio of two squares is frequently called duplicate ratio; of two square roots, subduplicate ratio; of two cubes, triplicate ratio; and of two cube roots, subtriplicate ratio; &c.

the first nail in his shoes, a half-penny for the second, a penny for the third, twopence for the fourth, and so on, doubling the price of every nail, to 32, the number of nails in his four shoes; what would the horse be sold for at that rate?

Ans. 44739241. 5s. 3\frac{3}{4}d.

OF EQUATIONS.

THE DOCTRINE OF EQUATIONS is that branch of algebra, which treats of the methods of determining the values of unknown quantities by means of their relations to others which are known.

This is done by making certain algebraic expressions equal to each other (which formula, in that case, is called an equation), and then working by the rules of the art, till the quantity sought is found equal to some given quantity

and consequently becomes known.

The terms of an equation are the quantities of which it is composed; and the parts that stand on the right and left of the sign =, are called the two members, or sides, of the equation.

Thus, if x=a+b, the terms are x, a, and b; and the meaning of the expression is, that some quantity x, standing on the left hand side of the equation, is equal to the sum of the quantities a and b on the right hand side.

A simple equation is that which contains only the first

power of the unknown quantity: as,

x+a=3b, or ax=bc, or $2x+3a^2=5b^2$;

Where x denotes the unknown quantity, and the other letters, or numbers, the known quantities.

A compound equation is that which contains two or more different powers of the unknown quantity; as,

 $x^2 + ax = b$, or $x^3 - 4x^2 + 3x = 25$.

Equations are also divided into different orders, or receive particular names, according to the highest power of the unknown quantity contained in any one of their terms: as quadratic equations, cubic equations, biquadratic equations, &c. Thus, a quadratic equation is that in which the unknown quantity is of two dimensions, or which rises to the second power: as,

 $x^2 = 20$; $x^2 + ax = b$, or $3x^2 + 10x = 100$.

A cubic equation is that in which the unknown quantity is of three dimensions, or which rises to the third power: as,

 $x^3 = 27$; $2x^3 - 3x = 35$; or $x^3 - ax^2 + bx = c$.

A biquadratic equation is that in which the unknown quantity is of four dimensions, or which rises to the fourth power: as, $x^4 = 25$; $5x^4 - 4x = 6$; or $x^4 - ax^3 + bx^2 - cx = d$.

And so on for equations of the 5th, 6th, and other higher orders, which are all denominated according to the highest power of the unknown quantity contained in any one of their terms.

The root of an equation is such a number, or quantity, us, being substituted for the unknown quantity, will make both sides of the equation vanish, or become equal to each other.

A simple equation can have only one root; but every compound equation has as many roots as it contains dimensions, or as is denoted by the index of the highest power of the unknown quantity, in that equation.

Thus, in the quadratic equation $x^2 + 2x = 15$, the root, or value of x, is either + 3 or - 5; and, in the cubic equation $x^3 - 9x + 26x = 24$, the roots are 2, 3, and 4, as will be found by substituting each of these numbers for x.

In an equation of an odd number of dimensions, one of its roots will always be real; whereas in an equation of an even number of dimensions, all it roots may be imaginary; as roots of this kind always enter into an equation by pairs.

Such are the equations $x^2-6x+14=0$, and $x^4-2x^3-9x^2+10x+50=0^*$.

^{*} See, for a more particular account of the general theory of equations, further on, or Vol. II. of Bonnycastle's Treatise on Algebra, 8vo. 1820.

OF THE

RESOLUTION OF SIMPLE EQUATIONS,

Containing only one unknown Quantity.

The resolution of simple, as well as of other equations, is the disengaging the unknown quantity, in all such ex pressions, from the other quantities with which it is con nected, and making it stand alone, on one side of the equation, so as to be equal to such as are known on the other side; for the performing of which, several axioms and processes are required, the most useful and necessary of which are the following:*

CASE I.

Any quantity may be transposed from one side of an equation to the other, by changing its sign; and the two members, or sides, will still be equal.

Thus, if x+3=7; then will x=7-3, or x=4.

And, if x-4+6=8; then will x=8+4-6=6.

Also, if x-a+b=c-d: then will x=a-b+c-d.

And, if 4x-8=3x+20; then 4x-3x=20+8, and consequently x=28.

From this rule it also follows, that if a quantity be found on each side of an equation, with the same sign, it may

^{*} The operations required, for the purpose here mentioned, are chiefly such

as are derived from the following simple and evident principles:

1. If the same quantity be added to, or subtracted from, each of two equal quantities, the results will still be equal; which is the same, in effect, as taking any quantity from one side of an equation, and placing it on the other

side, with a contrary sign.

2. If all the terms of any two equal quantities, be multiplied or divided, by the same quantity, the products, or quotients thence arising, will be equal.

3. If two quantities, either simple or compound, be equal to each other,

any like powers, or roots, of them will also be equal.

All of which axioms will be found sufficiently illustrated, by the processes arising out of the several examples annexed to the six different cases given in the text.

be left out of both of them; and that the signs of all the terms of any equation may be changed from + to -, or from - to +, without altering its value.

Thus, if x+5=7+5; then, by cancelling, x=7.

And, if a-c=b-c; then, by changing the signs, x-a=c-b, of x=a+c-b.

EXAMPLES FOR PRACTICE.

- 1. Given 2x+3=x+17 to find x. Ans. x=14.
- 2. Given 5x 9 = 4x + 7 to find x. Ans. x = 16.
- 3. Given x+9-2=4 to find x. Ans. x=-3.
- 4. Given 9x-8=8x-5 to find x. Ans. x=3.
- 5. Given 7x+8-3=6x+4 to find x. Ans. x=-1.

CASE II.

If the unknown quantity, in any equation, be multiplied by any number, or quantity, the multiplier may be taken away, by dividing all the rest of the terms by it; and if it be divided by any number, the divisor may be taken away, by multiplying all the other terms by it.

Thus, if ax=3ab-c; then will $x=3b-\frac{c}{a}$. And, if 2x+4=16; then will x+2=3,

or x=8-2=6. Also, if $\frac{x}{a}=5+3$; then will x=10+6=16.

And, if $\frac{2x}{3}$ = 2=4; then 2x = 6=12, or, by division, x = 3=6, or x = 9.

EXAMPLES FOR PRACTICE.

- 1. Given 16x+2=34 to find x. Ans. x=2.
- 2. Given 4x-8=-3x+13 to find x. Ans. x=3.
- 3. Given 10x-19=7x+17 to find x. Ans. x=12.
- 4. Given 8x-3+9=-7x+9+27 to find x. Ans. x=2.

5. Given 3ax - 3ab = 12d. Ans. $x = b + \frac{4d}{a^2}$.

CASE III.

Any equation may be cleared of fractions, by multiplying each of its terms, successively, by the denominators of those fractions, or by multiplying both sides by the product of all the denominators, or by any quantity that is a multiple of them.

Thus, if $\frac{x}{3} + \frac{x}{4} = 5$, then, multiplying by 3, we have $x + \frac{3x}{4} = 15$; and this, multiplied by 4, gives 4x + 3x = 60; whence, by addition, 7x = 60, or $x = \frac{60}{7} = 8\frac{4}{7}$

And, if $\frac{x}{4} + \frac{x}{6} = 10$; then, multiplying by 12, (which is a multiple of 4 and 6,) 3x + 2x = 120, or 5x = 120, or $x = \frac{120}{5} = 24$.

It also appears, from this rule, that if the same number, or quantity, be found in each of the terms of an equation, either as a multiplier or divisor, it may be expunged from all of them, without altering the result.

Thus, if ax=ab+ac; then by cancelling, x=b+c.

And, if
$$\frac{x}{a} + \frac{b}{a} = \frac{c}{a}$$
; then $x+b=c$, or $x=c-b$.

EXAMPLES FOR PRACTICE.

1. Given
$$\frac{3x}{2} = \frac{x}{4} + 24$$
 to find x. Ans $x = 19\frac{1}{2}$.

2. Given
$$\frac{x}{3} + \frac{x}{5} + \frac{x}{2} = 62$$
 to find x. Ans. $x = 60$.

3. Given
$$\frac{x-3}{2} + \frac{x}{3} = 20 - \frac{x+19}{2}$$
 to find x.

Ans. $x=29\frac{1}{4}$.

4. Given
$$\frac{x+1}{2} + \frac{x+2}{3} = 16 - \frac{x+3}{4}$$
 to find x.

Ans. x = 13.

5. Given
$$\frac{x+a}{b} + \frac{x}{c} = \frac{2x}{a} + \frac{a+b}{d}$$
 to find x.

Ans.
$$x = \frac{a^2 cb - acb^2 - a^2 cd}{acd + abd - 2cbd}$$

CASE IV.

If the unknown quantity, in any equation, be in the form of a surd, transpose the terms so that this may stand alone, on one side of the equation, and the remaining terms on the other (by Case 1); then involve each of the sides to such a power as corresponds with the index of the surd, and the equation will be rendered free from any irrational expression.

Thus, if $\sqrt{x-2}=3$; then will $\sqrt{x}=3+2=5$, or, by

squaring, $x=5^2=25$.

And if $\sqrt{(3x+4)}=5$; then will 3x+4=25, or 3x=25-4=21, or $x=\frac{21}{2}=7$.

Also, if $\sqrt[3]{(2x+3)+4}=8$; then will $\sqrt[3]{(2x+3)}=8-4$ =4, or $2x+3=4^3=64$, and consequently 2x=64-3=61, or $x = \frac{61}{9} = 30\frac{1}{9}$.

EXAMPLES FOR PRACTICE.

1. Given $2\sqrt{x+3}=9$ to find x. Ans. x=9. 2. Given $\sqrt{(x+1)} - 2 = 3$ to find x. Ans. x = 24.

Ans. $x = 7\frac{2}{3}$.

3. Given $\sqrt[3]{(3x+4)}+3=6$ to find x. 4. Given $\sqrt{(4+x)}=4-\sqrt{x}$ to find x. Ans. $x=2\frac{1}{4}$.

5. Given
$$\sqrt{(4a^2+x^2)} = \sqrt[4]{(4b^4+x^4)}$$
 to find x .

Ans. $x = \sqrt{\frac{b^4-4a^4}{2a^2}}$

CASE V.

If that side of the equation, which contains the unknown quantity, be a complete power, the equation may be reduced to a lower dimension, by extracting the root of the said power on both sides of the equation.

Thus, if $x^2 = 81$; then $x = \sqrt{81} = 9$; and if $x^3 = 27$,

then $x = \sqrt[3]{27} = 3$.

Also, if $3x^2-9=24$; then $3x^2=24+9=33$, or $x^2=$

 $\frac{33}{3}$ = 11, and consequently $x = \sqrt{11}$.

And, if $x^2+6x+9=27$; then, since the left hand side of the equation is a complete square, we shall have, by extracting the roots, $x+3=\sqrt{27}=\sqrt{(9\times3)}=3\sqrt{3}$, or $x=3\sqrt{3}-3$.

EXAMPLES FOR PRACTICE.

1. Given $9x^2 - 6 = 30$ to find x. Ans. x = 2.

2. Given $x^3 + 9 = 36$ to find x. Ans. x = 3.

3. Given $x^2 + x + \frac{1}{4} = \frac{81}{4}$ to find x. Ans. x = 4.

4. Given $x^2 + ax + \frac{a^2}{4} = b^2$ to find x. Ans. $x = b - \frac{a}{2}$.

5. Given $x^2 + 14x + 49 = 121$ to find x. Ans. x = 4.

CASE VI.

Any analogy, or proportion, may be converted into an equation, by making the product of the two extreme terms equal to that of the two means.

Thus, if 3x : 16 :: 5 :: 6; then $3x \times 6 = 16 \times 5$, or $18x \times 6 = 40 \times 5$

=80, or $x = \frac{80}{18} = \frac{40}{9} = 4\frac{4}{9}$.

And, if $\frac{2x}{3}$: a::b:c; then will $\frac{2cx}{3}$ =ab, or 2cx=3ab;

or, by division, $x = \frac{3ab}{c}$.

Also, if $12-x: \frac{x}{0}: : 4:1$; then $12-x=\frac{4x}{0}=2x$, or 2x+x=12; and consequently $x=\frac{12}{9}=4$.

EXAMPLES FOR PRACTICE.

- 1. Given $\frac{3}{4}x$; a :: 5bc : cd to find x. Ans. $x = \frac{20ab}{3d}$.
- 2. Given $10-x:\frac{2}{3}x::3:1$ to find x. Ans. $x=3\frac{1}{3}$.
- 3. Given 8+8x:4x::8:2 to find x.
- Ans. x=1. Ans. x=2. 4. Given x:6-x:2:4 to find x.
- 5. Given 4x : a :: 9 / x : 9 to find x. Ans. x =

MISCELLANEOUS EXAMPLES.

- 1. Given 5x-15=2x+6, to find the value of x. Here 5x-2x=6+15, or 3x=6+15=21; and therefore $x = \frac{21}{3} = 7$.
- 2. Given 40-6x-16=120-14x, to find the value of x. Here 14x-6x=120-40+16; or 8x=136-40=96; and therefore $x = \frac{96}{9} = 12$.
- 3. Given $3x^2 10x = 8x + x^2$, to find the value of x. Here 3x-10=8+x, by dividing by x; or 3x-x=8+10=18, by transposition.

And consequently 2x=18, or $x=\frac{18}{9}=9$.

4. Given $6ax^3 - 12abx^2 = 3ax^3 + 6ax^2$, to find the value of x.

Here 2x-4b=x+2, by dividing by $3ax^3$; or 2x-x=2+4b; and therefore x=4b+2.

5. Given $x^2+2x+1=16$, to find the value of x

Here x+1=4, by extracting the square root of each side.

And therefore, by transposition, x=4-1=3.

6. Given 5ax-3b=2dx+c, to find the value of x. Here 5ax-2dx=c+3b; or (5a-2d)x=c+3b; and c+3b

therefore, by division, $x = \frac{c+3b}{5a-2d}$.

7. Given $\frac{x}{2} - \frac{x}{2} + \frac{x}{4} = 10$, to find the value of x.

Here $x - \frac{2x}{3} + \frac{2x}{4} = 20$; and $3x - 2x + \frac{6x}{4} = 60$; or 12x - 8x + 6x = 240; whence 10x = 240, or x = 24.

8. Given $\frac{x-3}{2} + \frac{x}{3} = 20 - \frac{x-19}{2}$, to find the value of x.

Here $x-3+\frac{2x}{3}=40-x+19$; or 3x-9+2x=120-3x+57; whence 3x+2x+3x=120+57+9; that is 8x=186, or $x=23\frac{1}{2}$.

9. Given $\sqrt{\frac{2x}{3}} + 5 = 7$, to find the value of x.

Here $\sqrt{\frac{2x}{3}} = 7 - 5 = 2$; whence, by squaring, $\frac{2x}{3} = 2^2 = 4$, and 2x = 12, or x = 6.

10. Given $x + \sqrt{(a^2 + x^2)} = \frac{2a^2}{\sqrt{(a^2 + x^2)}}$, to find the value of x.

Here $x\sqrt{(a^2+x^2)+a^2+x^2}=2a^2$; or $x\sqrt{(a^2+x)}=a^2-x^2$, and $x^2(a^2+x^2)=a^4-2a^2x^2+x^4$; whence $a^2x^2+x^4=a^4-2a^2x^2+x^4$, and $a^2x^2=a^4-2a^2x^2$; therefore $3x^2+x^2=a^4$, or $x=\frac{a^4}{3a^2}=\frac{a^2}{3}$; and consequently $x=\sqrt{\frac{a^2}{3}}$

 $=a\sqrt{\frac{1}{3}}=a\sqrt{\frac{3}{9}}=\frac{a}{3}\sqrt{3}$, the answer required.

EXAMPLES FOR PRACTICE.

1. Given 3x-2+24=31, to find the value of x.

Ans. x=3

2. Given 4-9y=14-11y, to find the value of y.

3. Given x+18=3x-5, to find the value of x.

4. Given $x + \frac{x}{2} + \frac{x}{2} = 11$, to determine the value of x.

Ans. x = 0

5. Given $2x-\frac{x}{2}+1=5x-2$, to find the value of x.

Ans. x=

6. Given $\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = \frac{7}{10}$, to determine the value of x.

x = 3 + 4 + 10 Ans. x = 1

7. Given $\frac{x+3}{2} + \frac{x}{3} = 4 - \frac{x-5}{4}$, to find the value of x.

Ans. $x = 3\frac{x}{4}$

8. Given $2+\sqrt{3x}=\sqrt{4+5x}$, to find the value of x.

Ans. x=1

9. Given $x + a = \frac{x^2}{a+x}$, to find the value of x.

Ans. x = -1

10. Given $\sqrt{x+\sqrt{a+x}} = \frac{2a}{\sqrt{(a+x)}}$ to find the valu of x. Ans. x=

11. Given $\frac{ax-b}{4} + \frac{a}{3} = \frac{bx}{2} - \frac{bx-a}{3}$, to find the value of x.

Ans. $x = \frac{3b}{3a-2}$

12. Given $\sqrt{a^2 + x^2} = \sqrt[4]{b^4 + x^4}$, to find the value of x.

Ans. $x = \sqrt{\frac{b^4 - a}{2 a^2}}$

13. Given $\sqrt{a+x} + \sqrt{a-x} = \sqrt{ax}$, to find the value of x.

Ans. $x = \frac{4a^2}{a^2+4}$

14. Given $\frac{a}{1+x} + \frac{a}{1-x} = b$, to determine the value of x.

Ans.
$$x = \sqrt{\frac{b-2a}{b}}$$

15. Given $a+x=\sqrt{a^2+x}\sqrt{(b^2+x^2)}$, to find the value of x.

Ans. $x=\frac{b^2}{4a}-a$

16. Given $\frac{1}{2}\sqrt{(x^2+3a^2)} - \frac{1}{2}\sqrt{(x^2-3a^2)} = x\sqrt{a}$, to find the value of x.

Ans. $x = \sqrt[4]{\frac{9a^3}{4-4a}}$

17. Given $\sqrt{(a+x)} + \sqrt{(a-x)} = b$, to find the value of x.

Ans $x = \frac{b}{a} \sqrt{(4a-b^2)}$

18. Given $\sqrt[3]{(a+x)} + \sqrt[3]{(a-x)} = b$, to find the value of x.

Ans. $x = \sqrt{a^2 - \left(\frac{b^3 - 2a}{3b}\right)^3}$

19. Given $\sqrt{a} + \sqrt{x} = \sqrt{ax}$, to find the value of x.

Ans.
$$x = \frac{a}{(\sqrt{a-1})^2}$$

20. Given $\sqrt{\left(\frac{x+1}{x-1}\right)} + \sqrt{\left(\frac{x-1}{x+1}\right)} = a$, to determine the

value of x. Ans. $x = \frac{a}{\sqrt{a^2 - 4}}$ 21. Given $\sqrt{(a^2 + ax)} = a - \sqrt{(a^2 - ax)}$, to find the value of x.

of x. Ans. $x = \frac{a}{2}\sqrt{3}$.

22. Given $\sqrt{(a^2-x^2)}+x\sqrt{(a^2-1)}=a^2\sqrt{(1-x^2)}$, to find the value of x.

Ans. $x=*\sqrt{(\frac{a^2-1}{x^2+2})}$

^{*} Bonnycastle, in his Key, says this answer is wrong; the answer which he gives, is $\sqrt{\frac{a^2-1}{(a^4+2a^2-3)}}$; Now these two answers give the same value for

23. Given
$$\sqrt{(x+a)} = c - \sqrt{(x+b)}$$
, to find the value of x .

Ans. $x = \left(\frac{c^2 + b - a}{2c}\right)^2 - b$

24. Given $\sqrt{\frac{b}{a+x}} + \sqrt{\frac{c}{a-x}} = \sqrt[4]{\frac{4bc}{a^2 - x^2}}$, to find the value of x .

Ans. $x = a\left(\frac{b+c}{b-c}\right)$

Of the resolution of simple equations, containing two unknown quantities.

When there are two unknown quantities, and two independent simple equations involving them, they may be reduced to one, by any of the three following rules;

RULE I.

Observe which of the unknown quantities is the least involved, and find its value in each of the equations, by the methods already explained; then let the two values, thus found, be put equal to each other, and there will arise

Consequency both give the same answer. Or thus, by hypothesis; Let
$$\sqrt{\binom{a^2-1}{a^2+3}} = \sqrt{\frac{a^2-1}{(a^4+2a^2-3)}},$$
 By squaring $\frac{a^2-1}{a^2+3} = \frac{a^4-2a^2+1}{a^4+2a^2-3}$, Dividing by a^2-1 we get $\frac{1}{a^2+3} = \frac{a^2-1}{a^2+3}$ and by multiplication at $1 + 2a^2 = \frac{a^2-1}{a^2+3} = \frac{a^2-1}{a^2+3}$.

 $\frac{az-1}{4+2az-3}$, and by multiplication a4+2az-3=a4+2az-3. Therefore the hypothesis is correct.

x; that is, $x = \sqrt{\frac{a^2-1}{(a^2+3)}}$, or $\sqrt{\frac{a^2-1}{(a^4+2a^2-3)}}$; because if we assume any value for a, and substitute it in both, we get the same value for x. Let a=2, then $\sqrt{\frac{a^2-1}{a^2+3}} = \sqrt{\frac{2}{3}}$ and $\sqrt{\frac{a^3-1}{a^4+2a^2-3}} = \sqrt{\frac{2}{(21)}} = \sqrt{\frac{2}{3}}$; consequently both give the same answer. Or thus, by hypothesis; Let $\sqrt{\frac{a^2-1}{a^2-3}} = \sqrt{\frac{a^2-1}{a^2-3}} = \sqrt{\frac{a^2-1}{a^2-3}}$

a new equation with only one unknown quantity in it, the value of which may be found as before.*

EXAMPLES.

1. Given
$$\begin{cases} 2x+3y=23\\ 5x-2y=10 \end{cases}$$
 to find the values of x and y .

Here, from the first equation, $x=\frac{23-3y}{a}$,

rom the first equation,
$$x = \frac{2}{2}$$
,

And from the second,
$$x = \frac{10+2y}{5}$$

23-3y 10+2y

Whence we have
$$\frac{23-3y}{2} = \frac{10+2y}{5}$$
,

Or
$$115-15y=20+4y$$
, or $19y=115-20=95$,
That is, $y=\frac{95}{19}=5$, and $x=\frac{23-15}{2}=4$.

2. Given
$$\begin{cases} x+y=a \\ x-y=b \end{cases}$$
 to find the values of x and y.

Here, from the first equation,
$$x=a-y$$
,
And from the second, $x=b+y$,
Whence $a-y=b+y$, or $2y=a-b$,

Whence
$$a-y=b+y$$
, or $2y=a-b$,

And therefore
$$y = \frac{a-b}{2}$$
, and $x = a-y$,

Or, by substitution,
$$x=a-\frac{a-b}{2}=\frac{a+b}{2}$$
.

3. Given
$$\begin{cases} \frac{1}{2}x + \frac{1}{3}y = 7 \\ \frac{1}{3}x + \frac{1}{2}y = 8 \end{cases}$$
 to find the values of x and y .

Here, from the first equation,
$$x=14-\frac{2y}{3}$$
,

And from the second,
$$x=24-\frac{3y}{2}$$
,

Therefore, by equality,
$$14 - \frac{2y}{3} = 24 - \frac{3y}{2}$$
,

^{*} This rule depends upon the well known axiom, that things which are equal to the same thing, are equal to each other; and the two following methods are founded on principles which are equally simple and obvious.

And consequently $42-2y=72-\frac{9y}{2}$,

Or by multiplication 84-4y=144-9y;

And, therefore, also 5y=144-84=60,

Or, by division, $x=\frac{60}{5}=12$, and $x=14-\frac{24}{3}=6$.

EXAMPLES FOR PRACTICE.

1. Given 4x+y=34, and 4y+x=16, to find the values of x and y.

Ans. x=8, y=2.

2. Given 2x+3y=16, and 3x-2y=11, to find the values of x and y.

Ans. x=5, y=2.

1 dues of x and y. 3. Given $\frac{2x}{5} + \frac{3y}{4} = \frac{9}{20}$, and $\frac{3x}{4} + \frac{2y}{5} = \frac{61}{120}$ to find the values of x and y. Ans. $x = \frac{1}{2}, y = \frac{1}{3}$.

4. Given $\begin{cases} \frac{1}{2}x + 2y = a \\ \frac{1}{2}x - 2y = b \end{cases}$ to find x and y.

Ans. x = a + b, and $y = \frac{1}{4}a - \frac{1}{4}b$.

5. Given $\begin{cases} \frac{x}{2} + \frac{y}{3} = 8 \\ \frac{x}{3} - \frac{y}{2} = 1 \end{cases}$ to find x and y.

Ans. x = 12, and y = 6.

6. Given $\begin{cases} \frac{x}{2} + \frac{y}{3} = 9 \\ x : y :: 4 : 3 \end{cases}$ to find x and y.
Ans. x = 12, and y = 9.

7. Given x+y=80, and $\frac{2x}{3} = \frac{3y}{4}$, to find x and y.

Ans. $x=42\frac{x}{1}$, and $y=37\frac{11}{12}$.

3. Given $y-6=\frac{x}{2}$, and x=y+6, to find x and y,

Ans. x=24, and y=18.

RULE II.

Find the value of either of the unknown quantities in that equation in which it is the least involved; then substitute this value in the place of its equal in the other equation, and there will arise a new equation with only one unknown quantity in it; the value of which may be found as before.

EXAMPLES.

1. Given $\begin{cases} x+2y=17 \\ 3x-y=2 \end{cases}$ to find the values of x and y.

From the first equation, x=17-2y; which value, being substituted for x, in the second,

gives
$$3(17-2y)-y=2$$
.
Or $51-6y-y=2$, or $7y=51-2=49$,
Whence $y=\frac{49}{7}=7$, and $x=17-2y=3$.

2. Given $\begin{cases} x+y=13\\ x-y=3 \end{cases}$ to find the values of x and y. From the first equation, x=13-y; which value being substituted for x, in the second,

Gives 13 - y - y = 3, or 2y = 13 - 3 = 10, Whence $y = \frac{10}{2} = 5$, and x = 13 - y = 8.

3. Given $\begin{cases} x:y::a:b\\ x^2+y^2=c \end{cases}$ to find the values of x and y. Here the analogy in the first, turned into an equation,

gives
$$bx=ay$$
, or $x=\frac{ay}{b}$

And this value, substituted for x in the second,

gives
$$\left(\frac{ay}{b}\right)^2 + y^2 = c$$
, or $\frac{a^2y^2}{b^2} + y^2 = c$,

Whence we have $a^2y^2 + b^2y^2 = b^2c$, or $y^2 = \frac{b^2c}{a^2 + h^2}$

And, consequently, $y=b\sqrt{\frac{c}{a^2+h^2}}$, and $x=a\sqrt{\frac{c}{a^2+h^2}}$

EXAMPLES FOR PRACTICE ..

1. Given $\frac{x}{7} + 7y = 99$, and $\frac{y}{7} + 7x = 51$, to find the values Ans. x=7, and y=14... of x and y.

2. Given
$$\frac{x}{2} - 12 = \frac{y}{4} + 8$$
, and $\frac{x+y}{5} + \frac{x}{3} - 8 = \frac{2y-x}{4} + 27$,

to find the values of x and y.

Ans. x=60, y=40.3. Given x+y=s, and $x^2+y^2=d$, to find the values of

x and y. Ans.
$$x = \frac{s^2 + d}{2s}$$
, $y = \frac{s^2 - d}{2s}$.

4. Given 5x-3y=150, and 10x+15y=825, to find x and y. Ans x=45, and y=25.

5. Given x+y=16, and x:y::3:1, to find x and y. Ans. x = 12, and y = 4.

6. Given $x + \frac{y}{6} = 12$, and $y + \frac{x}{2} = 9$, to find x and y.

Ans. x=10, and y=4. 7. Given x:y::3:2, and $x^2-y^2=20$, to find x and y. Ans. x=6, and y=4.

8. Given $\frac{x}{9} - 12 = \frac{y}{4} + 13 + \frac{x+y}{5} + \frac{x}{3} + 16 = \frac{2x-y}{4} + 27$, to find x and y. Ans. x=60, and y=20.

RULE III.

Let one or both of the given equations be multiplied, or divided, by such numbers, or quantities, as will make the term that contains one of the unknown quantities the same in each of them; then, by adding, or subtracting, the two equations thus obtained, as the case may require, there will arise a new equation, with only one unknown quantity in it, which may be resolved as before*.

may be applied to any particular case of this kind, by substituting the numeral of a, b, a', b', in the place of the letters, and observing, when either of them is negative, to change the signs accordingly.

^{*} The values of the unknown quantities in the two literal ax → by=c, and a'x + b'y = c', may be found in general terms, by multiplying the first by a', and the second by a, and then working according to the last rule, when the ac'-ca' $\frac{ac'-ca'}{ab'-ba'}$, and $x=\frac{cb'-bc'}{ab'-ba'}$; which solution results, so determined, will be y=

Where the numerator is the difference of the products of the opposite coefficients in the order in which y is not found, and the denominator is the difference of the products of the opposite coefficients taken from the orders that involve the two unknown quantities. Coefficients are of the same order which

EXAMPLES.

1. Given $\begin{cases} 3x + 5y = 40 \\ x + 2y = 14 \end{cases}$ to find the values of x and y. First, multiply the second equation by 3, and it will give

3x + 6y = 42.

Then, subtract the first equation from this, and it will give 6y - 5y = 42 - 40, or y = 2.

Whence, also, x=14-2y=14-4=10.

2: Given $\begin{cases} 5x - 3y = 9 \\ 2x + 5y = 16 \end{cases}$ to find the values of x and y.

Multiply the first equation by 2, and the second by 5; then 10x - 6y = 18, and 10x + 25y = 30.

And if the former of these be subtracted from the latter there will arise 31y=62, or $y = \frac{62}{31} = 2$.

Whence, by the first equation, $x = \frac{9+3y}{z} = \frac{15}{z} = 3$.

EXAMPLES FOR PRACTICE.

- 1. Given $\frac{x+8}{4} + 6y = 21$, and $\frac{y+6}{3} = 23 5x$, to find x Ans, x=4, and y=3. and y.
 - 2. Given 3x + 7y = 79, and $2y = 9 + \frac{x}{9}$, to find x and y.

Ans. x = 10, and y = 7.

3. Given 30x + 40y = 270, and 50x + 30y = 340, to find x and y. Ans. x=5, and y=3.

4. Given 3x - 3y = 2x + 2y, and x + y : xy : 3 : 5, to

find x and y.

Ans. x=10, and y.

5. Given $x^2y+xy^2=30$, and $x^3+y^2=35$, to find x.

Ans. x=3, and y=2.

either affect no unknown quantity, as c and c'; or the same unknown quantity n the different equations, as a and a'. Coefficients are opposite when they affect the different unknown quantities in the different equations, as a and b', s' and b.

6. Given
$$\frac{3x-5y}{2} = \frac{2x+4}{5} = 3$$
, and $8 = \frac{x-2y}{4} = \frac{x}{2} + \frac{y}{3}$, to Ans. $x = 12$, and $y = 6$.

find x and y.

7. Given x+y:a::x-y:b, and $x^2-y^2=c$, to find the values of x and y.

Ans. $\alpha = \frac{a+b}{2} \sqrt{\frac{c}{ab}}, y = \frac{a-b}{2} \sqrt{\frac{c}{ab}}$

8. Given ax+by=c, and dx+ey=f, to find the values of x and y.

Ans. $x = \frac{ce - bf}{ae - bd}$, $y = \frac{af - de}{ae - bd}$

9. Given x+y=a, and $x^2-y^2=b$, to find the values of Ans. $x = \frac{a^2 + b}{2a}$, $y = \frac{a^2 - l}{2a}$ x and y.

10. Given $x^2 + xy = a$, and $y^2 + xy = b$, to find the values Ans. $x = \frac{a}{\sqrt{(a+b)}}$, $y = \frac{b}{\sqrt{(a+b)}}$ of x and y.

Of the resolution of simple equations, containing three of more unknown quantities.

When there are three unknown quantities, and three independent simple equations containing them, they may be reduced to one, by the following method*.

RULE.

Find the values of one of the unknown quantities i each of the three given equations, as if all the rest wer

similar cases are so proposed as to be interpretation.

believes from the following example: x-2y+z=5; 2x+y-z=7; x+3y-2z=2;

where, if it were required to determine the values of x, y and z, it will be only be eliminating x from each of them, and then equating the results, the found by eliminating x from each of them, and y-3z=-3; 5y-3z=-3, and 5y-3z=-3, the state of the same furnish no determinative.

which equations, being identical, or both the same, furnish no determinate answer. And, in effect, if the three equations be properly examined, it wi be found, that the third is merely the difference of the first and second, an consequently involves no condition but what is contained in the other two.

^{*} The necessity for observing that the given equations, in this and other similar cases are so proposed as to be independent of each other, will be

mown; then put the first of these values equal to the seond, and either the first or second equal to the third, and here will arise two new equations with only two unknown quantities in them, the values of which may be found as in he former case; and thence the value of the third.

Or, multiply each of the equations by such numbers, or quantities, as will make one of their terms the same in hem all; then, having subtracted any two of these resulting equations from the third, or added them together, as he case may require, there will remain only two equations, which may be resolved by the former rules.

And in nearly the same way may four, five, &c. unmown quantities be exterminated from the same numer of independent simple equations; but, in cases of this ind, there are frequently shorter and more commodious nethods of operation, which can only be learnt from pracice*.

EXAMPLES.

1. Given
$$\left\{ \begin{array}{l} x+y+z = 29 \\ x+2y+3z = 62 \\ \frac{1}{2}x+\frac{1}{3}y+\frac{1}{4}z = 10 \end{array} \right\}$$
 to find $x, y, \text{ and } z$.

* The values of the unknown quantities in the three literal equations ax + by + cz = d; a'x + b'y + c'z = d'; a'x + b''y + c''z = d''; nay be exhibited, in general terms, like those before mentioned, as follows:

$$\begin{aligned} & = \frac{db'c'' - dc'b'' + ca'b'' - ba'c'' + bc'a'' - cb'a''}{ab'c'' - ac'b'' + ca'b'' - ba'c' + bc'a'' - cb'a''} \\ & = \frac{db'c'' - ac'b'' + ca'b'' - ba'c' + bc'a'' - cb'a''}{ab'c'' - ac'b'' + ca'b'' - ba'c'' + bc'a' - cb'a''} \\ & = \frac{ab'c'' - ac'b'' + ca'b'' - ba'c'' + bc'a'' - cb'a''}{ab'c'' - ac'b'' + bc'a'' - cb'a''} \\ & = \frac{ab'd'' - ac'b'' + ba'b'' - ba'c'' + bc'a'' - cb'a''}{ab'c'' - ac'b'' + bc'a'' - cb'a''} \end{aligned}$$

which formula, by substitution, may be employed for the resolution of any numeral case of this kind, as in the instance of two equations before given. The numerator of any of these equations, such as z, consists of all the different products, which can be made of three opposite coefficients taken from he orders in which z is not found; and the denominator consists of all the products that can be made of the three opposite coefficients taken from the products that can be made of the three opposite coefficients taken from the orders which involve the three unknown quantities.

Here, from the first equation, x=29-y-z. From the second, x=62-2y-3z,

And from the third, $x=20-\frac{2}{3}y-\frac{1}{2}z$,

Whence 29-y-z=62-2y-3z, And also 29-y-z=90-2y-1z

And, also, $29 - y - z = 20 - \frac{2}{3}y - \frac{1}{2}z$.

From the first of which y=33-2z, And from the second, $y=27-\frac{3}{2}z$,

Therefore $33-2z=27-\frac{3}{2}z$, or z=12,

Whence, also, y=33-2z=9And x=29-y-z=8.

2. Given $\begin{cases} 2x+4y-3z=22\\ 4x-2y+5z=18\\ 6x+7y-z=63 \end{cases}$ to find x, y, and z.

Here, multiplying the first equation by 6, the second by 3, and the third by 2, we shall have

12x + 24y - 18z = 132, 12x - 6y + 15z = 54, 12x + 14y - 2z = 126.

And, subtracting the second of these equations successively from the first and third, there will arise

30y - 33z = 78, 20y - 17z = 72.

Or, by dividing the first of these two equations by 3, and then multiplying the result by 2,

$$20y - 22z = 52$$
,
 $20y - 17z = 72$.

Whence, by subtracting the former of these from the latter, we have 5z=20, or z=4.

And, consequently, by substitution and reduction, y=7, and x=3.

3. Given x+y+z=53, x+2y+3z=105, and x+3y+4z=134, to find the values of x, y, and z.

Ans. x=24, y=6, and z=23.

4. Given
$$x + \frac{1}{2}y + \frac{1}{3}z = 32$$
, $\frac{1}{3}x + \frac{1}{4}y + \frac{1}{5}z = 15$, and $\frac{1}{4}x + \frac{1}{5}z = 12$, to find the values of x , y , and z

 $y + \frac{1}{6}z = 12$, to find the values of x, y, and z.

Ans. x=12, y=20, z=30.

5. Given 7x+5y+2z=79, 8x+7y+9z=122, and x+4y+5z=55 to find the values of x, y, and z.

Ans. x=4, y=9, z=3.

6. Given x+y=a, x+z=b, and y+z=c, to find the values of x, y, and z.

Ans. $y = \frac{a-b+c}{2}$, $x = \frac{a+b-c}{2}$ and $z = \frac{b-a+c}{2}$.

7. Given $\frac{x}{2} + \frac{y}{3} + \frac{z}{4} = 62$, $\frac{x}{3} + \frac{y}{4} + \frac{z}{5} = 47$, and $\frac{x}{4} + \frac{y}{5} + \frac{z}{6} = 38$, to find x, y, and z. Ans. x = 24, y = 60, and z = 120.

8. Given z+y=x+100, y-2x=2z-100, and z+100=3x+3y, to find x, y, and z

=3x+3y, to find x, y, and z. Ans. $x=9\frac{1}{11}$, $y=45\frac{5}{11}$, and $z=63\frac{7}{11}$.

9. Given x+y+z=7, 2x-3=y+3z, and 5x+5z=3y+19, to find x, y, and z.

10. Given $3x+5y-4z=2\dot{o}$, 5x-2y+3z=46, and 3y+19

iz-x=62, to find x, y, and z.

Ans. x=7, y=8, and z=9. 11. * Given x+y+z=13, x+y+u=17, x+z+u=18,

and y+z+u=21, to find x, y, z, and u. Ans. x=2, y=5, z=6, and u=10.

MISCELLANEOUS QUESTIONS,

PRODUCING SIMPLE EQUATIONS.

The usual method of resolving algebraical questions, s first to denote the quantities, that are to be found, by x, or some of the other final letters of the alphabet;

^{*} This can be resolved by proceeding after the same manner, as equations wolving three unknown quantities; but the resolution of it may be greatly utilitated, by introducing into the calculation, beside the principal unknown unnities, a new unknown quantity arbitrarily assumed, such as, for example, ee sum of all the rest: and when a little practised in such calculations, they ecome easy.

then, having properly examined the state of the question, perform with these letters, and the known quantities, by means of the common signs, the same operations and reasonings, that it would be necessary to mak if the quantities were known, and it was required to verify them, and the conclusion will give the result sought.

Or, it is generally best, when it can be done, to denot only one of the unknown quantities by x or y, and the to determine the expression for the others, from the nature of the question; after which the same method creasoning may be followed, as above. And, in some cases the substituting for the sums and differences of quantities or availing ourselves of any other mode, that a prope consideration of the question may suggest, will greatly facilitate the solution.

1. What number is that whose third part exceeds it

fourth part by 16?

Let x = the number required, Then its $\frac{1}{3}$ part will be $\frac{1}{3}x$, and its $\frac{1}{4}$ part $\frac{1}{4}x$.

And therefore $\frac{1}{3}x - \frac{1}{4}x = 16$, by the question,

That is
$$x - \frac{3}{4}x = 48$$
, or $4x - 3x = 192$,

Hence x=192, the number required.

2. It is required to find two numbers such, that the sum shall be 40, and their difference 16.

Let x denote the least of the two numbers required, Then will x+16 = to the greater number,

And x+x+16=40, by the question,

That is 2x=40-16, or $x=\frac{24}{2}=12=$ least number.

And x+16=12+16=28 = the greater number required 3. Divide 1000l. between A. B., and c., so that A sha have 72l. more than B., and c 100l. more than A.

Let x=B's share of the given sum, Then will x+72=A's share. And x+172=c's share. Hence their sum is x+x+72+x+172, Or 3x+244=1000, by the question, That is 3x=1000-244=756,

Or $x = \frac{756}{3} = 252l. = B$'s share,

Hence x + 72 = 324l. = A.'s share. And x + 172 = 424l. = C.'s share. Also, as above, 252l. = B.'s share.

Sum of all =1000l. the proof.

4. It is required to divide 1000l. between two persons, so that their shares of it shall be in the proportion of 7 to 9. Let x = the first person's share,

Then will 1000 - x = second person's share,

And x : 1000 - x :: 7 : 9, by the question, That is $9x = (1000 - x) \times 7 = 7000 - 7x$,

Or 9x+7x=7000, or $x=\frac{7000}{16}=437l$. 10s.=1st share, and 1000-x=1000-437l. 10s.=562l. 10s.=2d share.

5. The paving of a square court with stones, at 2s. a yard, will cost as much as the enclosing it with pallisades, at 5s. a yard; required the side of the square.

Let x= length of the side of the square sought, Then 4x= number of yards of enclosure, And x^2 = number of yards of pavement, Hence $4x \times 5 = 20x$ = price of enclosing it, And $x^2 \times 2 = 2x^2$ = the price of the paving, Therefore $2x^2 = 20x$, by the question,

Therefore $2r^2 = 20x$, by the question, Or 2x = 20, and x = 10, the length of the side required.

6. Out of a cask of wine, which had leaked away a third part, 21 gallons were afterwards drawn, and the cask being then guaged, appeared to be half full; how much did it hold?

Let x = the number of gallons the cask is supposed to have held.

Then it would have leaked away $\frac{1}{3}x$ gallons.

Whence there had been taken out of it, altogether, $21 + \frac{1}{2}x$ gallons,

And therefore $21 + \frac{1}{3}x = \frac{1}{9}x$, by the question,

That is $63+x = \frac{3}{6}x$, or 126+2x=3x,

Consequently 3x-2x 126, or x=126, the number of gallons required.

7. What fraction is that, to the numerator of which if 1 be added, its value will be $\frac{1}{3}$, but if 1 be added to the de-

nominator, its value will be $\frac{1}{4}$.

Let the fraction required be represented by $\frac{x}{y}$,

Then $\frac{x+1}{y} = \frac{1}{3}$, and $\frac{x}{y+1} = \frac{1}{4}$, by the question.

Hence 3x+3=y, and 4x=y+1, or $x=\frac{y+1}{4}$,

Therefore $3\left(\frac{y+1}{4}\right) + 3 = y$, or 3y + 3 + 12 = 4y,

That is y=15, and $x=\frac{y+1}{4}=\frac{15+1}{4}=\frac{16}{4}=4$,

Whence the fraction that was to be found, is $\frac{4}{15}$.

8. A market woman bought in a certain number of eggs at 2 a penny, and as many others at 3 a penny, and having sold them out again, altogether, at the rate of 5 for 2d., found she had lost 4d.; how many eggs had she?

Let x=the number of eggs of each sort,

Then will $\frac{1}{2}x$ = the price of the first sort,

And $\frac{1}{3}x$ = the price of the second sort,

But 5:2::2x (the whole number of eggs): $\frac{4x}{5}$,

Whence $\frac{4r}{5}$ = the price of both sorts, when mixed toge-

ther at the rate of 5 for 2d.,

And consequently $\frac{1}{2}x + \frac{1}{3}x - \frac{4x}{5} = 4$, by the question,

That is 15x+10x-24x=120, or x=120, the number

of eggs of each sort, as required.

9. If a can perform a piece of work in 10 days, and B in 13; in what time will they finish it, if they are both set about it together?

Let the time sought be denoted by x,

Then $\frac{x}{10}$ = the part done by λ in one day,

And $\frac{x}{13}$ = the part done by B in one day,

Consequently $\frac{x}{10} + \frac{x}{13} = 1$ (the whole work).

That is 13x+10x=130, or 23x=130,

Whence $x = \frac{130}{23} = 5\frac{15}{23}$ days, the time required.

10. If one agent A, alone, can produce an effect e, in the time α, and another agent B, alone in the time b; in what time will both of them together produce the same effect?
Let the time sought be denoted by x,

Then $a:e::x:\frac{ex}{a}$ = part of the effect produced by A.

And $b:e::x:\frac{ex}{b}$ = part of the effect produced by B.

Hence $\frac{ex}{a} + \frac{ex}{b} = e$ (the whole effect) by the question,

Or $\frac{x}{a} + \frac{x}{b} = 1$ by dividing each side by e,

Therefore $x + \frac{ox}{b} = a$, or bx + ax = ab,

Consequently $x = \frac{ab}{a+b}$ = time required.

11. How much rye at 4s. 6d. a bushel, must be mixed

with 50 bushels of wheat, at 6s. a bushel so that the mixture may be worth 5s. a bushel?

Let x=the number of bushels required, Then 9x is the price of the rye in sixpences,

And 600 the price of the wheat in ditto,

Also $(50+x)\times 10$ the price of the mixture in ditto, Whence 9x+600=500+10x, by the question,

Or, by transposition, 10x-9x=600-500,

Or, by transposition, 10x - 9x = 600 - 500, Consequently x = 100 the number of bushels required.

12. A labourer engaged to serve for 40 days, on condition that for every day he worked he should receive 20d,, but for every day he was absent he should forfeit 8d.; now at the end of the time, he had to receive 11.118.8d.; how many days did he work, and how many was he idle?

Let the number of days that he worked be denoted by

x,

Then will 40-x be the number of days he was idle, Also 20x the sum earned, and $(40-x)\times 3$.

Or 320-8x the sum forfeited,

Whence 20x - (320 - 8x) = 380d. (=11. 11s. 8d.), by the question,

That is 20x - 320 + 8x = 380, Or 28x = 380 + 320 = 700.

Consequently $x = \frac{700}{28} = 25$, the number of days he worked, and 40 - x = 40 - 25 = 15, the number of days he was idle.

QUESTIONS FOR PRACTICE.

1. It is required to divide a line, of 15 inches in length, into two such parts, that one may be three fourths of the other.

Ans. 84 and 63

2. My purse and money together are worth 20s. and the money is worth 7 times as much as the purse, how much is there in it?

Ans. 17s. 6d.

3 A shepherd, being asked how many sheep he had in

his flock, said, if I had as many more, half as many more, and 7 sheep and a half, I should have just 500; how many had he?

Ans. 197.

4. A post is one fourth of its length in the mud, one third in the water, and 10 feet above the water, what is its whole length?

Ans. 24 feet

5. After paying away $\frac{1}{4}$ of my money, and then $\frac{1}{5}$ of the

remainder, I had 72 guineas left; what had I at first?

Ans. 120 guineas

6 It is required to divide 300l. between, A. B., and C. so that A may have twice as much as B., and C as much as A and B together.

Ans. A 100l. B 50l. C 150l.

7. A person, at the time he was married was 3 times as old as his wife; but after they had lived together 15 years, he was only twice as old; what were their ages on their wedding day?

Ans. Bride's age 15, bridegroom's 45

two thirds of the remainder will be 40?

Ans 65

9. At a certain election, 1296 persons voted, and the successful candidate had a majority of 120; how many

voted for each?

Ans. 708 for one, and 588 for the other 10. A's age is double of B's, and B's is triple of c's, and the sum of all their ages is 140; what is the age of each?

Ans. A's 84, B's 42, and c's 14

11. Two persons A and B, lay out equal soms of money in trade; A gains 1201, and B loses 871., and A's money

is now double of B's what did each lay out?

Ans. 3001.

12. A person bought a chaise, horse and harness, for 601.; the horse came to twice the price of the harness, and the chaise to twice the price of the horse and harness; what did he give for each?

Ans. 13l. 6s. 8d. for the horse, 6l. 13s. 4d. for the harness, and 40l. for the chaise

13. A person was desirous of giving 3d. apiece to some beggars, but found he had not money enough in his pocket

by 8d., he therefore gave them each 2d., and had then 3d. remaining; required the number of beggars?

Ans. 11

14. A servant agreed to live with his master for N. a

year, and a livery, but was turned away at the end of seven months and received only 2l. 13s. 4d. and his livery; what was its value?

Ans. 4l. 16s.

15. A person left 560l, between his son and daughter, in such a manner, that for every half crown, the son should have, the daughter was to have a shilling; what

were their respective shares?

Ans. Son 400l., daughter 160l.

16. There is a certain number, consisting of two places of figures which is equal to four times the sum of its digits; and if 18 be added to it the digits will be inverted; what is the number?

Ans. 24

17. Two persons, A and B, have both the same income; A saves a fifth of his yearly, but B, by spending 50l. per annum more than A, at the end of four years, finds himself 100l. in debt; what was their income?

Ans. 1251.

18. When a company at a tavern came to pay their reckoning, they found, that if there had been three persons more, they would have had a shilling apiece less to pay, and if there had been two less, they would have had a shilling a piece more to pay; required the number of persons, and the quota of each?

Ans. 12 persons, quota of each 5s.

19. A person at a tavern borrowed as much money as he had about him, and out of the whole spent 1s.; he then went to a second tavern, where he also borrowed as much as he had now about him, and out of the whole spent 1s.; and going on, in this manner, to a third and fourth tavern, he found, after spending his shilling at the latter, that he had nothing left; how much money had he at first?

Ans. 11½d.

20. It is required to divide the number 75 into two such parts, that three times the greater shall exceed seven times the less by 15.

Ans. 54 and 21

21. In a mixture of British spirits and water, ½ of the whole plus 25 gallons was spirits, and 1 part minus 5 gallons was water; how many gallons were there of each?

Ans. 85 of wine, and 35 of water

22. A bill of 1201, was paid in guineas and moidores, and the number of pieces of both sorts that were used were just 100; how many were there of each, reckoning the guinea at 21s., and the moidore at 27s. ? 23. Two travellers set out at the same time from Lon-

don and York, whose distance is 197 miles; one of them goes 14 miles a day, and the other 16; in what time will

they meet?

Ans. 6 days 133 hours. 24. There is a fish whose tail weighs 9lb., his head weighs as much as his tail and half his body, and his body weighs as much as his head and his tail; what is the whole weight of the fish ?

25. It is required to divide the number 10 into three such parts, that, if the first be multiplied by 2, the second by 3, and the third by 4, the three products shall be all equal. Ans. 4-8, 3-1, and 2-4

26. It is required to divide the number 36 into three such parts, that $\frac{1}{2}$ the first, $\frac{1}{3}$ of the second, and $\frac{1}{4}$ of the

third, shall be all equal to each other.

Ans. The parts are 8, 12, and 16

- 27. A person has two horses, and a saddle, which, of itself, is worth 50l.; now, if the saddle be put on the back of the first horse, it will make his value double that of the second, and if it be put on the back of the second, it will make his value triple that of the first: what is the value of each horse? Ans. One 30l. and the other 40l.
- 28. If A gives B 5s. of his money, B will have twice as much as the other has left; and if B gives A 5s. of his money, a will have three times as much as the other has left: how much had each? Ans. A 13s. and B 11s.
- 29. What two numbers are those whose difference, sum and product, are to each other as the numbers 2, 3, and 5, respectively? Ans. 10 and 2

30. A person in play lost a fourth of his money, and then won back 3s., after which he lost a third of what he now had, and then won back 2s; lastly, he lost a seventh of what he then had, and after this found he had but 12s. remaining; what he had at first?

Ans. 20s.

31. A hare is 50 leaps before a greyhound and takes 4 leaps to the greyhound's 3, but 2 of the greyhound's leaps are as much as 3 of the hare's how many leaps must the greyhound take to catch the hare?

Ans. 300.

32. It is required to divide the number 90 into four such parts, that if the first part be increased by 2, the second diminished by 2, the third multiplied by 2, and the fourth divided by 2, the sun, difference, product, and quotient, shall be all equal?

Ars. The parts are 18, 22, 10, and 40

33. There are three numbers whose differences are equal, (that is, the second exceeds the first as much as the third exceeds the second), and the first is to the third as 5 to 7; also the sum of the three numbers is 324, what are those numbers?

Ans. 90, 108, and 126.

34. A man and his wife usually drank out a cask of beer in 12 days, but when the man was from home it lasted the woman 30 days; how many days would the man alone be in drinking it?

Ans. 20 days

in drinking it?

35. A general ranging his army in the form of a solid square, finds he has 284 men to spare, but increasing the side by one man, he wants 25 to fill up the square; how

many soldiers had he?

Ans. 24000

36. If a and b together can perform a piece of work m

8 days, A and c together in 9 days, and B and c in 10 days, how many days will it take each person to perform the the same work alone?

Ans. A $14\frac{34}{49}$ days, B $17\frac{23}{41}$, and C $23\frac{7}{31}$

QUADRATIC EQUATIONS.

A QUADRATIC EQUATION, as before observed, is that in which the unknown quantity is of two dimensions, or which rises to the second power; and is either simple or compound.

A simple quadratic equation, is that which contains only the square, or second power, of the unknown quantity, as

$$ax^2 = b$$
, or $x^2 = \frac{b}{a}$; where $x = \sqrt{\frac{b}{a}}$.

A compound quadratic equation, is that which contains both the first and second power of the unknown quantity; as

$$ax^2 + bx = c$$
, or $x^2 + \frac{b}{a}x = \frac{c}{a}$.

In which case, it is to be observed, that every equation of this kind, having any real positive root, will fall under one or other of the three following forms:

.*
$$x^2 + ax = b$$
 ... where $x = -\frac{a}{2} \pm \sqrt{(\frac{a^2}{4} + b)}$.

* It may be observed, with respect to these forms, that

In the coase az + az - b = 0, where $az = \frac{1}{2}a + \sqrt{\frac{1}{3}ae + b}$, or $-\frac{1}{2}a - \sqrt{\frac{1}{3}(a^2 + b)}$, the first value of a, must be positive, because $\sqrt{\frac{1}{3}a^2 + b}$ is greater than $\sqrt{\frac{1}{4}a^2}$, or its equal $\frac{1}{2}a$; and its second value will evidently be negative, because each of the terms, of which it is composed in negative.

2. In the case $x^2 - ax - b = a$, where $x = \frac{1}{2}a + \sqrt{(\frac{1}{2}a^2 + b)}$, or $\frac{1}{4}a - \sqrt{(\frac{1}{4}a^2 + b)}$, the first value of x, is manifestly positive, being the sum of two positive terms: and the second value will be negative, because $\sqrt{(\frac{1}{4}a^2 + b)}$

5. In the case $x^2 - ax + b = 0$, where $x = \lambda a + \sqrt{(\lambda a^2 - b)}$, or $\frac{1}{2}a - \sqrt{(\lambda a^2 - b)}$, both the values of x will be positive, when $\frac{1}{4}a^2$ is greater than b; for its first value, is then evidently positive, being composed of two positive terms; and its second value, will also be positive; because $\sqrt{(\lambda a^2 - b)}$ is less than $\sqrt{\lambda a^2}$, or its equal $\frac{1}{4}a$.

But if $\frac{1}{4}a^2$, in this case be less than b the solution of the proposed equation is inpossible; because the quantity $\frac{1}{4}a^2 - b$, under the radical, is the negative; and consequently $\sqrt{\frac{1}{4}a^2 - b}$ will be imaginary, or of no assign-

able value.

4. It may be also further observed, that there is a fourth case of the form $x^2+xx+b=0$, where $x=-\frac{1}{2}a+\sqrt{(\frac{1}{4}a^2+b)}$, or $x-\frac{1}{2}a-\sqrt{(\frac{1}{4}a^2-b)}$, the two values of x will be both negative, or both imaginary, according as $\frac{1}{4a^2}$ is greater or less than b; the imaginary roots, when they occur, being here of the forms $-(a'+c'\sqrt{-1})$ and $-(a'-c\sqrt{-1})$. From which it follows, that if all the terms of a quadratic equation, when

From which it follows, that if all the terms of a quadratic equation, when brought to the left hand side, be positive, its two roots will be both negative, or both imaginary; and conversely, if each of the roots be negative or each

imaginary, the signs of all the terms will be positive.

So that of all quadratic equations, which can have any real positive root, that of the third form, $x^2 - ax$, b = o, is the only one, where the solution, for certain numeral values of α and b, will become impossible.

2.
$$x - ax = b_{\perp}$$
. where $x = +\frac{a}{2} \pm \sqrt{\left(\frac{a^2}{4} + b\right)}$.
3. $x^2 - ax = -b$. where $x = +\frac{a}{2} \pm \sqrt{\left(\frac{a^2}{4} - b\right)}$.

Or, if the second and last terms be taken either positively or negatively, as they may happen to be, the general equation

 $ax^{2} \pm bx = \pm c$, or $x^{2} \pm \frac{b}{a}x = \pm \frac{c}{a}$

which comprehends all the three cases above mentioned, may be resolved by means of the following rule:

RULE.

Transpose all the terms that involve the unknown quantity to one side of the equation, and the known terms to the other; observing to arrange them so that the term which contains the square of the unknown quantity may be positive, and stand first in the equation.

Then, if this square has any coefficient prefixed to it, let all the rest of the terms be divided by it, and the equation will be brought to one of the three forms above

mentioned.

In which case, the value of the unknown quantity x is always equal to half the coefficient, or multiplier of x, in the second term of the equation, taken with a contrary sign, together with \pm the square root of the square of this number and the known quantity that forms the absolute or third term. of the equation.*

 $x^2 \pm ax = \pm b$ if the square of half the coefficient a of the second term of the equation be added to each of its sides, so as to render it of the form

^{*} This rule, which is more commodious in its practical application, than that usually given, is founded upon the same principle; being derived from the well known property, that in any quadratic $x^2 \pm ax = \pm b \text{ if the square of half the coefficient } a$

 $x^2 \pm ax + \frac{1}{4}a^2 = \frac{1}{4}a^2 \pm b$ that side which contains the unkno va quantit will then be a complete square; and, consequently, by extracting the root of each side, we shall have $x \pm \frac{1}{4}a = \pm \sqrt{\frac{1}{4}a^2 \pm b}, \text{ or } x = \pm \frac{1}{4}a \pm \sqrt{\frac{1}{4}a^2 \pm b},$

Note. All equations, which have the index of the unknown quantity, in one of their terms, just double that of the other, are resolved like quadratics, by first finding the value of the square root of the first term, according to the method used in the above rule, and then taking such a root, or power of the result, as is denoted by the reduced index of the unknown quantity.

Thus, if there be taken any general equation of this

kind, as

$$x^{2m}+ax^m=b$$

we shall have, by taking the square root of x^{2m} , and observing the latter part of the rule,

$$x^{m} = -\frac{a}{2} \pm \sqrt{\left(\frac{a^{2}}{4} + b\right)}$$
, and $x = \left\{-\frac{a}{2} \pm \sqrt{\left(\frac{a^{2}}{4} + b\right)}\right\} \left(\frac{a^{2}}{m} + b\right)$

And if the equation, which is to be resolved, be of the following form.

$$x^m \rightarrow \alpha x^{\frac{m}{2}} = b$$

we shall necessarily have, according to the same principle.

$$x^{\frac{m}{2}} = \frac{a}{2} \pm \sqrt{\frac{a^2}{4} + b}$$
, and $x = \left\{ \frac{a}{2} \pm \sqrt{\frac{a^2}{4} + b} \right\}^{\frac{2}{m}}$

EXAMPLES.

1. Given $x^2 + 4x = 140$, to find the value of x. Here $x^2 + 4x = 140$, by the question,

Whence $x=-2\pm\sqrt{(4+140)}$, by the rule, Or, which is the same thing, $x=-2\pm\sqrt{144}$,

which is the same as the rule, taking a and b in + or - as they may happen to be.

To this we may also further add, that from the constant occurrence of the onble sign before the radical part of the above expression, it necessarily folws, that every qua fratic equation must have two roots; which are either

oth real, or both imaginary according to the nature of the question

It may here, also, be observed, that the ambiguous sign \pm , which denotes oth + and -, is prefixed to the radical part of the value of x in every expression of this kind, because the square root of any positive quantity, as $a \ge a$ in the square root of a negative quantity, $a \ge a \ge a \ge a$. It with esquare root of a negative quantity, $a \ge a \ge a \ge a \ge a$. It is the square root of a negative quantity, $a \ge a \ge a \ge a \ge a \ge a \ge a$. It is the square root of a negative product.

To this we may also further add, that from the constant occurrence of the

Wherefore x=-2+12=10, or -2-12=-14, Where one of the values of x is positive and the other negative.

2. Given $x^2 - 12x + 30 = 3$, to find the value of x. Here $x^2-12x=3-30=-27$, by transposition, Whence $x=6\pm\sqrt{(36-27)}$, by the rule, Or, which is the same thing, $x=6\pm \sqrt{9}$,

Therefore x=6+3=9, or =6-3=3,

Where it appears that x has two positive values. 3. Given $2x^2 + 8x - 20 = 70$, to find the value of x. Here $2x^2 + 8x = 70 + 20 = 90$, by transposition, And $x^2 + 4x = 45$, by dividing by 2, Whence $x=-2\pm\sqrt{(4+45)}$, by the rule, Or which is the same thing, $x=-2\pm\sqrt{49}$.

Therefore x=-2+7=5, or =-2-7=-9, Where one of the values of x is positive and the other negative.

4. Given $3x^2 - 3x + 6 = 5\frac{1}{3}$, to find the value of x.

Here $3x^2 - 3x = 5\frac{1}{3} - 6 = -\frac{2}{3}$ by transposition.

And $x^2 - x = -\frac{2}{9}$ by dividing by 3,

Whence $x = \frac{1}{2} \pm \sqrt{\left(\frac{1}{4} - \frac{2}{9}\right)}$, by the rule,

Or, by subtracting $\frac{2}{9}$ from $\frac{1}{4}$, $x = \frac{1}{9} \pm \sqrt{\frac{1}{26}}$,

Therefore $x = \frac{1}{9} + \frac{1}{6} = \frac{2}{3}$, or $= \frac{1}{9} - \frac{1}{6} = \frac{1}{3}$.

In which case x has two positive values.

5. Given $\frac{1}{9}x^2 - \frac{1}{3}x + 20\frac{1}{2} = 42\frac{9}{3}$ to find the value of x.

Here $\frac{1}{6}x^2 - \frac{1}{2}x = 42\frac{2}{3} - 20\frac{1}{2} = 22\frac{1}{6}$ by transposition,

And $x^2 = \frac{2}{9}x = 44\frac{1}{3}$, by dividing by $\frac{1}{9}$, or multiplied

by 2, Whence we have $x=\frac{1}{3}\pm\sqrt{(\frac{1}{9}+44\frac{1}{3})}$, by the rule,

Or, by adding
$$\frac{1}{9}$$
 and $44\frac{1}{8}$ together, $x = \frac{1}{3} \pm \sqrt{\frac{400}{9}}$,

Therefore
$$x = \frac{1}{3} + 6\frac{2}{3} = 7$$
, or $= \frac{1}{3} - 6\frac{2}{3} = -6\frac{1}{3}$,

Where one value of x is positive, and the other negative.

6. Given $ax^2 + bx = c$, to find the value of x.

Here $x^2 + \frac{b}{a}x = \frac{c}{a}$ by dividing each side by a.

Whence, by the rule,
$$x = -\frac{b}{2a} \pm \sqrt{\left(\frac{b^2}{4a^2} + \frac{c}{a}\right)}$$
,

Or, multiplying c and a by 4a, $x = -\frac{b}{2a} \pm \sqrt{\frac{b^2 + 4ac}{4a^2}}$,

Therefore
$$x=-\frac{b}{2a}\pm\frac{1}{2a}\sqrt{(b^2+4ac)}$$
.

7. Given $ax^2 - bx + c = d$, to find the value of x. Here $ax^2 - bx = d - c$, by transposition,

And
$$x^2 - \frac{b}{a}x = \frac{d-c}{a}$$
 by dividing by a.

Whence
$$x = \frac{b}{2a} \pm \sqrt{\left(\frac{d-c}{a} + \frac{b^2}{4a^2}\right)}$$
 by the rule,

Or, mult
$$d-c \& a$$
 by $4a$, $x = \frac{b}{2a} \pm \frac{1}{2a} \sqrt{(4a(d-c)+b^2)}$

8. Given $x^4 + ax^2 = b$, to find the value of x. Here $x^4 + ax^2 = b$, by the question,

Or
$$x^2 = -\frac{a}{2} \pm \sqrt{(a^2 + b)} = -\frac{a}{2} \pm \frac{1}{2} \sqrt{(a^2 + 4b)}$$
, by the rule,

Whence $x = \pm \sqrt{(-\frac{a}{2} \pm \frac{1}{2} \sqrt{(4b + a^2)})}$ by extraction of roots.

9 Given $\frac{1}{2}x^{6} - \frac{1}{4}x^{3} = -\frac{1}{32}$, to find the value of x.

Here
$$\frac{1}{2}x^6 - \frac{1}{4}x^3 = -\frac{1}{32}$$
, by the question,

And
$$x^{0} - \frac{1}{2}x^{3} = -\frac{1}{16}$$
, by multiplying by 2,
Whence $x^{3} = \frac{1}{4} \pm \sqrt{(\frac{1}{16} - \frac{1}{16})}, = \frac{1}{4}$ by the rule,
And consequently $x = \sqrt[3]{\frac{1}{4}} = \sqrt[3]{\frac{2}{9}} = \frac{1}{9}\sqrt[3]{2}$.

10. Given $2x^{\frac{2}{3}} + 3x^{\frac{1}{3}} = 2$, to find the value of x. Here $2x^{\frac{2}{3}} + 3x^{\frac{1}{3}} = 2$, by the question, And $x^{\frac{2}{3}} + \frac{3}{6}x^{\frac{1}{3}} = 1$, by dividing by 2,

Whence
$$x^{\frac{1}{3}} = \frac{2}{4} + \sqrt{\left(\frac{9}{16} + 1\right)} = -\frac{3}{4} + \frac{5}{4} = \frac{1}{2}$$
, or -2 ,
Therefore $x = (\frac{1}{2})^3 = \frac{1}{8}$, or $(-2)^3 = -8$.

11. Given $x^4 - 12x^3 + 44x^2 - 48x = 9009(a)$, to find the value of a.

This equation may be expressed as follows, $*(x^2 - 6x)^2 + 8(x^2 - 6x) = a$,

* The method of expressing a biquadratic equation, by the form of a

quadratic, as above, is found thus: $\cot x^4 + 2ax^2 + 5ax^2 + 4a^3x = d$, be the biquadratic, which I am to resolve by a quadratic; first, find the two first terms of the square root of the left hand side of the equation, the square of which take from said side; then, if the remainder be divisible by the two terms of the root thus found; the equation can be resolved by a quadratic equation.

EXAMPLE. $x4 + 2ax^3 + 5a^2x^2 + 4a^3x = d$ $2x^{2} + ax)2ax^{3} + 5a^{2}x^{2}$ $2ax^{3} + a^{2}x^{2}$ (x2 + ax)

$$x^2 + ax$$
) $4a^2x^2 + 4a^3x(4a^2 + 4a^3x)$

Hence $(x^2 + ax)^3 + 4az(x^2 + ax) = x^4 + 2ax^3 + 5azx^2 + 4a3x$, and consequently $(x^3 + ax)^2 + 4a^2(x^2 + ax) = d$; which can be resolved by a quadratic equation.

Whence $x^2-6x=-4\pm\sqrt{(16+a)}$, by the common rule, And, by a second operation, $x=3\pm\sqrt{(9-4\pm\sqrt{(16+a)})}$. Therefore, by restoring the value of a, we have $x=3\pm\sqrt{(5\pm\sqrt{9025})}$. Or by extraction of roots, x=13, the Ans.

EXAMPLES FOR PRACTICE.*

1. Given $x^2 - 8x + 10 = 19$, to find the value of x. Aps. x = 9.

2. Given $x^2 = x - 40 = 170$, to find the value of x.

Ans. x=15.

3. Given $3x^2+9x-9=76$ to find the value of x.

3. Given $3x^2+2x-9=76$, to find the value of x. Ans. x=5

4. Given $\frac{1}{2}x^2 - \frac{1}{3}x + 7\frac{3}{6} = 8$, to find the value of x.

Ans. $x=1\frac{1}{2}$.

5. Given $\frac{1}{2}x - \frac{1}{3}\sqrt{x} = 22\frac{1}{6}$, to find the value of x.

Ans. x = 49.

6.† Given $x+\sqrt{(5x+10)}=8$, to find the value of x. Ans, x=3.

^{*} The unknown quantity in each of the following examples, as well as in those given above, has always two values, as appears from the common rule; but the negative and imaginary roots being, in general, but seldom used in practical questions of this kind, are here suppressed.

[†] Two values, of x are found, according to the process in resolving this question; that is, 18 and 3; but it appears, that 18 does not answer the condition of the equation, if we suppose that $\sqrt{(5x+10)}$ represents the positive square root of 5x+10. The reason is, that 5x+10 is the square of $-\sqrt{(5x+10)}$; thus, by squaring both sides of the equation $\sqrt{(5x+10)} = 8-x$, a new condition is introduced, and a new value of the unknown quantity corresponding to it, which had no place before. Here, 18 is the value which corresponds to the supposition that $x-\sqrt{(5x+10)}$

Esistes what is already said, it may be farther remarked, that one of the two positive roots of a quadratic equation of this kind, is also sometimes excluded by a condition in the question. Thus, if it were required to find three numbers in geometrical proportion, such that the sum of the first and second shall be 10, and the difference of the second and third 24.

It will be found, by putting x10-x and 34-x for the three numbers, that the values of x, in the resulting equation $x(34-x)=(10-x)^2$, as found by the usual rule, will be 25 and 2.

x.

7. Given $\sqrt{(10+x)-\sqrt[3]{(10+x)}}=2$, to find the value of x. Ans. x=6.

8. Given $2x^4-x^2+96=99$, to find the value of x.

Ans. $x=\frac{1}{2}\sqrt{6}$.

9. Given $x^6 + 20x^3 - 10 = 59$, to find the value of x.

10. Given $3x^{2^n}-2x^n+3=11$, to find the value of x.

Ans. $x=\sqrt[n]{2}$.

11. Given $5\sqrt[3]{x-3}\sqrt{x-1}\frac{1}{3}$, to find the value of x.

Ans. $3\frac{13}{81}$ or $\frac{1}{81}$.

12. Given $\frac{2}{3}x\sqrt{(3+2x^2)} = \frac{1}{2} + \frac{2}{3}x^2$, to find the value of

Ans. $x = \frac{1}{2}\sqrt{(-3+3\sqrt{2})}$.

13. Given $x\sqrt{(\frac{6}{x}-x)} = \frac{1+x^2}{\sqrt{x}}$ to find the value of x.

Ans. $x=(1+\frac{1}{2}\sqrt{2})^{\frac{1}{2}}$

14. Given $\frac{1}{x}\sqrt{(1-x^3)}=x^2$, to find the value of x.

Ans. $x = (\frac{1}{2}\sqrt{5-\frac{1}{2}})^{\frac{1}{3}}$.

15. Given $x\sqrt{(\frac{a}{x}-1)} = \sqrt{(x^2-b^2)}$, to find the value

of x. Ans. $x = \frac{1}{4}a + \frac{1}{4}\sqrt{(8b^2 + a^2)}$

16. Given $\sqrt{(1+x-x^2)} - 2(1+x-x^2) = \frac{1}{9}$ to find the

value of x. Ans. $x = \frac{1}{2} + \frac{1}{6} \sqrt{41}$.

But as the sum of the first and second of the required numbers is, by the question 10, it is manifest that 25 cannot be one of them; therefore the only proper root of the sum in this case is 2; and, consequently, the proportional sought are 2, 5, and 32.

17. Given $\sqrt{(x-\frac{1}{x})} + \sqrt{(1-\frac{1}{x})} = x$, to find the value of x.

Ans. $x = \frac{1}{2} + \frac{1}{2}\sqrt{5}$.

18. Given $x^{4^n} - 2x^3 + x^n = 6$, to find the value of x. Ans. $x = \sqrt[n]{(\frac{1}{2} + \frac{1}{2} \sqrt{13})}$.

19. Given $x^4 - 2x^3 + x = a$, to find the value of x.

Ans. $x = \frac{1}{2} \pm \sqrt{\frac{3}{4}} \pm \sqrt{(\alpha + \frac{1}{4})}$

When there are more equations and unknown quantities than one, a single equation, involving only one of the unknown quantities, may sometimes be obtained, by the rules before laid down for the solution of simple equations and, in this case, one of the unknown quantities being determined, the others may be found, by substituting its value in the remaining equations.

EXAMPLES.

1. Given $\begin{cases} x^2 + y^3 = 65 \\ xy = 28 \end{cases}$ to find the values of x and y.

Here, from the second equation, we have $y=\frac{28}{x}$; and by substituting this in the first $x^2 + \frac{784}{x^2} = 65$, or $x^4 - 65x^2 = -784$.

Whence, by the common rule before given, we have $x=\pm\sqrt{\frac{65}{2}}\pm\sqrt{\frac{4225}{4}-784}$.

Or, by reducing the parts under the last radical, and extracting the root $x=\pm\sqrt{\left(\frac{65}{2}\pm\frac{33}{2}\right)}=7$, or -1, and con-

sequently, $y = \frac{28}{7}$, or $-\frac{28}{4} = 4$ or -7.

Or the solution, in cases of this kind, may often be more readily obtained, by some of the artifices frequently made use of upon these occasions; which can only be learned from experience: thus, taking as before, (1.) $x^2 + y^2 = 65$,

(2.) xy=28, we shall have, as in the former method, by multiplying by 2, 2xy=56, and, by adding this equation to the first, and subtracting it from the same, $x^2+2xy+y^2=121$, and $x^2-2xy+y^2=9$. Whence, by extracting the square roots of each of these last equations, there will arise, $x+y=\pm 11$, and $x-y=\pm 3$, and, consequently by adding and subtracting these we shall have $2x=\pm 14$, or x=7, or x=7, and x=7, and x=7. It will also, sometimes facilitate the operation by substituting for one of the unknown quantities the product of the other, and a third unknown quantity which method may be applied with advantage, whenever the sum of the dimensions of the unknown quantities is the same in every term of the equation.

2. Given $\begin{cases} x^2 + xy = 56 \\ xy + 2y^2 = 60 \end{cases}$ to find the value of x and y.

Here, agreeably to the above observation, let x=vy, then $v^2y^2+vy^2=56$, and $vy^2+2y^2=60$, whence, from the first of these equations, $y^2=\frac{56}{v^2+v}$, and from the second $y^2=\frac{60}{v+2}$. Therefore, by equating the right hand member of these two expressions, we shall have $\frac{60}{v+2}=\frac{56}{v^2+v}$, or $60v^2+60v=56v+112$. And, by transposing 56v, and dividing the result by 60, $v^2+\frac{1}{15}v=\frac{28}{15}$. Hence by the common rule, for quadratics, we have $v=-\frac{1}{30}\pm\sqrt{\left(\frac{1}{900}+\frac{28}{15}\right)}=-\frac{1}{30}+\frac{41}{30}=\frac{4}{3}$. And, consequently, by the former part of the process, $y^2=\frac{60}{v+2}=\frac{60}{1\frac{1}{3}+2}=18$, or $y=\sqrt{(13)}=3\sqrt{2}$, and $x=vy=\frac{4}{3}\times 3\sqrt{2}=4\sqrt{2}$. The work may also be sometimes shortened, by substituting for the unknown quantities, the sum or difference of two other

quantities; which method may be used, when the unknown quantities, in each equation, are similarly involved

3. Given $\begin{cases} \frac{x^2}{y} + \frac{y^2}{x} = 18 \\ x + y = 12 \end{cases}$ to find the values of x and y.

Here, according to the above observation, let there be assumed x=z+v, and y=z-v. Then by adding these two equations together, we shall have x+y=2z=12, or z=6, also, since x=6+v, y=6-v, and by the first equation. $x^2+y^3=18xy$, we shall obtain, by substitution, $(6+v)^3+(6-v)^3=18(6+v)(6-v)$, or, by involving the two parts of the first member, and multiplying those of the second, $432+36v^2=648-18v^2$, whence, by transposition

 $54v^2 = 216$; and by division, $v^2 = \frac{216}{54} = 4$; or $v = \pm 2$.

And, therefore, by the first assumption, and the first part of the process, we have $x=z+v=6\pm 2=8$, or 4, and $y=z-v=6\pm 2=4$, or 8.

QUESTIONS PRODUCING QUADRATIC EQUATIONS.

The methods of expressing the conditions of questions of this kind, and the consequent reduction of them, till they are brought to a quadratic equation, involving only one unknown quantity and its square, are the same as those already given for simple equations.

1. To find two numbers such that their difference shall

be 8, and their product 240.

Let x equal the least number. Then will x + 8 = the greater.

And $x(x + 8) = x^2 + 8x = 240$, by the question, Whence $x = -4 + \sqrt{(16 + 240)} = -4 + \sqrt{256}$ by the common rule, before given,

Therefore x = 16-4 = 12, the less number, And x + 8 = 12 + 8 = 20, the greater. 2. It is required to divide the number 60 into two such parts, that their product shall be 864.

Let x = the greater part, Then will 60 - x = the less,

And $x(60-x)=60x-x^2=864$, by the question, Or by changing the signs on both sides of the equation $x^2-60x=-864$,

Whence $x=30\pm\sqrt{(900-864)}=30\pm\sqrt{36}=30\pm6$, by the rule.

And consequently x=30+6=36, or 30-6=24, the two parts sought.

 It is required to find two numbers such that their sum shall be 10(a), and the sum of their squares 58(b).
 Let x=the greater of the two numbers,

Then will a-x—the less,

And $x^2 + (a-x)^2 = 2x^2 - 2ax + a^2 = b$, by the question, Or $2x^2 - 2ax = b - a^2$, by transposition,

And $x^2 - ax = \frac{b - a^2}{a}$, by division.

Whence
$$x = \frac{a}{2} \pm \sqrt{(\frac{a}{4} + \frac{b - a^2}{2})} = \frac{a}{2} \pm \frac{1}{2} \sqrt{(2b - a^2)}$$

by the rule, And if 10 be put for a, and 58 for b, we shall have

 $x = \frac{10}{2} + \frac{1}{2} \checkmark (116 - 100) = 7$, the greater number,

And
$$10-x=\frac{10}{2}-\frac{1}{2}$$
 (116-100)=3, the less.

4. Having sold a piece of cloth for 24l., I gained as much per cent as it cost me; what was the price of the cloth?

Let x= pounds the cloth cost, Then will 24-x= the whole gain, But 100:x::x:24-x, by the question. Or $x^2=100(24-x)=2400-100x$, That is, $x^2+100x=2400$, Whence $x=-50+\sqrt{(2500+2400)}=-50+70=20$ by the rule, And consequently 201 .= price of the cloth.

5. A person bought a number of sheep for 801., and if he had bought 4 more for the same money, he would have paid 11. less for each; how many did he buy?

Let x represent the number of sheep,

Then will $\frac{80}{x}$ be the price of each,

And $\frac{80}{x+4}$ = price of each, if x+4 cost 80l.

But $\frac{80}{x} = \frac{30}{x+4} + 1$, by the question, Or $80 = \frac{80x}{x+4} + x$, by multiplication.

And $80x+320=80x+x^2+4x$. by the same, Or, by leaving out 80x on each side, $x^2 + 4x = 320$, Whence $x=-2+\sqrt{(4+320)}=-2+18$, by the rule, And consequently x=16, the number of sheep.

6. It is required to find two numbers, such that their sum, product, and difference of their squares, shall be all equal to each other.

Let
$$x$$
=the greater number, and y = the less.
Then
$$\begin{cases} x+y=xy\\ x+y=x^2-y^2 \end{cases}$$
 by the question.

Hence $1 = \frac{x^2 - y^2}{x + y} = x - y$, or x = y + 1, by 2d equation. And (y+1)+y=y(y+1) by 1st equation, That is, $2y+1=y^2+y$; or $y^2-y=1$,

Whence $y = \frac{1}{2} + \sqrt{(\frac{1}{4} + 1)} = \frac{1}{6} + \frac{1}{6} \checkmark 5$, by the rule,

Therefore
$$y = \frac{1}{2} + \frac{1}{2} \sqrt{5} = 1.6180$$
 . . . And $x = y + 1 = \frac{3}{2} + \frac{1}{2} \sqrt{5} = 2.6180$. . .

Where . . . denotes that the decimal does not end.

7. It is required to find four numbers in arithmetical progression, such that the product of the two extremes hall be 45, and the product of the means 77.

Let x = least extreme, and y = common difference, Then x, x+y, x+2y, and x+3y, will be the four numbers Hence $\begin{cases} x(x+3y) = x^2 + 3xy = 45\\ (x+y)(x+2y) = x^2 + 3xy + 2y^2 = 77 \end{cases}$ by the question,

And $2y^2 = 77 - 45 = 32$, by subtraction, Or $y^2 = \frac{32}{2} = 16$ by division, and $y = \sqrt{16} = 4$,

Therefore $x^2+3xy=x^2+12x=45$, by the 1st equation And consequently $a=-6+\sqrt{(36+45)}=-6+9=3$, by the rule,

Whence the numbers are 3, 7, 11, and 15.

3. It is required to find three numbers in geometrics progression, such that their sum shall be 14, and the sum of their squares 84.

Let x, y, and z be the three numbers, Then $xz=y^2$, by the nature of proportion,

And $\begin{cases} x+y+z=14\\ x^2+y^2+z^2=84 \end{cases}$ by the question, Hence x+z=14-y, by the second equation,

And $x^2+2zx+z^2=196-28y+y^2$, by squaring bot sides,

Or $x^2+z^2+2y^2=196-28y+y^2$ by putting $2y^2$ for it equal 2xz,

That is $x^2 + y^2 + z^2 = 196 - 28y$ by subtraction, Or 196 - 28y = 84 by equality,

Hence $y = \frac{196 - 84}{98} = 4$, by transposition and division,

Again $xz=y^2=16$, or $x=\frac{16}{z}$, by the 1st. equation,

And $x+y+z=\frac{16}{z}+4+z=14$, by the 2d equation,

Or $16+4z+z^2=14z$, or $z^2-10z=-16$, Whence $z=5\pm\sqrt{(25-16)}=5\pm3=8$, or 2 by the rule

Therefore the three numbers are 2, 4, and 8.

9. It is required to find two numbers, such that thei sum shall be 13(a), and the sum of their fourth power 4721 (b).

Let x= the difference of the two numbers sought,

Then will $\frac{1}{2}a + \frac{1}{2}x$, or $\frac{a+x}{2}$ = the greater number,

And
$$\frac{1}{2}a - \frac{1}{2}x$$
, or $\frac{a-x}{2}$ = the less,
 $(a+x)^4$, $(a-x)^4$

But
$$\frac{(a+x)^4}{16} + \frac{(a-x)^4}{16} = b$$
, by the question,

Or $(a+x)^4+(a-x)^4=16b$, by multiplication, or $2a^4+12a^2x^2+2x^4=16b$, by involution and addition,

and $x^4 + 6a^2x^2 = 8b - a^4$, by transposition and division, Whence $x^2 = -3a^2 + \sqrt{(9a^4 + 8b - a^4)} = -3a^2 +$

 $\sqrt{8(a^4+b)}$, by the rule,

And $x=\sqrt{-3a^2+2\sqrt{2(a^4+b)}}$, by extracting the root, Where, by substituting 13 for a, and 4721 for b.

we shall have x=3,

Therefore $\frac{13+x}{2} = \frac{16}{2} = 8$, the greater number,

And $\frac{13-x}{2} = \frac{10}{2} = 5$, the less number,

The sum of which is 13, and $8^4 + 5^4 = 4721$.

10. Given the sum of two numbers equal s, and their product = p, to find the sum of their squares, cubes, biquadrates, &c.

Let x and y denote the two numbers; then

(1.) x+y=s, (2.) xy=p. From the square of the first of these equations take wice the second, and we shall have

(3.) $x^2 + y^2 = s^2 - 2p = \text{sum of the squares}$.

Multiply this by the 1st equation, and the product will be $v^3+xy^2+x^2y+y^3=s^3-2sp$.

From which subtract the product of the first and second

equations, and there will remain

(4.) $x^3+y^3=s^3-3sp=$ sum of the cubes. Multiply this likewise by the 1st, and the product will be $t^4+xy^3+x^3y+y^4=s^4-3s^2p$; from which subtract the

 $x^4 + xy^3 + x^3y + y^4 = s^4 - 3s^2p$; from which subtract the product of the second and third equations, and there will remain

(5.) $x^4 + y^4 = s^4 - 4s^2p + 2p^2 = \text{sum of the biquadrates}$. And, again multiplying this by the 1st equation, and subtracting from the result the product of the second and fourth, we shall have

(6.) $x^5 + y^5 = s^5 - 5s^3p + 5sp^2 = \text{sum of the fifth powers.}$ And so on; the expression for the sum of any powers in general being $x^m + y^m = s^m - ms^{m-2}p + \frac{m(m-3)}{2}s^{m-4}p^2 -$

$$\frac{n(m-4)(m-5)}{2 \cdot 3} s^{m-6} p^3 + \frac{m(m-5)(m-6)(m-7)}{2 \cdot 3 \cdot 4} s^{m-8} p^4 - ,$$

&c.

Where it is evident that the series will terminate when the index of s becomes = o.

QUESTIONS FOR PRACTICE.

1. It is required to divide the number 40 into two such parts, that the sum of their squares shall be 818.

Ans. 23 and 17

2. To find a number such, that if you subtract it from 10, and then multiply the remainder by the number itself, the product shall be 21. Ans. 7 or 3

3. It is required to divide the number 24 into two such parts, that their product shall be equal to 35 times their

Ans. 10 and 14 4. It is required to divide a line, of 20 inches in length, into two such parts that the rectangle of the whole and one of the parts shall be equal to the square of the other.

Ans. 10 15 - 10

5. It is required to divide the number 60 into two such parts, that their product shall be to the sum of their squares in the ratio of 2 to 5. Ans. 20 and 40

6. It is required to divide the number 146 into such two parts, that the difference of their square roots shall be 6.

Ans. 25 and 121

7. What two numbers are those whose sum is 20 and their product 36? Ans. 2 and 18

8. The sum of two numbers is 11, and the sum of their reciprocals 31; required the numbers. Ans. 1 and 5

9. The difference of two numbers is 15, and half their product is equal to the cube of the less number; required the numbers.

Ans. 3 and 13

the numbers.

Ans. 3 and 13

10. The difference of two numbers is 5, and the differ-

ence of their cubes 1685; required the numbers.

Ans. 8 and 13

11. A person bought cloth for 33l. 15s. which he sold again at 2l. 8s. per piece, and gained by the bargain as much as one piece cost him; required the number of pieces.

Ans. 15

12. What two numbers are those, whose sum, multi-

plied by the greater, is equal to 77, and whose difference, multiplied by the less, is equal to 12.

Ans. 4 and 7

13. A glazier bought as many sheep as cost bim 60l., and after reserving 15 out of the number, sold the remainder for 54l., and gained 2s. a head by them: how many sheep did he buy?

Ans. 75

14. It is required to find two numbers, such that their product shall be equal to the difference of their squares, and the sum of their squares equal to the difference of their cubes.

Ans. $\frac{1}{2}\sqrt{5}$ and $\frac{1}{4}(5+\sqrt{5})$

15. The difference of two numbers is 8, and the difference of their fourth powers is 14560; required the numbers.*

Ans. 3 and 11

RULE.

Whenever, in the cubic equation of the form $x_0 + ax = b$; b can be divided into two factors, m and n so that, m2 + a = n, then the cubic can be resolved by a quadratic: thus, let the equation be $x^3 + 6x = 20$; now let 20 be divided into two factors 10 and 2; it is evident that 22; or 4 added to 6, the coefficient of x, is equal 10, the other factor. And, consequently, by multiplying by x, the equation becomes $x^4 + 6x^2 = 20x$, or $x^4 + 6x^2 = 10 \times 2x$; then squaring 2x and adding this square to both sides we get $x + 16x^2 = (2x)^2 + 10$ (2x), by completing the square $x^4 + 10x^2 + 25 = (2x)^2 + 10$ (2x) + 25, and extracting the square $x^4 + 5x^2 + 5x^2$

^{*} In the solution of this question, the process brings out the answer in the form a3+ax=b; which is a cubic equation, and, therefore cannot be resolved by the ordinary rules for quadratics; but we can sometimes reduce such equations to the form of a quadratic, and then resolve it according to the rules already given.

- 16. A company at a tavern had 8l. 15s. to pay for their reckoning; but, before the bill was settled, two of them went away; in consequence of which those who remained had 10s. apiece more to pay than before: how many were there in company?

 Ans. 7
- 17. A person ordered 7l. 4s. to be distributed among some poor people; but, before the money was divided, there came in, unexpectedly, two claimants more, by which means the former received a shilling a piece less than they would otherwise have done; what was their number at first?

 Ans. 16 persons

13. It is required to find four numbers in geometrical progression such, that their sum shall be 15, and the sum of their squares 85.

Ans. 1, 2, 4, and 3

19. The sum of two numbers is 11, and the sum of

their fifth powers is 17831; required the numbers?

20. It is required to find four numbers in arithmetical progression such, that their common difference shall be 4,

and their continued product 176985.

Aus 15, 19, 23, and 27

21. Two detachments of foot being ordered to a station at the distance of 39 miles from their present quarters, begin their march at the same time; but one party, by travelling \(\frac{1}{2}\) of a mile an hour faster than the other, arrive there an hour sooner; required their rates of marching?

Ans. 3\(\frac{1}{2}\) and 3 miles per hour

22. It is required to find two numbers such that the square of the first plus their product, shall be 140, and the square of the second minus their product 78.

Ans. 7 and 13.

23. It is required to find two numbers, such that their difference shall be $13\frac{65}{210}$, and the difference of their cube roots $1\frac{1}{2}$.

Ans. $15\frac{5}{2}$, and $2\frac{1}{2}$

24. It is required to find three numbers in arithmetical progression, such that the sum of their squares shall be 93; and if the first be multiplied by 3, the second by 4, and the third by 5, the sum of the products shall be 66.

Ans. 2, 5, and 8.

25. The sum of three numbers in harmonical propor-

tion is 191, and the product of the first and third is 4032; required the numbers.

Ans. 72, 63, and 56.

26. It is required to find four numbers in arithmetical progression, such that if they are increased by 2, 4, 8 and 15 respectively, the sums shall be in geometrical progression.

Ans. 6, 8, 10 and 12.

27. It is required to find two numbers, such, that if their difference be multiplied into their sum, the product will be 5; but if the difference of their squares be multiplied into the sum of their squares the product will be 65.

Ans. 3 and 2

23. It is required to divide the number 10 into two such parts, that if the square root of the greater part be taken from the greater part, the remainder shall be equal to the square root of the less part added to the less part.

Ans. $5+\frac{1}{2}\sqrt{19}$ and $5-\frac{1}{2}\sqrt{19}$. 29. It is required to find two numbers, such that if their product be added to their sum it shall make 61, and if their sum be taken from the sum of their squares it shall leave 88. Ans. $7+\sqrt{2}$ and $7-\sqrt{2}$.

30. It is required to find two numbers, such that their difference multiplied by the difference of their squares shall be 576, and their sum multiplied by the sum of their squares shall be 2336.

Ans. 5 and 11.

31. It is required to find three numbers in continual proportion, whose sum shall be 20, and the sum of their squares 140. Ans. $6\frac{3}{4} + \sqrt{3}\frac{5}{16}$, $6\frac{1}{2}$, and $6\frac{3}{4} - \sqrt{3}\frac{5}{16}$.

32. It is required to find two numbers whose product

32. It is required to find two numbers whose product shall be 320, and the difference of their cubes to the cube of their difference, as 61 is to unity.

Ans. 20 and 16.

33. The sum of 700 dollars was divided among four persons, A, B, G and D, whose shares were in geometrical progression; and the difference between the greatest and least, was to the difference between the two means, as 37 to 12. What were all the several shares?

Ans. 108, 144, 192, and 256 Dollars.

OF CUBIC EQUATIONS.

A cubic equation is that in which the unknown quantity rises to three dimensions; and like quadratics, or those of the higher orders, is either simple or compound.

A simple cubic equation is of the form

$$ax^3 = b$$
, or $x^3 = \frac{b}{a}$; where $x = \sqrt[3]{\frac{b}{a}}$

A compound cubic equation is of the form $x^3+ax=b$, $x^3+ax^2=b$, or $x^3+ax^2+bx=c$, in each of which, the known quantities a, b, c, may be either + or -.

Or, either of the two latter of these equations may be reduced to the same form as the first, by taking away its second term; which is done as follows:

RULE.

Take some new unknown quantity, and subjoin to it a third part of the coefficient of the second term of the equation with its sign changed; then if this sum, or difference as it may happen to be, be substituted for the original unknown quantity and its powers, in the proposed equation, there will arise an equation wanting its second term.

Note. The second term of any of the higher orders of equations may also be exterminated in a similar manner, by substituting for the unknown quantity some other unknown quantity, and the 4th, 5th, &c. part of the coefficient of its second term, with the sign changed, according as the equation is of the 4th, 5th, &c. power.*

^{*} Equations may be transformed into a variety of other new equations; the principal of which are as follows:

^{1.} The equation x4-4x3-19x2+106x-120=0, the roots of which are 2, 3, 4, and -5; by changing the signs of the second and fourth terms, becomes x4+4x3-19x2-106x-120=0, the roots of which are 5, -2, -3, and -4.

EXAMPLES.

1. It is required to exterminate the second term of the equation $x^3 + 3ax^2 = b$, or $x^3 + 3ax^2 - b = 0$.

Here
$$x=z-\frac{3a}{3}=z-a$$
,
 $x^3=z^3-3az^2+3a^2z-a^3$
Then
$$\begin{cases} 3ax^2=+3az^2-6a^2z+3a^3\\-b=-b \end{cases}$$

Whence
$$z^3 - 3a^2z + 2a^3 - b = 0$$
,
Or $z^3 - 3a^2z = b - 2a^3$,

in which equation the second power (z^2) , of the unknown quantity, is wanting.

2. Let the equation $x^3-12x^2+3x=-16$, be transformed into another, that shall want the second term.

Then
$$\begin{cases} (z+4)^3 = z^3 + 12z^2 + 48z + 64 \\ -12(z+4)^2 = -12z^2 - 96z - 192 \\ +3(z+4) = +3z + 12 \end{cases}$$

Whence
$$z^3 - 45z - 116 = -16$$

Or $z^3 - 45z = 100^{-1}$

which is an equation where z2; or the second term, is wanting, as before.

^{2.} The equation x3 + x2 - 10x + 3 = 0, is transformed, by assuming x = y 4, into y3 - 11y2 + 30y = 0, or y2 - 11y + 30 = 0; the roots of which are

greater than those of the former by 4.

3. The equation x3 - 6x2 + 9x - 1 = 0, may be transformed into one which shall want the third term, by assuming x = y + e, and in the resulting equation, let $3e^2 - 12e + 9$, or $e^2 - 4e + 3 = 0$, in which the values of e are 1 and 3; then assume z = y + 3, or y + 1, and the resulting equation will be y3 + 3y2 - 1 = 0, an equation wanting the third term.

^{4.} The equation $6x^3-11x^2+6x-1=0$ by assuming $x=\frac{1}{y}$, may be tansformed into $y^3-6y^2+11y-6=0$; the roots of which are the reciprocals of

the former.

5. The equation $3x^3 - 13x^2 + 14x + 16 = 0$ by assuming $x = \frac{1}{3}$, into $y^3 - \frac{1}{3}x^2 + \frac{1}{4}x + \frac{1}{4}x$

3. Let the equation $x^3 - 6x^2 = 10$, be transformed into another that shall want the second term.

Ans. $y^3-15y^2+81y=243$, be transformed into an equation that shall want the second term.

Ans. $x^3 + 6x = 88$

5. Let the equation $x^3 + \frac{3}{4}x^2 + \frac{7}{8}x - \frac{9}{16} = 0$, be trans-

formed into another, that shall want the second term.

Ans. $y^3 + \frac{11}{16}y = \frac{3}{4}$

6. Let the equation $x^4 + 8x^3 - 5x^2 + 10x - 4 = 0$, be transformed into another, that shall want the second term.

Ans. $y^4 - 29y^2 + 94y - 92 = 0$. 7. Let the equation $x^4 - 3x^3 + 3x^2 - 5x - 2 = 0$, be transformed into another that shall want the third term.

Ans. $y^4 + y^3 - 4y - 2 = 0$.

8. Let the equation $3x^3-2x+1=0$, be transformed into another, whose roots are the reciprocals of the former.

Ans. $y^3 - 2y^2 + 3 = 0$.

9. Let the equation $x^4 - \frac{1}{2}x^3 + \frac{1}{3}x^2 - \frac{3}{3}x + \frac{7}{13} = 0$, be transformed into another, in which the coefficient of the highest term shall be unity, and the remaining terms integers.

Ans. $y^4 - 3y^3 + 12y^2 - 162y + 72 = 0$.

OF THE SOLUTION OF CUBIC EQUATIONS.

RULE.

Take away the second term of the equation when necessary, as directed in the preceding rule. Then, if the numeral coefficients of the given equation, or of that arising from the reduction above mentioned, be substituted for a and b in either of the following formulæ, the result will give one of the roots, as required*.

^{*} If, instead of the regular method of reducing a cubic equation of the general form.

$$x = \begin{cases} \sqrt[3]{\left\{\frac{b}{2} + \sqrt{\left(\frac{b^3}{4} + \frac{a^3}{27}\right)}\right\} + \sqrt[3]{\left\{\frac{b}{2} - \sqrt{\left(\frac{b^2}{4} + \frac{a^3}{27}\right)}\right\}}} & \text{or} \\ \sqrt[3]{\left\{\frac{b}{2} + \sqrt{\left(\frac{b^2}{4} + \frac{a^3}{27}\right)}\right\} - \frac{\frac{1}{3}a}{\sqrt[3]{\left\{\frac{b}{2} + \sqrt{\left(\frac{b^2}{4} + \frac{a^3}{27}\right)}\right\}}} \end{cases}$$

Where it is to be observed, that when the coefficient a, of the second term of the above equation, is negative, $\frac{a^3}{27}$, as also $\frac{a}{3}$, in the formula, will be negative; and if the

absolute b be negative, $\frac{b}{2}$ in the formula, will, also, be negative; but $\frac{b^2}{4}$ will be positive.*

 $x^3 + ax + bx + c = 0$.

* This method of solving cubic equations is usually ascribed to Caydan, a celebrated Italian analyst of the 16th century; but the authors of it were Scipio Ferreus, and Nicolas Tartalea, who discovered it about the same time, independently of each other, as is proved by Montucla, in his Histoire des Mathematiques, Vol. I. p. 568, and more at large in Hutton's Mathe-

matical Dictionary, Art. Algebra.

The rule above given, which is similar to that of Cardan, may be demonstrated as follows:

Let the equation, whose root is required, be x3 + ax=b.

It may, likewise be remarked, that when the equation is of the form

$$x^3-ax=\pm b$$

and $\frac{a^3}{27}$ is greater than $\frac{b^2}{4}$, or $4a^3$ greater than 27b, the solution of it cannot be obtained by the above rule; as the question, in this instance, falls under what is usually called the Irreducible Case of cubic equations.*

And if, from the square of this last equation, there be taken 4 times the cube of the equation $yz=-\frac{1}{3}a$, we shall have y6-2y3z3+z6=b2+Aga2, or

 $y_3 - z_3 = \sqrt{(b_2 + \frac{4}{27}a_3)}$ But the sum of this equation and $y_3 + z_3 = b$, is $2y_3 = b + \sqrt{(b_2 + \frac{4}{27}a_3)}$ $\frac{4}{27}a^3$) and their difference is $2z^3 = b - \sqrt{(b^2 + \frac{4}{27}a^3)}$; whence $y = \sqrt[3]{(\frac{1}{2}a^3)}$ $b + \checkmark (\frac{1}{4}b^2 + \frac{1}{27}\pi^3)), \text{ and } z = \sqrt[3]{\left(\frac{1}{2}b - \checkmark\left(\frac{1}{4}b^2 + \frac{1}{27}\pi^3\right)\right)}.$ From which it appears, that y+z, or its equal x, is = $\sqrt[3]{\left(\frac{1}{2}b + \checkmark\left(\frac{1}{4}b^2 + \frac{1}{27}\pi^3\right)\right) + \sqrt[3]{\left(\frac{1}{2}b - \checkmark\left(\frac{1}{4}b^2 + \frac{1}{27}\pi^3\right)\right)}, \text{ which is the theo-}$

rem;

Or, since z is
$$= -\frac{a}{3y}$$
, it will be $y + z = y - \frac{a}{3y}$, or $x = \frac{2}{3}\sqrt{(\frac{1}{2}b + \sqrt{(\frac{1}{2}b^2 + \frac{1}{27}a^3)})} - \frac{\frac{1}{3}a}{\sqrt[3]{(\frac{1}{2}b + \sqrt{(\frac{1}{2}b^2 + \frac{1}{27}a^3)})}}$, the same as the

* It may here be farther observed as a remarkable circumstance in the history of this science, that the solution of the Irreducible Case above mentioned, except by means of a table of sines, or by infinite series, has hitherto baffled the united efforts of the most celebrated mathematicians in Europe; although it is well known that all the three roots of the equation are, in this case, real; whereas, in those that are resolvable by the above formula, only one of the roots is real, so that in fact, the rule is only applicable to such cubics, as have two equal, or two impossible roots.

The reason why the assumptions, inade in the note to the former part of this article with respect to the solution of the equation $x^3 - ax = b$, are found to fail in the case in question (and it does not appear that any other can be adopted) is, that the two auxiliary equations 3yz = -a and y3 + z3 = b, which in this case, become 3yz = a, and y3 + z3 = b, or y3z3 = a

 $\frac{z}{27}$, and $y^3 + z^2 = b$, cannot take place together; being inconsistent with

For the greatest product that can be formed of the two quantities 43 + 23 is, when they are all equal to each other; or since y3 +23=b, when each of these = $\frac{1}{2}b$; in which case their product is= $\frac{1}{2}b^2$.

But, as above shown $y \cdot z^2 = \frac{a^3}{27}$, by the question, therefore when $\frac{a^3}{27} > \frac{b^2}{4}$ the two conditions are incompatible with each other; and consequently the solution of the problem, upon that supposition, can only be obtained by imaginary quantities.

EXAMPLES.

Given $2x^3 - 12x^2 + 36x = 44$, to find the value of x. Here $x^3 - 6x^2 - 18x = 22$, by dividing by 2.

And, in order to exterminate the second term.

Put
$$x=z+\frac{6}{3}=z+2$$
,

$$\begin{vmatrix} (z+2)^3=z^3+6z^3+12z+8\\ -6(z+2)^2=-6z^3-24z-24\\ 18(z+2)= & 18z+36 \end{vmatrix} = 22$$

Whence $z^3+6z+20=22$, or $z^3+6z=2$, And, consequently, by substituting 6 for a, and 2 for b, in

the first formula, we shall have,

$$x = \sqrt[3]{\frac{2}{2} + \sqrt{\frac{4}{4} + \frac{216}{27}}} + \sqrt[3]{\frac{2}{2} - \sqrt{\frac{4}{4} + \frac{216}{27}}} = \sqrt[3]{(1 + \sqrt{(1+8)}) + \sqrt[3]{(1-\sqrt{(1+8)})}} + \sqrt[3]{(1-\sqrt{9})} + \sqrt[3]{(1+3) + \sqrt[3]{(1-3)}} + \sqrt[3]{4 - \sqrt[3]{2}},$$
Therefore $x = z + 2 = \sqrt[3]{4 - 2} + 2 + 1.587401 - 1.259921 = 2.32748$, the answer.

2. Given $x^3 - 6x = 12$, to find the value of x.

Here a being equal to -6, and b equal to 12, we shall have, by the formula.

$$x = \sqrt[3]{(6+\sqrt{36-8})} - \frac{-2}{\sqrt[3]{(6+\sqrt{36-8})}} = \frac{2}{\sqrt[3]{(6+\sqrt{28})} + \frac{2}{\sqrt[3]{(6+\sqrt{28})}} - \sqrt[3]{(6+5.2915)} + \frac{2}{\sqrt[3]{(6+5.2915)}} = \sqrt[3]{(11.2915)} + \frac{2}{\sqrt[3]{(11.2915)}} = 2.2435 + \frac{2}{2.2435} = 2.2435 + .8957 = 3.1392$$
Therefore $x = 3.1392$, the answer.

Therefore x=3.1392, the answer.

3. Given $x^3 - 2x = -4$, to find the value of x. Here a being =-2, and b=-4, we shall have, by the formula.

$$x=\sqrt[3]{\left\{-2+\sqrt{(4-\frac{8}{27})}\right\}}+\sqrt[3]{\left\{-2-\sqrt{(4-\frac{8}{27})}\right\}}$$
, or

by reduction,
$$\sqrt{(-2+\frac{10}{9}\sqrt{3})} - \sqrt[3]{(2+\frac{10}{9}\sqrt{3})} = \sqrt[3]{(-2+1.9245)} - \sqrt[3]{(2+1.9245)} = \sqrt[3]{(-.0755 - \sqrt{3}.9245)} = -.4226 - 1.5773 = -1.9999$$
, or -2

Therefore $x = -2$, the answer.*

Note. When one of the roots of a cubic equation has been found, by the common formula as above, or in any other way, the other two roots may be determined, as fol-

lows:

Let the known root be denoted by r, and put all the terms of the equation, when brought to the left hand side, =0; then if the equation, so formed, be divided by $x\pm r$, according as r is positive or negative, there will arise a quadratic equation, the roots of which will be the other two roots of the given cubic equation.

4. Given $x^3-15x=4$, to find the three roots, or values of x.

Here x is readily found, by a few trials, to be equal to 4, and therefore

$$\begin{array}{r}
x-4)x^3 - 15x - 4(x^2 + 4x + 1) \\
x^3 - 4x^2 \\
\hline
4x^2 - 15x \\
4x^2 - 16x \\
\hline
x - 4 \\
x - 4
\end{array}$$

down in Surds, Case 12, the result will be found equal to -2 as it ought.

When the root of the given equation is a whole number, this method only determines it by an approximation of 9s, in the decimal part, which sufficiently indicates the entire integer; but in most instances of this kind, its value may be more readily found, by a few trials, from the equation itself.

Or if, as in the above example, the roots, or numeral values of $\sqrt[3]{(-2+\frac{10}{9})}$, and $-\sqrt[3]{(2+\frac{10}{9})}$ be determined according to the rule laid

Whence, according to the note above given, $x^2+4x+1=0$, or $x^2+4x=-1$;

the two roots of which quadratic are $-2+\sqrt{3}$ and -2-√3; and consequently

4, -2+ 13, and -2-13,

are the three roots of the proposed equation. Or, putting a=-15 and b=4, we shall have,

or, putting
$$a = -15$$
 and $b = 4$, we shall have,

$$d = \sqrt[3]{\left\{\frac{b}{2} + \sqrt{\left(\frac{b^2}{4} - \frac{a^3}{27}\right)}\right\}} = \sqrt[3]{\left\{2 + \sqrt{(4 - 12b)}\right\}}$$

$$\sqrt[3]{(2+\sqrt{-121})} = 2+\sqrt{-1},$$
and $c = \sqrt[3]{\frac{b}{2}} - \sqrt{(\frac{b^2}{4} - \frac{a^3}{27})} = \sqrt[3]{(2-\sqrt{(4-125)})} =$

$$\frac{3}{2}(2-3/-121)=2-3/-1$$

 $\sqrt[3]{(2-\sqrt{-121})} = 2 - \sqrt{-1},$ as will be found either by cubing $2 + \sqrt{-1}$ and $2 - \sqrt{-1}$, or by the rule given in case 12 surds.

Whence
$$a+c=2+\sqrt{-1+2}-\sqrt{-1}=4$$
, $-\frac{1}{2}(d+c)+\frac{1}{2}(d-c)\sqrt{-3}=-2+\sqrt{-1}\times\sqrt{-3}=-2+\sqrt{3}$,

$$-\frac{1}{2}(d+c) - \frac{1}{2}(d-c)\sqrt{-3} = -\frac{2}{2}-\sqrt{-1}\times\sqrt{-3} = -\frac{2}{2}$$

and consequently 4, $-2+\sqrt{3}$, and $-2-\sqrt{3}$ are the three roots of the equation, as before found.

EXAMPLES FOR PRACTICE.

- 1. Given $x^3 + 3x^2 6x = 8$, to find the root of the equation, or the value of x. Ans. x=2.
- 2. Given $x^3 + x^2 = 500$, to find the root of the equation. or the value of x. Ans. x = 7.616789.
- 3. Given $x^3 3x^2 = 5$, to find the root of the equation. or the value of x. Ans. x = 3.103803.
- 4. Given $x^3 6x = 6$, to find the root of the equation, or the value of x. Ans. 3/4+3/2.
- 5. Given $x^3 + 9x = 6$, to find the root of the equation, or the value of x. Ans. 3/9-3/3.
- 6. Given $x^3 + 2x^2 23x = 70$, to find the root of the equation, or the value of x. Ans. x = 5.134899.
- 7. Given $x^3 17x^2 + 54x = 350$, to find the root of the equation, or the value of x. Ans. x = 14.954068.

8. Given $x^3-6x=4$, to find the three roots of the equation, or the three values of x.

Ans. -2, $1+\sqrt{3}$, and $1-\sqrt{3}$.

9. Given $x^3 - 5x^2 + 2x = -12$, to find the three roots of the equation, or the three values of x.

Ans. -3, 1+15, and 1-15.

OF THE

SOLUTION OF CUBIC EQUATIONS

BY

CONVERGING SERIES.

This method, which, in some cases, will be found more convenient in practice than either of the former, consists in substituting the numeral parts of the given equation, in the place of the literal, in one of the following general formulæ, accorling to which it may be found to belong, and then collecting as many terms of the series as are sufficient for determining the value of the unknown quantity, to the degree of exactness required.*

1. $x^3 + ax = b.1$

$$x = \sqrt[3]{\left\{\sqrt{\left(\frac{1}{4}b^2 + \frac{1}{27}a^3\right) + \frac{1}{2}b}\right\}} - \sqrt[3]{\left\{\sqrt{\left(\frac{1}{4}b^2 + \frac{1}{27}a^3\right) - \frac{1}{2}b}\right\}}; \text{ or, by putting, for the sake of greater simplicity, } \sqrt{\left(\frac{1}{4}b^2 + \frac{1}{27}a^3\right)}$$

$$\Rightarrow^{q}, \text{ and reducing the expression, } x = s^{\frac{1}{3}}\left\{\sqrt{\left(1 + \frac{b}{2s}\right)} - \sqrt[3]{\left(1 - \frac{b}{2s}\right)}\right\}$$

^{*} The method laid down in this article, of solving cubic equation, by means of series was first given by Nicole, in the Memoirs of the Academy of Sciences, an. 1738, p. 99; and afterwards at greater length, by Clairaur in his Elemens d'Algebra.

[†] With respect to the determination of the roots of cubic equations by means of series, let there be given, as above, the equation $x^3 + ax = b$, where the root by transposing the terms of each of the two branches of the common formula. is

$$\begin{array}{c} x = \frac{2b}{\sqrt[3]{(2(27b^3 + 4a^3))}} \left\{ 1 + \frac{2.5}{6.9} \left(\frac{27b^2}{27b^2 + 4a^3} \right) + \frac{2.5.8.11}{6.9.12.15} \right. \\ \left(\frac{27b^2}{27b^2 + 4a^3} \right)^2 + \frac{2.5.8.11.14.17}{6.9.12.15.18.21} \left(\frac{27b^2}{27b^2 + 4a^3} \right)^3 + \&c. \right\} \cdot Or, \\ x = \frac{2b}{\sqrt[3]{(2(27b^2 + 4a^3))}} \left\{ 1 + \frac{2.5}{6.9} \left(\frac{27b^2}{27b^2 + 4a^3} \right)^4 + \frac{8.11}{12.15} \right. \\ \left(\frac{27b^2}{27b^2 + 4a^3} \right)^3 + \frac{14.17}{18.21} \left(\frac{27b^2}{27b^2 + 4a^3} \right)^2 + \frac{20.23}{24.27} \\ \left(\frac{27b^2}{27b^2 + 4a^3} \right)^3 + \&c. \right\} \cdot Or, \\ \left(\frac{27b^2}{27b^2 + 4a^3} \right) + \frac{14.17}{18.21} \left(\frac{27b^2}{27b^2 + 4a^3} \right)^2 + \frac{20.23}{24.27} \right. \\ \left(\frac{27b^2}{27b^2 + 4a^3} \right)^3 + \&c. \right\} \cdot Or,$$

In which case, as well as in all the following ones, A, E, c, &c. denote the terms immediately preceding those in which they are first found.

2. $x^3 - ax = \pm b$, where $\frac{1}{4}b^2$ is supposed to be greater than $\frac{1}{24}a^3$, or $27b^2 > 4a^3$.

$$x = \pm 2\sqrt[3]{\frac{b}{2}} \left\{ 1 - \frac{2}{3.6} \left(\frac{27b^2 - 4a^3}{27b^2} \right) - \frac{2.5.8}{3.6.9.12} \left(\frac{27b^2 - 4a^3}{27b^2} \right)^3 - \frac{2.5.8}{3.6.9.12.15.18} \left(\frac{27b^2 - 4a^3}{27b^3} \right)^3 - \&c. \right\}^*. \text{ Or,}$$

Hence, extracting the roots of the right hand member of this equation, by the binomial theorem, there will arise $\sqrt[3]{(1+\frac{b}{2s})} = 1 + \frac{1}{3}(\frac{b}{2s}) - \frac{2}{3.6}$

$$(\frac{b}{2s})^2 + \frac{2 \cdot 5}{3 \cdot 6 \cdot 9} (\frac{b}{s})^3 - \frac{2 \cdot 5 \cdot 8}{3 \cdot 6 \cdot 9 \cdot 12} (\frac{b}{2s})^4 + &c.$$

$$\sqrt[3]{(1 - \frac{b}{2s})} = 1 - \frac{1}{3} (\frac{b}{2s}) - \frac{2}{3 \cdot 6} (\frac{b}{2s})^2 - \frac{2 \cdot 5}{3 \cdot 6 \cdot 9} (\frac{b}{2s})^3 - \frac{2 \cdot 5 \cdot 8}{3 \cdot 6 \cdot 9 \cdot 12}$$

$$(\frac{b}{2s})^4 - &c.$$

And consequently, if the latter of these two series be taken from the former, the result, by making the first term of the remainder a multiplier, will give, $x = \frac{2bs_1^2}{6s_2} \left\{ 1 + \frac{2.5}{6.9} \left(\frac{b}{2s_2} \right)^2 + , &c. \right\}.$

where, since $s = \sqrt{(\frac{3}{4}b^2 + \frac{1}{27}a^3)}$, we shall have $(\frac{b}{2s})^2 = \frac{27b^2}{27b^2 + 4a^3}$, we shall have $(\frac{b}{2s})^2 = \frac{27b^2}{27b^2 + 4a^3}$, &c. And $\frac{2bx^{\frac{1}{3}}}{6s} = \frac{2b}{6s_3^2} = \frac{2b}{\sqrt{(2(27b^2 + 4a^3))}}$. Whence, also, by substitution, we have the above formula.

* The root, as found by the common formula, when properly reduced, is $x = \pm \sqrt[3]{\frac{b}{2}} \left\{ \sqrt[3]{\left(1 + \frac{2}{b} \checkmark \left(\frac{1}{4}b^2 - \frac{1}{27}a^3\right)\right)} + \sqrt[3]{\left(1 - \frac{2}{b} \checkmark \left(\frac{1}{4}b^2 - \frac{1}{27}a^3\right)\right)} \right\}$

$$x = \pm 2\sqrt[3]{\frac{b}{2}} \left\{ 1 - \frac{2}{3.6} \left(\frac{27b^2 - 4a^3}{27b^2} \right) A - \frac{5.8}{9.12} \left(\frac{27b^2 - 4a}{27b^2} \right) B - \frac{11.14}{15.18} \left(\frac{27b^2 - 4a^3}{27b^2} \right) C - \frac{17.20}{21.24} \left(\frac{27b^2 - 4a^3}{27b^2} \right) D - &c.$$

In which case the upper sign must be taken when b is positive, and the under sign when it is negative; and the same for the first root in the two following cases.

3.
$$x^3 - ax = \pm b$$
,

where $\frac{1}{4}b^2$ is supposed to be less than $\frac{1}{27}a^3$, or $27b^2 \angle 4a^3$.

$$x = \pm 2\sqrt[3]{\frac{b}{2}} \left\{ 1 + \frac{2}{3.6} \left(\frac{4a^3 - 27b^2}{27b^2} \right) - \frac{2.5.8}{3.6.9.12} \left(\frac{4a^3 - 27b^2}{27b^2} \right)^2 + \frac{2.5.8.11.14}{3.6.9.12.15.18} \left(\frac{4a^3 - 27b^2}{27b^2} \right)^3 - &c. \right\}^* \cdot \text{Or},$$

 a^{3}) \}. Or, putting, as in the last case $\frac{2}{b}\sqrt{\left(\frac{1}{4}b^{2}-\frac{1}{27}a^{3}\right)}$, or its equal $\left(\frac{27b2-4a^{3}}{97b2}\right)^{\frac{1}{2}}=s$, we shall have $x=\pm\sqrt[3]{\frac{b}{2}}\left\{\sqrt[3]{(1+s)}+\sqrt[3]{(1-s)}\right\}$.

Whence, extracting the roots of the right hand member of this equation, there will arise $3\sqrt{(1+s)} = 1 + \frac{1}{3}s - \frac{2}{3.6}s^2 + \frac{2.5}{3.6.9}s^3 - \frac{2.5.8}{3.6.9.12}s^4 + &c.$

$$\sqrt[3]{(1-s)} = 1 - \frac{1}{3}s - \frac{2}{3.6}s^2 - \frac{2.5}{3.6.9}s^3 - \frac{2.5.8}{3.6.9.12}s^4 - &c.$$

And, consequently, by adding the two series together, and taking the first term of the result as a multiplier, we shall have $x=\pm2\sqrt[3]{\frac{b}{2}}\left\{1-\frac{2}{3.6}s^2\right\}$

$$\frac{2.5.8}{3.6.9.12}s^4 - \frac{2.5.8.11.14}{3.6.9.12.15.18}s^6 - &c.$$
 Or, by substituting $\left(\frac{27bz - 4a3}{27bz}\right)$

for its equal s, we get the above expression.

* This expression is obtained from the last series, by barely changing the signs of the numerator and denominator in each of its terms; which does not alter their value.

after their value. Hence, in order to determine the other two roots of the equation, let that above found, or its equivalent expression $\sqrt[3]{\frac{1}{2}b + \sqrt{\frac{1}{4}b^2 - \frac{1}{27}a^3}}$

$$+\frac{3}{4}\left\{\frac{1}{2}b-\sqrt{\left(\frac{1}{4}b^2-\frac{1}{27}a^3\right)}\right\}=\pm r.$$

Then, according to the formula that has been before given for these roots,

$$\begin{split} x &= \pm 2 \sqrt[3]{\frac{b}{2}} \left\{ 1 + \frac{2}{3.6} \left(\frac{4a^3 - 27b^2}{27b^2} \right) \mathbf{A} - \frac{5.8}{9.12} \left(\frac{4a^3 - 27b^2}{27b^2} \right) \mathbf{E} \right. \\ &+ \frac{11.14}{15.18} \left(\frac{4a^3 - 27b^2}{27b^2} \right) \mathbf{c} - \frac{17.20}{21.24} \left(\frac{4a^3 - 27b^2}{27b^2} \right) \mathbf{D} + \&c. \end{split}$$

which series answers to the irreducible case, and must be used when 2a3 is less than 27b2.

And if the root thus found be put =r, the other two

roots may be expressed as follows:

roots may be expressed as follows:
$$x = \frac{r}{2} \pm \frac{\sqrt{(4a^3 - 27b^2)}}{93\sqrt{2b^2}} \left\{ 1 - \frac{2.5}{6.9} \left(\frac{4a^3 - 27b^2}{27b^2} \right) + \frac{2.5}{6.9.12.15} \left(\frac{4a^3 - 27b^2}{27b^2} \right)^2 - \frac{2.5.8.11.14.17}{3.6.9.12.15.18.21} \left(\frac{4a^3 - 27b^2}{27b^2} \right)^3 + &c. \right\}.$$

in the former part of the present article, we shall have $x=\mp\frac{r}{2}\pm\frac{\sqrt{r-3}}{2}$ $\left\{\frac{3}{4}\left(\frac{1}{4}b+\sqrt{\left(\frac{1}{4}b^2-\frac{1}{27}a^3\right)}\right)-\frac{3}{4}\left(\frac{1}{2}b-\sqrt{\left(\frac{1}{4}b-\frac{1}{27}a^3\right)}\right)\right\}$. Or, putting $\frac{2}{\hbar}\sqrt{\left(\frac{1}{4}b^2-\frac{1}{27}a^3\right)}=s$, and reducing the expression, $x=\mp\frac{r}{2}\pm$ $\frac{(\frac{1}{b}b)\frac{1}{3}\sqrt{-3}}{2}$ $\left\{\frac{3}{\sqrt{(1+s)}}-\frac{3}{\sqrt{(1-s)}}\right\}$. Whence, extracting the cube roots of the right hand member of this equation, there will arise $\sqrt[3]{(1+s)} = 1 + \frac{1}{3}s - \frac{2}{3.6}s^2 + \frac{2.5}{3.6.9}s^3 - \frac{2.5.8}{3.6.9.12}s^4 + &c.$

 $\frac{3}{4}/(1-s) = 1 - \frac{1}{3}s - \frac{2}{36}s^3 - \frac{2.5}{3.6.9}s^3 - \frac{2.5.8}{3.6.9.12}s^4 - \&c.$

And, consequently, by taking the latter of these series from the former, and making the first term of the remainder a multiplier, we shall have $x = \mp$ $\frac{r}{2} \pm \frac{s(\frac{1}{2}b)\frac{1}{3}\sqrt{-3}}{3} \left\{ 1 + \frac{2.5}{6.9}s^2 + \frac{2.5.8.11}{6.9.12.15}s^4 + \frac{2.5.8.11.14.17}{6.9.12.15.18.21}s^6 + \&c. \right\}$ But since $s = \frac{2}{h} \sqrt{(\frac{1}{4}b^2 - \frac{1}{27}a^3)} = (\frac{27b^2 - 4a^3}{27b^2})\frac{1}{2}, s^2 = \frac{27b^2 - 4a^3}{27b^2} = -\frac{1}{27b^2}$ $\frac{4a_3-27b_2}{27b_2}$, $s_4 = \left(\frac{4a_3-27b_2}{27b_2}\right)^2$, &c., and also $\sqrt[3]{\frac{1}{2}}b \times \frac{s}{3}\sqrt{-3} = \sqrt[3]{\frac{1}{2}}$ $b \times \frac{\sqrt{-3}}{2} \sqrt{\left(\frac{27b^2 - 4a^3}{27b^2}\right)} = \frac{1}{3} \sqrt[3]{\frac{1}{2}} b \times \sqrt{\left(\frac{4a^3 - 27b^2}{9b^2}\right)} =$

 $\frac{\sqrt{(4a^3-27b^2)}}{9\sqrt[3]{2b^2}}$, if these values be substituted for their equals, in the last series, the result will give the above expressions, for the two remaining roots of

the equation.

$$\begin{array}{l} x = \mp \frac{r}{2} \pm \frac{\sqrt{(4a^3 - 27b^2)}}{9\sqrt[3]{2b^3}} \left\{ 1 - \frac{2 \cdot 5}{6 \cdot 9} \left(\frac{4a^3 - 27b}{27b^2} \right) \mathbf{A} + \frac{8 \cdot 11}{12 \cdot 15} \right. \\ \left. \left(\frac{4a^3 - 27b^2}{27b^2} \right) \mathbf{B} - \frac{14 \cdot 17}{18 \cdot 21} \left(\frac{4a^3 - 27b^2}{27b^2} \right) \mathbf{c} + \frac{20 \cdot 23}{24 \cdot 27} \left(\frac{4a^3 - 27b^2}{27b^2} \right) \mathbf{p} - &c. \end{array} \right\}$$

Where $-\frac{1}{2}r$, or $+\frac{1}{2}r$, most be taken according as b is positive or negative; and the double signs ± must be considered as + in one case, and - in the other, as usual. 4. $x^3 - ax = \pm b$,

where $\frac{1}{4}b^2$, is still supposed to be less than $\frac{1}{27}a^3$, or $27b^3$ 1 4a3.

$$x = \pm 2 \frac{2b}{\sqrt[3]{(2(4a^3 - 27b^3))}} \left\{ 1 - \frac{2.5}{6.9} \left(\frac{27b^2}{4a^3 - 27b^2} \right) + \frac{2.5.8.11}{6.9.12.15} \left(\frac{27b^2}{4a^3 - 27b^2} \right)^2 - \frac{2.5.8.11.14.17}{6.9.12.15.18.21} \left(\frac{27b^2}{4a^3 - 27b^2} \right)^3 + &c. \right\}. * Or,$$

* By transposing the terms of the common formula, as in the first case, we shall have $x = \sqrt[3]{\left\{\sqrt{\left(\frac{1}{4}b^2 - \frac{1}{27}a^3\right) + \frac{1}{2}b\right\}} - \sqrt[3]{\left(\frac{1}{4}b^2 - \frac{1}{27}a^3\right)} - \frac{1}{2}a^3}$ $\frac{1}{2}b$. Or, by putting, for the sake of simplicity, as before, $\sqrt{\frac{1}{4}b^2-\frac{1}{2\sqrt{3}}}$ $(a^3) = s$, and reducing the equation $x = \frac{3}{s} / s \left\{ \frac{3}{s} / (1 + \frac{b}{a_s}) - \frac{3}{s} / (1 - \frac{b}{a_s}) \right\}$

Whence, extracting the roots of the right hand member, as in the former instances, $(1 + \frac{b}{2s}) = 1 + \frac{1}{3}(\frac{b}{2s}) - \frac{1}{3.6}(\frac{b}{2s})^2 + \frac{2.5}{3.6.9}(\frac{b}{2s})^3 - \frac{2.5.8}{3.6.9.12}(\frac{b}{2s})^4$ +&c. $\frac{3}{2}\left(1-\frac{b}{2a}\right)=1-\frac{1}{3}\left(\frac{b}{2a}\right)-\frac{2}{66}\left(\frac{b}{2a}\right)^2-\frac{2.5}{369}\left(\frac{b}{2a}\right)^3-\frac{2.5.8}{3.6.9.12}$

$$(\frac{b}{2s})^4 - \&c.$$

And, consequently, by taking the latter of these series from the former,

And, consequently, by taking the latest of these states and making the first term of the result a multiplier, we shall have
$$x = \frac{2bs_3^{\perp}}{6s} \left\{ 1 + \frac{2.5}{6.9} (\frac{b}{2s})^2 + \frac{2.5.8.11}{6.9.12.15} (\frac{b}{2s})^4 + \frac{2.5.8.11.14.17}{6.9.12.15.18.21} (\frac{b}{2s})^6 + \frac{2.5.8.11.14.17}{6.9.12.15.18.21} (\frac{b}{2s})^6 + \frac{2.5.8.11}{6.9.12.15} (\frac{b}{2s})^4 + \frac{2.5.8.11.14.17}{6.9.12.15.18.21} (\frac{b}{2s})^6 + \frac{2.5.8.11}{6.9.12.15} (\frac{b}{2s})^6 + \frac{2.5.8.11}{6.9.12} (\frac{b}{2s})^8 + \frac{2.5.8.11}{6.9.12$$

$$\begin{array}{l} x = \mp \frac{2b}{\sqrt[3]{(2(4a^3 - 27b^2))}} \left\{ 1 - \frac{2.5}{6.9} \left(\frac{27b^2}{4a^3 - 27b^2} \right) \mathbf{A} + \frac{8.11}{12.15} \right. \\ \left. \left(\frac{27b^2}{4a^3 - 27b^2} \right) \mathbf{B} - \frac{14.17}{16.21} \left(\frac{27b^2}{4a^3 - 27b^2} \right) \mathbf{c} + \frac{20.23}{24.27} \left(\frac{27b^2}{4a^2 - 27b^2} \right) \\ \mathbf{B} - \&c. , \text{ which series also answers to the irreducible case,} \end{array}$$

 $=-\frac{27b^2}{4a^3-27b^2}$, $(\frac{b}{2s})^4=(\frac{27b^2}{4a^3-27b^2})^2$ &c., and $\frac{2bs\frac{1}{3}}{6s}=\frac{2b}{6s^2}=$

 $6\sqrt[3]{(\frac{1}{4}b^2-\frac{1}{2}7^a)}$ $\sqrt{2(4a^3-27b^2)}$ Whence, if these values be substituted for their equals in the last series, there will arise the above expression for the first root of the equation. And, if we put the root thus found, or its equivalent expression

 $\sqrt[3]{\left(\frac{1}{4}b^2 - \frac{1}{27}a^3\right) + \frac{1}{2}b\left(-\frac{3}{2}\right)} \left(\sqrt[4]{b^2 - \frac{1}{27}a^3} - \frac{1}{2}b\right) = \pm r,$ we shall have, according to the formula before given for the other two roots, $x = \mp \frac{r}{2} \pm \frac{\sqrt{-3}}{2} \left\{ \sqrt[3]{(\sqrt{(\frac{1}{4}b^2 - \frac{1}{2}7}a^3)} + \frac{1}{2}b) + \sqrt[3]{(\frac{1}{4}b^2 - \frac{1}{2}7}a^3)} \right\}$ $-\frac{1}{2}b$. Or, taking, as before, $\sqrt{(\frac{1}{4}b^2-\frac{1}{27}a^3)}=s$, and simplifying the result, $x = \mp \frac{r}{2} \pm \frac{s \pm \sqrt{-3}}{2} \left\{ \sqrt[3]{(1 + \frac{b}{2s})} + \sqrt[3]{(1 - \frac{b}{2s})} \right\}$

Whence, by extracting the roots of the right hand side of this equation,

$$\begin{array}{l} \sqrt[3]{(1+\frac{b}{2s})} = 1 + \frac{1}{3}(\frac{b}{2s}) - \frac{2}{3.6}(\frac{b}{2s})^2 + \frac{2.5}{3.6.9}(\frac{b}{2s})^3 - \frac{2.5.8}{3.6.9.12}(\frac{b}{2s})^4 + \text{\&c.} \\ + \text{\&c.} \quad \sqrt[3]{(1-\frac{b}{2s})} = 1 - \frac{1}{3}(\frac{b}{2s}) - \frac{2}{3.6}(\frac{b}{2s})^2 - \frac{2.5}{3.6.9}(\frac{b}{2s})^3 - \frac{2.5.8}{3.6.9.12}(\frac{b}{2s})^4 - \text{\&c.} \end{array}$$

And, consequently, if the latter of these series be added to the former, we shall have, by making the first term of the result a multiplier, $x = \mp \frac{r}{2} \pm 3\frac{1}{3} \sqrt{-3} \begin{cases} 1 - \frac{2}{3.6} (\frac{b}{2s})^2 - \frac{2.5.8}{3.6.9.12} (\frac{b}{2s})^4 - \frac{2.5.8.11.14}{3.6.9.12.15.13} \end{cases}$ $\left(\frac{b}{2s}\right)^6$ - &c. But since $s = \sqrt{\left(\frac{1}{4}b^2 - \frac{1}{27}a^3\right)} = \left(\frac{27b^2 - 4a^3}{497}\right)^{\frac{1}{2}}$, we shalf also have $(\frac{b}{2s})^2 = \frac{27b^2}{27b^2 - 4a^3} = -\frac{27b^2}{4a^3 - 27b^2}$ &c., and consequently, s $\frac{1}{3}\sqrt{-3} = \sqrt[6]{\left(\frac{4a_3 - 27b^2}{b}\right)}$

Hence, if these values be substituted for their equals in the above series, the result will give the above expressions for the two remaining roots of the equation.

and must be used when $2a^{2}$ is greater than $27b^{2}$. And if the root thus found, be put =r, as before, the other two roots may be expressed thus: $x=\mp\frac{r}{2}\pm\sqrt{\frac{4a^{3}-27b^{2}}{4}}\left\{\frac{1+\frac{2}{3.6}(\frac{27b^{3}}{4a^{3}-27b^{2}})-\frac{2.5.8}{3.6.9.12}(\frac{27b^{2}}{4a^{3}-27b^{2}})^{2}+\frac{2.5.8.11.14}{3.6.9.12.15.18}(\frac{27b^{2}}{4a^{3}-27b^{2}})^{3}-\&c.\right\}.$ Or, $x=\pm\frac{r}{2}\pm\sqrt[3]{\frac{4a^{3}-27b^{2}}{4}}\left\{1+\frac{2}{3.6}(\frac{27b^{2}}{4a^{2}-27b^{2}})\Lambda-\frac{5.8}{9.12}(\frac{27b^{2}}{4a^{3}-27b^{2}})\mathrm{E}+\frac{11.14}{15.18}(\frac{27b^{2}}{4a^{3}-27b^{2}})\mathrm{C}-\frac{17.20}{21.24}(\frac{27b^{2}}{4a^{3}-27b^{2}})\mathrm{E}+\&c.\right\}$

Where the signs are to be taken as in the latter part of

the preceding case.

EXAMPLES.

of x.
; and
nsequen
000 (A) 61 (B)
77 (c)
82 (D)
62 (E)

tly,

CODIC EQ	
$\frac{26.29}{30.33} \times \frac{1}{9} E$.0000005 (F)
	1.0217787
Log. 1.0217787 Log. 3/648 Colog. 27	0.0093570 0.9371916 8.5686362
No. 3274801	-1.5151848
2. Given $x^3 - 9x = 12$ to find Here $a = 9$ ar	ad b=12;
nence $2\sqrt[3]{\frac{12}{2}} = 2\sqrt[3]{6}$ and $\frac{27b^2}{2}$	$\frac{7b^2}{27 \times 144}$
$\frac{44 - 108}{144} = \frac{36}{144} = \frac{1}{4}.$	
Consequently by formula 2 w	ve shall have 1.0000000 (A)
$-\frac{2}{3.6}\times\frac{1}{4}(A)$	—.0277778 (в)
$-\frac{5.8}{9.12} \times \frac{1}{4}$ (B)	0025720 (c)
$-\frac{11.14}{15.18} \times \frac{1}{4}$ (c)	-:0003667 (D)
$-\frac{17.20}{21.24} \times \frac{1}{4}$ (D)	0000619 (E)
$-\frac{23.26}{27.30} \times (E)$	0000114 (F)
$-\frac{29.32}{33.36} \times \frac{1}{4} (F)$	0000022 (G)
Sum	0307920
Comp.	.9692080

oble Ego	ATIONS.
Log. 969208 Log. 23/6 or Log. 3/48	-1.9864137 0.5604137
No. 3.522334	.5468274
therefore $x=$ 3. Given $x^3-12x=15$ to fin Here $a=12$ an	d the three values of $b=15$;
whence $2\sqrt[3]{\frac{b}{2}} = 2\sqrt[3]{\frac{15}{2}} = \sqrt[3]{\frac{15}{2}} = \sqrt[3]{\frac{15}{2}$	
$\frac{12 - 27.15^2}{27.15^2} = \frac{256 - 225}{225} = \frac{3}{225}$	1 .
27.152 225 29 Consequently by formula 3,	
1	+1.0000000 (A
$+\frac{2}{3.6}\times\frac{31}{225}$ A	+0.0153086 (в
$-\frac{5.8}{9.12} \times \frac{31}{225}$ B	-0.0007812 (c)
$+\frac{11.14}{15.18} \times \frac{31}{225} c$	+0.0000614 (D)
$-\frac{17.20}{21.24} \times \frac{31}{225}$ D	-0.0000057 (E
$+\frac{23.26}{27.30} \times \frac{31}{225} E$	+0.0000006 (F)
Sum of + Terms Sum of - Terms	+1.0153706 0007869
Difference	1.0145837
Log. 1.0145837 Log. 3/60	.0062880
No. 3.971362	.59900.11

Therefore the affirmative value of x or first root, r=3.971962.

Whence the three roots or values of x are 3.971962,—1.5760365 and -2.395925.

4. Given $x^3 - 6x = 2$ to find the three values of x.

Here
$$\frac{-2b}{\sqrt[3]{(2(4a^3-27b^2))}} = \frac{-4}{\sqrt[3]{(2(4.6^3-27.4))}} =$$

Colog. 21

No. 339870

Therefore one of the negative roots or values of x, is

8.6777807

1.5313214

-.339870=-r.
Again
$$\sqrt[6]{\frac{4a^3-27b^2}{4}} = \sqrt[6]{\frac{4.6^3-27.4}{4}} = \sqrt[6]{(6^3-27)} = \sqrt[6]{189}$$
 and $\frac{27b^2}{4a^3-27b^2} = \frac{1}{7}$.
Hence.

1 1.0000000 (A)
$$+\frac{2}{3.6} \times \frac{1}{7}$$
 A +0.0158730 (B)

$-\frac{5.8}{9.12} \times \frac{1}{7} B$.0008398 (c)
$+\frac{11.14}{15.18}\times\frac{1}{7}c$	+.0000684 (д)
$-\frac{17.20}{21.24} \times \frac{1}{7}$ D	0000066 (E
$+\frac{23.26}{27.30}\times\frac{1}{7}$ E	+.0000002 (r
K.	Sum 1.0150952
Log. 1.0150952	.0065070
Log. 189 No. 2.431741	.3794103
	.3609173
Therefore $\frac{r}{2}$	+.169935
Last number	±2.431741
Result	+2.601676

Or — 2.261806 And consequently +2.601676, -2.261806, and -.339876 are the three roots required.

EXAMPLES FOR PRACTICE.

1. Given $x^3+9x=30$, to find the root of the equation, or the value of x.

Ans. x=2.180849.

2. Given $x^3-2x=5$, to find the root of the equation, or the value of x.

Ans. x=2.0945515.

3. Given $x^3-3x=3$, to find the root of the equation, or the value of x. Ans. 2.103803.

4. Given $x^3 - 27 x = 36$, to find the three roots or values of x. Ans. 5.765722, --4.320634, and --1.445038, 5. Given $x^3 - 48x^2 = -200$, to find the root of the

equation, or the value of x. Ans. 47.9128.

6. Given $x^3 - 22x = 24$, to find the root of the equation, or the value of x.

Ans. 5.162277.

OF BIQUADRATIC EQUATIONS.

A biquadratic equation, as before observed, is one that rises to the fourth power, or which is of the general form $x^4 + ax^3 + bx^2 + cx + d = 0$.

The root of which may be determined by means of the following formula; substituting the numbers of the given equation, with their proper signs, in the places of the literal coefficients a, b, c, d.

BULE LA

Find the value of z in the cubic equation $z^3 + (\frac{1}{4}ac \frac{1}{18}b^2-d$) $z=\frac{1}{108}b^3+\frac{1}{8}(c^2+da^2)-\frac{b}{24}(ac+8d)$ by one of the former rules; and let the root, thus determined, be denoted by r. Then find the two values of x in each of the following quadratic equations.

^{*} This method is that given by Simpson, p. 150 of his Algebra, which consists in supposing the given biquadratic to be formed by taking the difference of two complete squares, being the same in principle as that of Fer-

Thus, let the proposed equation be of the form x4 + ax3 + bx2 + cx + d==0 (1), having all its terms complete; and assume $(x^2 + \frac{1}{2}ax + p)^2 - (qx)$ $+r)^2 = x^4 + ax^3 + bx^2 + cx + d.$

Then, if $x^2 + \frac{1}{2}ax + p$ and qx + r be actually involved, we shall have Then, if $x^3 + 2a^2 + r$ and $x^2 + ap$ $\begin{vmatrix} x + p^2 \\ -r^2 \end{vmatrix} = x^4 + ax^3 + bx^2 + cx + d$.

And, consequently, by equating the homologous terms, there will arise

where, since the product of the first and last of the absolute terms of these equations is evidently equal to a of the square of the second, we shall have

 $²p3 + (\frac{1}{4}a^2 - b) p^2 - 2dp - d(\frac{1}{4}a^2 - b) = \frac{1}{4}(a^2p^2 - 2acp + c^2).$ Or, by bringing the unknown quantities to the left hand side, and the known to the right, and then dividing by 2.

$$x^{2} + \left(\frac{1}{2}a + \sqrt{\frac{1}{4}a^{2} + 2(r - \frac{1}{3}b)}\right)x = -\left(r + \frac{1}{6}b\right) + \sqrt{\left(r + \frac{1}{6}b\right)^{2} - d}$$

$$x^{2} + \left(\frac{1}{2}a - \sqrt{\frac{1}{4}a^{2} + 2(r - \frac{1}{3}b)}\right)x = -\left(r + \frac{1}{6}b\right) - \sqrt{\left(r + \frac{1}{6}b\right)^{2} - d}$$

and they will be the four roots of the biquadratic required,

EXAMPLES.

1. Given the equation $x^4 - 10x^3 + 35x^2 - 50x + 24 = 3$, to find its roots.

Here a = -10, b = 35, c = -50, and d = 24;

Whence, by substituting these numbers in the cubic equation

$$p_3 - \frac{b}{a}p_2 + \frac{1}{2}(ac - 4d)p = \frac{1}{8}(c_2 + a_2d) - \frac{1}{2}bd(2)$$

From which last equation p can be determined by the rules before given for cubics.

And since, from the preceding equations, it appears that

$$q = \sqrt{(2p + \frac{1}{4}a^2 - b)}$$
 and $r = \frac{ap - c}{2a}$, or $\sqrt{(p^2 - d)}$,

it is evident that the several values of x can be obtained from the quantities thus found.

For, because $x4 + ax^3 + bx^2 + cx + d$, or its equal $(x2 + \frac{1}{2}ax + p)2 - (x + p)2 = 0$, it is plain that $(x2 + \frac{1}{2}ax + p)2 = (x + p)2$. And, therefore, by extracting the roots of each side of this equation, there will arise

 $x^2 + \frac{1}{2}ax + p = qx + r$; or $x^2 + (\frac{1}{2}a - q)x = r - p$. Whence, by substituting the above values of p, q, and r, for their equals, and transposing the terms, we shall have $a^2 + \frac{1}{3}a + \sqrt{(2p + \frac{1}{4}a^2 - b)}$ $x+p \mp \sqrt{(p^2-d)}=0$, for the case where ap-c is positive; and

 $x^2 + \frac{1}{2}a + \sqrt{(2p + \frac{1}{4}a^2 - b)} \left\{ x + p + \sqrt{(p^2 - d)} = 0, \right\}$

for the case where ep-e is negative; which two quadratics give the four roots of the proposed equation.

And by putting $p=z+\frac{b}{\epsilon}$, in the reducing equation (2), in order to destroy its second term, the several steps of the investigation may be made to agree with the expressions given in the above rule.

$$z^{3} + \left(\frac{1}{4}ac - \frac{1}{12}b^{2} - d\right)z = \frac{1}{108}b^{3} + \frac{1}{8}(c^{2} + du^{2}) - \frac{b}{24}(ac + 8d),$$

we shall have the following reduced equation,

$$z^3 - \frac{13}{12}z = \frac{35}{108}$$

which being resolved, according to the rule before laid down for that purpose, gives

$$z = \frac{1}{6} \left\{ \sqrt[3]{35 + 18} \sqrt{-3} + \sqrt[3]{35 - 18} \sqrt{-3} \right\}.$$

But, by the rule for binomial surds, given in the former part of the work,

$$\sqrt[3]{(35+18\sqrt{-3})} = \frac{7}{2} + \frac{1}{2}\sqrt{-3}$$
, and $\sqrt[3]{(35-18\sqrt{-3})} = 7 - \frac{1}{2}\sqrt{-3}$;

Wherefore $z = \frac{1}{6} \left\{ \frac{7}{2} + \frac{1}{2} \sqrt{-3} + \frac{7}{2} - \frac{1}{2} \sqrt{-3} \right\} = \frac{7}{6}$.

And if this number be substituted for $r_1 = 10$ for $a_1 = 35$ for b_1 , and 24 for d_2 , in the two quadratic equations,

$$x^{2} + \left(\frac{1}{2}a + \sqrt{\frac{1}{4}a^{2} + 2(r - \frac{1}{3}b)}\right)x = -(r + \frac{1}{6}b) + \sqrt{\frac{1}{6}b}$$

$$+ \frac{1}{6}b)^{2} - d,$$

$$x^{2} + \left(\frac{1}{2}a - \sqrt{\frac{1}{4}a^{2} + 2(r - \frac{1}{3}b)}\right)x = -(r + \frac{1}{6}b) - \sqrt{\frac{1}{6}a^{2} + \frac{1}{6}b^{2}}$$

$$+ \frac{1}{6}b^{2} - d,$$

they will become, after reducing them to their most simple terms,

 $x^2 - 3x = -2$, and $x^2 - 7x = -12$: from the first of which $x = \frac{3}{2} \pm \sqrt{\frac{1}{4}} = \frac{3}{2} \pm \frac{1}{2} = 2$ or 1, and

from the second $x = \frac{7}{2} \pm \sqrt{\frac{1}{4}} = \frac{7}{2} \pm \frac{1}{2} = 4$ or 3;

Whence the four roots of the given equation are 1, 2, 3, and 4.

Or, when its second term is taken away, it will be ef

the form $x^4+bx^2+cx+d=0$, to which it can'always be reduced; and in that case, is solution may be obtained by the following rule:

RULE II.

Find the value of z in the cubic equation

$$z^{2} - \left(\frac{1}{12}b^{2} + d\right)z = \frac{1}{108}b^{3} + \frac{1}{8}c^{2} - \frac{1}{3}bd,$$

and let the root thus determined be denoted by r.

Then find the two values of x, in each of the following quadratic equations.

$$x^{3} + (\sqrt{2(r - \frac{1}{5}b)})x = -(r + \frac{1}{6}b) + \sqrt{(r + \frac{1}{6}b)^{2} - d}$$

$$x^{2} + (\sqrt{2(r - \frac{1}{3}b)})x = -(r + \frac{1}{6}b) - \sqrt{(r + \frac{1}{6}b)^{2} - d}$$

and they will be the four roots of the biquadratic equation required*.

 The method of solving biquadratic equations was first discovered by Louis Fermin, a disciple of the colebrated Chritim, before mentioned: but the above rule is herived from that grown by Descartes in his Gamatery, published in 1637, the ruth of which may be shown as follows:
 Let the thereby proposed equation in

as the first property and the second and c = 0.

and conceive it to be produced by the multiplication of the two quadratics $xx^2 + yx + q = 0$, and $xx^2 + rx + s = 0$.

Then, since these equations, as well as the given one, are each =0, there will arise, by taking their product,

x = +(y+r)x 3 + (x + y + pr)x 2 + (pr + pr)x + px = x 4 + xx 2 + λx 2 + x 2 + λx 2 + x 2 + λx 3 + x 2 + λx 3 + x

Whence, subtracting the square of the third of these from that of the second, and then changing the sides of the equation, we shall have

Where the value of p may be found by the rule before given for cubic equations.

Hence, also, since $s+\varsigma=s+pz$, and $s-q=\frac{s}{p}$, there will arise, by addition and advanction,

Or the four roots of the given equation, in this last case, will be as follows:

$$\begin{split} x &= -\frac{1}{2} \sqrt{\left\{ 2(r - \frac{1}{3}b) \right\}} + \sqrt{\left\{ -\frac{r}{2} + \frac{b}{3} + \sqrt{[(r + \frac{1}{6}b)^2 - d]} \right\}} \\ x &= -\frac{1}{2} \sqrt{\left\{ 2(r - \frac{1}{3}b) \right\}} - \sqrt{\left\{ -\frac{r}{2} - \frac{b}{3} + \sqrt{[(r + \frac{1}{6}b)^2 - d]} \right\}} \\ x &= -\frac{1}{2} \sqrt{\left\{ 2(r - \frac{1}{3}b) \right\}} + \sqrt{\left\{ -\frac{r}{2} - \frac{b}{3} - \sqrt{[(r + \frac{1}{6}b)^2 - d]} \right\}} \\ x &= -\frac{1}{2} \sqrt{\left\{ 2(r - \frac{1}{3}b) \right\}} - \sqrt{\left\{ -\frac{r}{2} - \frac{b}{3} - \sqrt{[(r + \frac{1}{6}b)^2 - d]} \right\}} \end{split}$$

2. Given $x^4 + 12x - 17 = 0$, to find the four roots of the equation.

Here a=0, b=0, c=12, and d=-17;

Whence, by substituting these numbers in the cubic equation

 $z^3 - (\frac{1}{12}b^2 + d)z = \frac{1}{108}b^3 + \frac{1}{8}c^2 - \frac{1}{3}bd,$

we shall have, after simplyfying the results, $z^3+17z=18$,

Where it is evident, by inspection, that z=1,

And if this number be substituted for r, 0 for b, and - 17 for d in the two quadratic equations in the above rule, their solution will give

$$\begin{array}{l} x = -\frac{1}{2}\sqrt{2} + \sqrt{(-\frac{1}{2} + \sqrt{18})} = -\frac{1}{2}\sqrt{2} + \sqrt{(-\frac{1}{2} + 3\sqrt{2})} \\ x = -\frac{1}{2}\sqrt{2} - \sqrt{(-\frac{1}{2} + \sqrt{18})} = -\frac{1}{2}\sqrt{2} - \sqrt{(-\frac{1}{2} + 3\sqrt{2})} \\ x = +\frac{1}{2}\sqrt{2} + \sqrt{(-\frac{1}{2} - \sqrt{18})} = +\frac{1}{2}\sqrt{2} + \sqrt{(-\frac{1}{2} - 3\sqrt{2})} \\ x = +\frac{1}{2}\sqrt{2} - \sqrt{(-\frac{1}{2} - \sqrt{18})} = +\frac{1}{2}\sqrt{2} - \sqrt{(-\frac{1}{2} - 3\sqrt{2})} \end{array}$$

 $x=+\frac{1}{2}\sqrt{2}-\sqrt{(-\frac{1}{2}-\sqrt{18})}=+\frac{1}{2}\sqrt{2}-\sqrt{(-\frac{1}{2}-3\sqrt{2})}$ Which are the four roots of the proposed equation; the two first being real, and the two last imaginary.

$$s = \frac{1}{2}a + \frac{1}{2}p^2 + \frac{b}{2a}; \ \eta = \frac{1}{2}a + \frac{1}{2}p^2 - \frac{b}{2a};$$

where p being known, s and q are likewise known. And, consequently, by extracting the roots of the two assumed quadratics $x^2 + px + q = 0$, and $x^2 + rx + s = 0$, or its equal x - px + s = 0, we shall

 $x=-3p+\sqrt{(3p^2-q)}$; $x=\frac{1}{2}p\pm\sqrt{(3p^2-s)}$; which expressions, when taken in + and -, give the four roots of the proposed biquadratic, as was required.

RULE* III.

The roots of any biquadratic equation of the forms $x^4 +$ $ax^2+bx+c=0$, may also be determined by the following general formulæ first given by EULER; which are remarkable for their elegance and simplicity.

* This method, which differs considerably from either of the former, consists in supposing the root of the given equation,

 $x^4 + ax^2 + bx + c = 0(1),$

to be of the following trinomial surd form

 $x = \sqrt{p} + \sqrt{q} + \sqrt{r};$ where p, q, r, denote the roots of the cubic equation,

 $y^3 + fy^2 + gy = h(2),$ of which the coefficients f, g, and the absolute term h, are the unknown

quantities that are to be determined. Then, agreeably to the theory of equations before given, we shall have p

+q+r=-f; pq+pr+qr=g; pqr=h. And, by squaring each side of the formula expressing the value of x, $x^2 = p + q + r + 2\sqrt{pq} + 2\sqrt{pr} + 2\sqrt{qr}$.

Or, by substituting for its equal -(p+q+r), and bringing the term, so obtained, to the other side of the equation

 $x^2 + f = 2\sqrt{pq} + 2\sqrt{pr} + 2\sqrt{qr}$. Also, by again squaring each side of this last expression, we shall have x^2

 $+2fx^2+fy^2=4pq+4pr+4qr+3\sqrt{p^2qr+8\sqrt{q^2pr+8\sqrt{r^2pq}}}$. Or substituting 4g for its equal 4pq+4pr+4qr, and bringing the term to the other side as before,

 $x^4 + 2fx^2 + f^2 - 4g = 3 \sqrt{pqr}(\sqrt{p} + \sqrt{q} + \sqrt{r}).$ But since, from what has been above laid down, we have

 $\sqrt{p} + \sqrt{q} + \sqrt{r} = x$, and $\sqrt{pqr} = \sqrt{h}$, if these be put for their equals in the last equation, it will become, by this substitution,

 $x4 + 2fx2 - 8h^{\frac{1}{2}}x + f^2 - 4g = 0$. Whence, comparing these coefficients with those of the given equation,

there will arise

there will arise $2f=a; -8 \sqrt{h}=b; f^2-cg=c, \text{ or,} \\ f=\frac{a}{2}; h=\frac{b^2}{64}; g=\frac{a^2}{16}-\frac{c}{4}.$ And, consequently, by substituting these values in the assumed cubic equation (2), we shall have

 $y^3 + \frac{1}{2}ay^2 + \frac{1}{16}(a^2 - 4c)y = \frac{b^2}{61}(3)$.

the three roots of which last equation, when substituted for p, q, and r, in the formula $x=\sqrt{p+\lambda}+\sqrt{q}$, will give, by taking each term of the expression both in + and -, all the four values of x. Or, in order to render this result more commodious in practice, by freeing it from tractions, let $y=\pm x$. Then by substitution and reduction, we shall

have the corresponding equation

Find the three roots of the cubic equation $z^3 + 2az^2 + (a^2 - 4c)z = b^2$, by one of the former rules, before given for this purpose; and let them be denoted by r', r', and r'''. Then, we shall have

When b is positive, $x = \frac{-\sqrt{r'} - \sqrt{r''} - \sqrt{r'''}}{2}$ $x = \frac{-\sqrt{r'} + \sqrt{r''} + \sqrt{r''}}{2}$ $x = \frac{+\sqrt{r'} - \sqrt{r''} + \sqrt{r''}}{2}$ $x = \frac{+\sqrt{r'} - \sqrt{r''} + \sqrt{r''}}{2}$ $x = \frac{+\sqrt{r'} + \sqrt{r'} - \sqrt{r''}}{2}$ $x = \frac{-\sqrt{r'} + \sqrt{r'} - \sqrt{r''}}{2}$ $x = \frac{-\sqrt{r'} - \sqrt{r''} + \sqrt{r''}}{2}$ $x = \frac{-\sqrt{r'} - \sqrt{r''} + \sqrt{r''}}{2}$

Note. If the three roots r', r'', r''', of the auxiliary cubic equation be all real and positive, the four roots of the proposed equation will, also, be real; and if one of these roots be positive, and the other two imaginary, or both of them negative, and equal to each other, two of the roots of the given equation will be real, and two imaginary; which are the only cases that produce real results.

3. Given $x^4 - 25x^2 + 60x - 36 = 0$, to find the four roots of the equation.

Here a = -25, b = 60, and c = -36;

Whence, by substituting these values for their equals, in the cubic equation above given, we shall have $z^3 - 2 \times 25$ $z^2 + (25^2 + 4 \times 36) z = 60^2$, or $z^3 - 50z^2 + 769z = 3600$:

 $z^3 + 2az^2 + (a^2 - 4c)z = b^2$, (4)

the three roots of which are each, evidently, four times those of the former. Hence using this instead of equation (3), and denoting its roots by r', r'', r'', r'', be last mentioned formula, taking each of its terms in + and -, as before,

will give the values of α , as in the above expressions.

Note. If we were to take all the possible changes of the signs, in this case, which the terms of the assumed formula admit of, it would appear that x should have eight different values; but it is to be observed, that, according to the first part of the above investigation, the product $\sqrt{p} \times \sqrt{q} \times \sqrt{r} + \sqrt{p}$, or $\frac{1}{8}b$; and, consequently, that when b is positive, either all the three radicals must be taken in +, or two in—and one in +; and when b is acgarize, they must either be all—or two + and one -; which considerations reduce the number of roots to four.

the three roots of which last equation, as found by trial, or by one of the former rules, are 9, 16, and 25, respectively; whence

$$x = \frac{1}{2}(-\sqrt{9} - \sqrt{16} - \sqrt{25}) = \frac{1}{3}(-3 - 4 - 5) = -6$$

$$x = \frac{1}{2}(-\sqrt{9} + \sqrt{16} + \sqrt{25}) = \frac{1}{2}(-3 + 4 + 5) = +3$$

$$x = \frac{1}{2}(+\sqrt{9} - \sqrt{16} + \sqrt{25}) = \frac{1}{2}(+3 - 4 + 5) = +2$$

$$x = \frac{1}{2}(+\sqrt{9} + \sqrt{16} - \sqrt{25}) = \frac{1}{2}(+3 + 4 - 5) = +1$$

And consequently the four roots of the proposed equation are 1, 2, 3, and -6.

EXAMPLES FOR FRACTICE.

- 1. Given $x^4 55x^2 30x + 504 = 0$, to find the four
- roots, or values of x. Ans. 3, 7, -4, and -6. 2. Given $x^4 + 2x^3 7x^2 8x = -12$, to find the four Ans. 1, 2, -3, and -2. roots, or values of x.
- 3. Given $x^4 8x^3 + 14x^2 + 4x = 8$, to find the four roots, Ans. $\begin{cases} 3+\sqrt{5}, 3-\sqrt{5} \\ 1+\sqrt{3}, 1-\sqrt{3} \end{cases}$ or values of x.
- 4. Given $x^4 17x^2 20x 6 = 0$, to find the four roots,
- Ans. $\begin{cases} 2+\sqrt{7}, & 2-\sqrt{7} \\ -2+\sqrt{2}, & -2-\sqrt{2} \end{cases}$ or values of x.
- 6. Given $x^4 3x^2 4x = 3$, to find the four roots, or values of x. Ans. $\begin{cases} -\frac{1}{2} + \frac{1}{2}\sqrt{13}, & \frac{1}{2} \frac{1}{2}\sqrt{13} \\ -\frac{1}{2} + \frac{1}{2}\sqrt{-3}, & -\frac{1}{2} \frac{1}{2}\sqrt{-3} \end{cases}$ 6. Given $x^4 19x^3 + 132x^2 302x + 200 = 0$, to find the
- four roots, or the values of x. Ans. $\begin{cases} 1.02304, 4.00000 \\ 6.57653, 7.39543 \end{cases}$
- 7. Given $x^4 27x^3 + 162x^2 + 356x 1200 = 0$, to find the four roots, or values of x. Ans. $\begin{cases} 2.05608, -3.00000 \\ 13.15306, 14.79086 \end{cases}$
- 8. Given $x^4 12x^2 + 12x 3 = 0$, to find the four roots, or values of x.
- 9. Given $x^4 36x^2 + 72x 36 = 0$, to find the four roots, Ans $\begin{cases} 0.8729836, & 1.2679494 \\ 4.7320506, & -6.8729836 \end{cases}$ or values of x.
- 10. Given $x^4 12x^3 + 47x^2 72x + 36 = 0$, to find the Ans. 1, 2, 3 and 6. roots, or values of x.

11. Given $x^4 + 24x^3 - 114x^3 - 24x + 1 = 0$, to find the roots, or values of x.

Ans. $\begin{cases} +\sqrt{197} - 14, \ 2+\sqrt{5} \\ -\sqrt{197} - 14, \ 2-\sqrt{5} \end{cases}$ 12. Given $x^4 - 6x^3 - 58x^3 - 114x - 11 = 0$, to find the

12. Given $x^2 - 6x^3 - 58x^3 - 114x - 11 = 0$, to find the roots, or the values of x.

Ans. $\pm \frac{5}{2} \sqrt{3} + \frac{3}{2} \pm \sqrt{(17 \pm \frac{2}{2} \sqrt{3})}$.

OF THE

RESOLUTION OF EQUATIONS

BY APPROXIMATION.

Equations of the fifth power, and those of higher dimensions, cannot be resolved by any rule or algebraic formula that has yet been discovered; except in some particular cases, where certain relations subsist between the coefficients of their several terms, or when the roots are rational; and, for that reason, can be easily found by means of a few trials.

In these cases, therefore, recourse must be had to some of the usual methods of approximation; among which that commonly employed is the following, which is universally applicable to all kinds of numeral equations, whatever may be the number of their dimensions, and though mot strictly accurate, will give the value of the root sought to any required degree of exactness.

RULE.

Find, by trials, a number nearly equal to the root sought, which call r; and let z be made to denote the difference between this assumed root, and the true root x.

Then instead of x, in the given equation, substitute its equal $r \pm z$, and there will arise a new equation, involving only z and known quantities.

Reject all the terms of this equation in which z is of two or more dimensions; and the approximate value of z may then be determined by means of a simple equation.

And if the value, thus found, be added to, or subtracted from that of r according as r was assumed too little or too great, it will give a near value of the root required.

But as this approximation will seldom be sufficiently exact, the operation must be repeated, by substituting the number thus found for r, in the abridged equation exhibiting the value of z; when a second correction of z will be obtained, which, being added to, or subtracted from r, will give a nearer value of the root than the former.

And by again substituting this last number for r in the above mentioned equation, and repeating the same process as often as may be thought necessary, a value of x may be

found to any degree of accuracy required.

Note. The decimal part of the root, as found both by this and the next rule, will, in general, about double itself at each operation; and therefore it would be useless as well as troublesome, to use a much greater number of figures than these in the several substitutions for the values of r.*

EXAMPLES.

1. Given $x^3 + x + x = 90$, to find the value of x by approximation.

Here the root, as found by a few trials, is nearly equal to 4.

^{*}It may here be observed, that if any of the roots of an equation be whole numbers, they may be determined by substituting 1, 2, 3, 4, &c. successively, both in plus and in minus, for the unknown quantity, till a result is obtained equal to that in the question; when those that are found to succeed, will be the roots required.

Or, since the last term of any equation is always equal to the continued product of all its roots, the number of these trials may be generally diminished, by finding all the divisors of that term, and then substituting them both in plus and minus, as before, for the unknown quantity, when those that give the proper result will be the rational roots sought; but if none of them are found to succeed, it may be concluded that the equation cannot be resolved by this method; the roots in that case, being either irrational or imaginary.

Let therefore
$$4 = r$$
, and $r + z = x$.
Then $\begin{vmatrix} x^3 = r^3 + 3r^2z + 3rz^2 + z^3 \\ x^2 = r^2 + 2rz + z^2 \end{vmatrix} = 90$.
 $x = r + z$

And by rejecting the terms z^3 , $3rz^2$ and z^2 , as small in:

comparison with z, we shall have

$$Whence z = \frac{90 - r^3 - r^2 - r}{3r^2 + 2r + 1} = \frac{90 - 64 - 16 - 4}{48 + 3 + 1} = \frac{6}{57} = .10.$$
And consequently $x = 4.1$, nearly.

Again, if 4.1 be substituted in the place of r, in the last equation, we shall have

z =
$$\frac{90 - r^3 - r^2 - r}{3r^2 + 2r + 1} = \frac{90 - 68.921 - 16.81 - 4.1}{50.43 + 6.2 + 1} = .00283$$
;

And consequently x = 4.1 + .00283 = 4.10283 for a second.

approximation.

And if the first four figures, 4.102, of this number be again substituted for r, in the same equation, a still nearer value of the root will be obtained; and so on, as far as may be thought necessary.

2. Given $x^2 + 20x = 100$, to find the value of x by approximation.

Ans. x = 4.1421356

3. Given $x^3+9x^2+4x=80$, to find the value of x by approximation.

Ans. x=2.4721359

4. Given $x^4 - 38x^3 + 210x^2 + 538x + 289 = 0$, to find the value of x by approximation.

Ans. x 30.53565375

5. Given $x^5 + 6x^4 - 10x^3 - 112x^2 - 207x + 110 = 0$, to find the value of x by approximation.

Ans. 4.46410161

The roots of equations, of all orders, can also be determined, to any degree of exactness, by means of the following easy rule of double position; which though it has not been generally employed for this purpose, will be found in some respects, superior to the former, as it can be applied, at once, to any unreduced equation, consisting of surds, or compound quantities, as readily as if it had been brought to its usual form.

RULE.

Find, by trial, two numbers as near the true root as possible, and substitute them in the given equation instead of the unknown quantity, noting the results that are obtained from each.

Then, as the difference of these results is to the difference of the two assumed numbers, so is the difference between the true result, given by the question, and either of the former, to the correction of the number belonging to the result used; which correction being added to that number when it is too little, or subtracted from it when it is too great, will give the root required nearly.

And if the number thus determined, and the nearest of the two former, or any other that appears to be more accurate, be now taken as the assumed roots, and the operation be repeated as before, a new value of the unknown quantity will be obtained still more correct than the first; and so on, proceeding in this manner, as far as may be judged necessary.*

^{*} The above rule for Double Position, which is much more simple and commodious than the one commonly employed for this purpose, is the same as that which was first given at p. 311 of the octavo edition of my Arithmetic, published in 1810.

To this we may farther add, that when one of the roots of an equation has been found, either by this method or the former, the rest may be determined as follows:

Bring all the terms to the left hand side of the equation, and divide the whole expression, so formed, by the difference between the unknown quantity (x) and the root first found; and the resulting equation will then be depressed a degree lower than the given one.

Find a root of this new equation, by approximation, as in the first instance, and the number so obtained will be a second root of the original equation.

Then, by means of this root, and the unknown quantity, depress the se-

Then, by means of this root, and the unknown quantity, depress the second equation a degree lower, and thence find a third root; and so on, till the equation is reduced to a quadratic; when the two roots of this, together with the former, will be the roots of the equation required.

Thus in the equation $x_3 - 15x_2 + 63x = 50$, the first root is found, by approximation to be 1.02804. Hence,

 $x-1.02804(x_3-15x_2+63x-50(x_2-13.97196x+48.63627=0.$

And the two roots of the quadratic equation, $\alpha^2 - 13.97196\alpha = -48.63627$, found in the usual way, are 6.57653 and 7. 39543.

So that the three roots of the given cubic equation x3-15x2+63x=50, are 1.02804, 6.37553, and 7.39543; their sum being=15, the coefficient of the second term of the equation, as it ought to be when they are right.

EXAMPLES.

1. Given $x^3+x^2+x=100$, to find an approximate value of x.

Here it is soon found, by a few trials, that the value of

x lies between 4 and 5.
Hence, by taking these as the two assumed numbers, the

And consequently x=4+.225=4.225, nearly.

Again, if 4.2 and 4.3 be taken as the two assumed num-

bers, the operation will stand thus :

	Sup. Secon				nd Sun.	
4.2 .		x			4.3	
17.64		x^2			18.49	
74.088 .		x^3			79.507	
	_				-	
102.297		4.3			102.297	
95.928		4.2			100	
	First Sup. 4.2 17.64 .74.088 95.928 102.297	First Sup. 4.2 17.64 .74.088 .95.928 Re 102.297 95.928	First Sup. 4.2 . x 17.64 . x ² 74.088 . x ³ 95.928 Results 102.297 . 4.3 95.928 . 4.2	First Sup. Se 4.2 . x 17.64 . x ² .74.088 . x ³ 95.928 Results 102.297 . 4.3 .	First Sup. Seco 4.2 . x	First Sup. Second Sup. 4.2 . x . 4.3 17.64 . x ² . 18.49 74.088 . x ³ . 79.507 95.928 Results 102.297 102.297 . 4.3 . 102.297 95.928 . 4.2 . 100

And consequently x=4.3-.036=4.264, nearly.

Again, let 4.264 and 4.265 be the two assumed numbers: then

First Sup.	1				Second Sup.			
			x			4.265		
18.181696			x^2			18.190225		
77.526752			x^3			77.581310		
99 979448		Ð	0611	140		100.03653		

Therefore

100.036535 4.265 100 99.972448 4.264 99.972448

.064087 : .001 :: .027552 : .0004299 And consequently

x=4.264+.0004299=4.2644299, very nearly.

2. Given $(\frac{1}{2}x^2-15)^2+x\sqrt{x}=90$, to find an approxi-

mate value of x.

Here, by a few trials, it will be soon found, that the value of x lies between 10 and 11; which let, therefore, be the two assumed numbers, agreeably to the directions given in the rule.

Hence

And consequently x=11 ... 31.122 : .482 And consequently x=11 ... 482=10.518.

Again, let 10.5 and 10.6 be the two assumed numbers,

Then

First Sup.

Second Sup.

49.7025 . . $(\frac{1}{8}x^2 - 15)^2$. . 55.830784 34.0239 . . $x\sqrt{x}$. . 34.511099

83.7264 . . Results . . 90.341883 Hence

90.341883 . . 10.6 . . 90.341883 83.7264 . . 10.5 . . 90.

6.615483 : .1 :: .341883 : .0051679 And consequently

x=10.6—.0051679=10.5948321, very nearly.

EXAMPLES FOR PRACTICE.

1. Given $x^3 + 10x^2 + 5x = 2600$, to find a near approximate value of x.

Ans. = 11.00673

2. Given $2x^4 - 16x^3 + 40x^2 - 30x + 1 = 0$, to find a near value of x. Ans. x = 1.284724

3. Given $x^5+2x^4+3x^3+4x^2+5x=54321$, to find the value of x.

4. Given $\sqrt[3]{(7x^3+4x^2)} + \sqrt{(20x^2-10x)} = 28$, to find the value of x. Ans. 4.510661

5. Given $\sqrt{(144x^2-(x^2+20)^2)}+\sqrt{(196x^2-(x^2+24)^2)}$ = 114, to find the value of x. Ans. 7.123883

OF EXPONENTIAL EQUATIONS.

An exponential quantity is that which is to be raised to some unknown power, or which has a variable quantity for its index; as

 a^x , $a^{\overline{x}}$, a^x , or $x^{\overline{x}}$, &c.

And an exponential equation is that which is formed between any expression of this kind and some other quantity, whose value is known; as

 $a^{x} = b, x^{x} = a, \&c.$

Where it is to be observed, that the first of these equations, when converted into logarithms, is the same as

 $x \log a = b$, or $x = \frac{\log b}{\log a}$; and the second equation $x^x = a$ is the same as $x \log x = \log a$.

In the latter of which cases, the value of the unknown quantity x may be determined, to any degree of exactness, by the method of double position, as follows:

BULE.

Find by trial, as in the rule before laid down, two numbers as near the number sought as possible, and substitute them in the given equation

 $x \log_{\bullet} x = \log_{\bullet} a$,

instead of the unknown quantity, noting the results obtain-

ed from each.

Then, as the difference of these results is to the difference of the two assumed numbers, so is the difference between the true result, given in the question, and either of the former, to the correction of the number belonging to the result used; which correction being added to that number, when it is too little, or subtracted from it, when it is too great, will give the root required, nearly.

And, if the number, thus determined, and the nearest of the two former, or any other that appears to be nearer, be taken as the assumed roots and the operation be repeated as before, a new value of the unknown quantity will be obtained still more correct than the first; and so on, proceeding in this manner, as far as may be thought

necessary.

EXAMPLES.

1. Given $x^x = 100$, to find an approximate value of x. Here, by the above formula, we have

 $x \log_{100} x = \log_{100} = 2$.

And since x is readily found, by a few trials, to be nearly in the middle between 3 and 4, but rather nearer the latter than the former, let 3.5 and 3.6 be taken for the two assumed numbers.

Then log. 3.5=.5440680, which, being multiplied by

3.5, gives 1.904238=first result; And log. 3.6= 5563025, which, being multiplied by 3.6, gives 2.002689 for the second result.

Whence

2.002689 . . 3.6 . . 2 002689 1.904238 . . 3.5 . . 2.

.098451 : .1 :: .002689 : .00273

for the first correction; which, taken from 3.6, leaves x=3.59727, nearly.

And as this value is found, by trial, to be rather too small, let 3.59727 and 3.59728 be taken as the two assumed numbers.

Then log. 3.59728=0.555974243134677 to 15 places The log. 3.59727=0.555973035847267 to 15 places which logarithms multiplied by their respective numbers give the following products:

1.999995025343512 both true to the last figure.

Therefore the errors are 4974656488 and 14877337702 and the difference of errors 9902681214

Now since only 6 additional figures are to be obtained, we may omit the three last figures in these errors; and state thus: as difference of errors 9902681: difference of sup. 1:: error 4974656: the correction 502354, which united to 3.59728 gives us the true value of $x = 3.59728502354^*$.

- 2. Given $x^x = 2000$, to find an approximate value of x.

 Ans. x = 4.82782263
- 3. Given $(6x)^x = 96$, to find the approximate value of x.

 Ans. x=1.8826432
- 4. Given $x^x = 123456789$, to find the value of x.

 Ans. 8.6400268
- 5. Given $x^x x = (2x x^x)^{\frac{1}{x}}$, to find the value of x.

 Ans. x = 1.747933.

OF THE

BINOMIAL THEOREM.

The binomial theorem is a general algebraical expression, or formula, by which any power, or root of a given quantity, consisting of two terms, is expanded into a series; the form of which, as it was first proposed by Newton, being as follows:

^{*} The correct answer to this question has been first given by *Doctor Advain*, in his edition of Hutton's Mathematics, who plainly proves that Hutton's answer, which is the same as Bonnycaste, is incorrect; See Hutton's Mathematics, Vol. 1. p. 263. N. Y. Edit.

$$(P+PQ)_{n}^{m} = P_{n}^{m} \left[1 + \frac{n}{n}Q + \frac{m}{n}(\frac{m-n}{2n})Q^{2} + \frac{m}{n}(\frac{m-n}{2n})\right]$$

$$(\frac{m-2n}{3n})Q^{3}, \&c.$$

$$Or,$$

$$(P+PQ)_{n}^{m} = P_{n}^{m} + \frac{m}{n}AQ + \frac{m-n}{2n}BQ + \frac{m-2n}{3n}CQ + \frac{m-3n}{4n}DQ, \&c.$$

Where P is the first term of the binomial, Q the second term divided by the first, $\frac{m}{n}$ the index of the power, of root, and A, B, C, &c. the terms immediately preceding those in which they are first found, including their signs + or -.

Which theorem may be readily applied to any particular case, by substituting the numbers, or letters, in the given example, for p, q, m, and n, in either of the above formulæ, and then finding the result according to the rule.*

$$\frac{m}{(a+x)^{\frac{m}{n}}} = \frac{m}{n} \left[1 + \frac{m}{n} \frac{x}{a} \right] + \frac{m}{n} \frac{m-n}{2n} \frac{x}{a} + \frac{m-n}{n} \frac{m-n}{2n} \frac{m-2n}{3n} \frac{m-2n}{a} \frac{m-2n}{3n} \frac{m-2n}{3n} \frac{m-2n}{a} \frac{m-2n}{3n} \frac{m-2n}{a} \frac{m-2n}{3n} \frac{m-2n}{a} \frac{m-2n}{3n} \frac{m-2n}{a} \frac{m-2n}{3n} \frac{m-2n}{a} \frac{m-2n}{3n} \frac{m-2n}{3n} \frac{m-2n}{a} \frac$$

$$a\frac{m}{n}\left[1+\frac{m}{n}\left(\frac{x}{a+x}\right)+\frac{m}{n}\frac{m+n}{2n}\left(\frac{x}{a+x}\right)^2+\frac{m}{n}\frac{m+n}{2n},\frac{m+2n}{3n}\left(\frac{x}{a+x}\right)^3\&c.\right]$$
Or, $(a+x)\frac{m}{n}=$

$$2a\frac{m}{n}[1-\frac{m}{n}(\frac{a-x}{a+x})+\frac{m}{n}\frac{m+n}{2n}(\frac{a-x}{a+x})2-\frac{m}{n}\frac{m+n}{2n}\frac{n+2n}{3n}(\frac{a-x}{a+x})3] \ \&c.$$
 It may here also be observed, that if m be made to represent any whole, or

It may here also be observed, that if m be made to represent any whole, or fractional number, whether positive or negative, the first of these expressions may be exhibited in a more simple form

$$(a+x)^m = a^m + ma^{m-1}x + \frac{m(m-1)}{1 \cdot 2}a^{m-2}x^2 + \frac{m(m-1)(m-2)}{1 \cdot 2}a^{m-3}x^3$$

$$m(m-1)(m-2) \qquad [m-(n-1)]a^nx^{m-n}$$

^{*} This celebrated theorem, which is of the most extensive use in algebra...
and various other branches of analysis, may be otherwise expressed as follows:

Where the last term is called the general term of the series, because if $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{3}$ c. be substituted successively for n, it will give all the rest.

EXAMPLES.

1. It is required to convert $(a^2+x)^{\frac{1}{2}}$ into an infinite series.

Here $r=a^2$, $q=\frac{x}{a^2}$, $\frac{m}{n}=\frac{1}{2}$, or m=1, and n=2; whence

$$\begin{split} & \frac{m}{r_n} = \left(a^2\right)^{\frac{m}{n}} = \left(a^2\right)^{\frac{1}{2}} = a = \lambda, \\ & \frac{m}{n} \Delta \mathbf{q} = \frac{1}{2} \times \frac{a}{1} \times \frac{x}{a^2} = \frac{x}{2a} = \mathbf{B}, \\ & \frac{m-n}{2n} \mathbf{B} \mathbf{q} = \frac{1-2}{4} \times \frac{x}{2a} \times \frac{x}{a^2} = -\frac{x^2}{2.4a^3} = \mathbf{c}, \\ & \frac{m-2n}{3n} \mathbf{c} \mathbf{q} = \frac{1-4}{6} \times -\frac{x^2}{2.4a^3} \times \frac{x}{a^2} = \frac{3x^3}{2.4.6a^5} = \mathbf{p}, \\ & \frac{m-3n}{4n} \mathbf{p} \mathbf{q} = \frac{1-8}{6} \times \frac{3x^3}{2.4.6a^5} \times \frac{x}{a^2} = -\frac{3.5x^4}{2.4.6.8a^7} = \mathbf{e}, \\ & \frac{m-4n}{5n} \mathbf{e} \mathbf{q} = \frac{1-8}{10} \times -\frac{3.5x^4}{2.4.6.8a^7} \times \frac{x}{a^2} = \frac{3.5.7x^5}{2.4.6.810a^9} = \mathbf{f}, \\ & \&c. \end{split}$$

Therefore $(a^2+x)^{\frac{1}{2}} = a + \frac{x}{2a} - \frac{x^2}{2.4a^3} + \frac{3x^3}{2.4.6a^5} - \frac{3.5x^4}{2.4.6.8a^7} + \frac{3.5.7x^5}{2.4.6.8.10a^9} - &c.$

Where the law of formation of the several terms of the series is sufficiently evident.

2. It is required to convert $\frac{1}{(a+b)^2}$, or its equal $(a+b)^2$, into an infinite series.

Here r=a, $q=\frac{b}{a}$, and $\frac{m}{n}=-2$, or m=-2, and n=1;

$$\frac{m}{r_{n}} = (a)^{\frac{m}{n}} = a^{-2} = \frac{1}{a^{2}} = \lambda,$$

$$\frac{m}{r_{n}} \land Q = -\frac{2}{1} \times \frac{1}{n^{2}} \times \frac{b}{a} = -\frac{2b}{a^{3}} = b.$$

$$\begin{split} \frac{m-n}{2n} & \text{Eq} = \frac{-2-1}{2} \times -\frac{2b}{a^3} \times \frac{b}{a} = \frac{3b^2}{a^4} = \epsilon, \\ \frac{m-2n}{3n} & \text{eq} = \frac{-2-2}{3} \times \frac{3b^2}{a^4} \times \frac{b}{a} = -\frac{4b^3}{a^5} = D, \\ \frac{m-3n}{4n} & \text{eq} = \frac{-2-3}{4} \times -\frac{4b^3}{a^5} \times \frac{b}{a} = \frac{5b^4}{a^6} = E, \\ & \&c. & \&c. & \&c. & \&c. \end{split}$$

Consequently $\frac{1}{(a+b)^2} = \frac{1}{a^2} - \frac{2b}{a^3} + \frac{3b^2}{a^4} - \frac{4b^3}{a^5} + \frac{5b^4}{a^6} &c.$

3. It is required to convert $\frac{a^2}{(a^2-x)^{\frac{1}{2}}}$, or its equal a^2 $(a^2-x)^{\frac{1}{2}}$, into an infinite series.

 $p=x^2, q=-\frac{x}{a^2}$, and $\frac{m}{n}=-\frac{1}{2}$, or m=-1 and n=2;

whence

$$\frac{m}{n} = (a^{2})^{\frac{m}{n}} = (a^{2})^{\frac{-1}{2}} = \frac{1}{a} = \lambda,$$

$$\frac{m}{n} A_{Q} = -\frac{1}{2} \times \frac{1}{a} \times -\frac{x}{a^{2}} = \frac{x}{2a^{3}} = B,$$

$$\frac{m-n}{2n} B_{Q} = \frac{-1-2}{4} \times \frac{x}{2a^{3}} \times -\frac{x}{a^{2}} = \frac{3x^{2}}{2.4a^{5}} = c,$$

$$\frac{m-2n}{3n} c_{Q} = \frac{-1-4}{6} \times \frac{3x^{2}}{2.4a^{5}} \times -\frac{x}{a^{2}} = \frac{3.5x^{3}}{2.4.6a^{7}} = D,$$

$$-3n = -1-6 \cdot 3.5x^{3} \quad x \quad 3.5.7x^{4}$$

 $\frac{an-3n}{4n}\log_{\mathbb{Q}} = \frac{-1-6}{8} \times \frac{3.5x^{2}}{2.4.6a^{7}} \times \frac{x}{a^{2}} = \frac{3.5.7x^{4}}{2.4.6.8a^{9}} = \varepsilon,$ &c. &c. &c.

Therefore

 $\frac{1}{(a^2-x)^2} = \frac{1}{a} + \frac{1}{2} \left(\frac{x}{a^3}\right) + \frac{3}{2.4} \left(\frac{x^2}{a^5}\right) + \frac{3.5}{2.4.6} \left(\frac{x^3}{a^7}\right) + \frac{3.5.7}{2.4.6.8} \left(\frac{x^4}{a^8}\right)$ &c.

 $\frac{\alpha^{3}}{(a^{2}-x)^{3}} = \alpha + \frac{1}{2} \left(\frac{x}{a}\right) + \frac{3}{2.4} \left(\frac{x^{2}}{a^{3}}\right) + \frac{3.5}{2.4.6} \left(\frac{x^{3}}{a^{3}}\right) + \frac{3.5.7}{2.4.6.8} \left(\frac{x^{4}}{a^{7}}\right)$

4. It is required to convert $\sqrt[3]{9}$, or its equal $(8+1)\frac{1}{9}$ into an infinite series.

Here
$$\mathbf{r} = 8$$
, $\mathbf{q} = \frac{1}{8}$, and $\frac{m}{n} = \frac{1}{3}$, or $m = 1$ and $n = 3$;

Whence
$$\mathbf{r}^n = (8)^n = 8^{\frac{1}{3}} = 2 = \kappa.$$

$$\frac{m}{n} \mathbf{A} \mathbf{q} = \frac{1}{3} \times \frac{2}{1} \times \frac{1}{2^3} = \frac{1}{3.2^2} = \mathbf{B},$$

$$\frac{m - n}{2n} \mathbf{B} \mathbf{q} = \frac{1 - 3}{6} \times \frac{1}{3.2^2} \times \frac{1}{2^3} = -\frac{1}{3.6.2^4} = \mathbf{c},$$

$$\frac{m - 2n}{2n} \mathbf{c} \mathbf{q} = \frac{1 - 6}{9} \times -\frac{1}{3.6.2^4} \times \frac{1}{2^3} = \frac{5}{3.6.9.2^7} = \mathbf{p},$$

$$\frac{m - 3n}{4n} \mathbf{p} \mathbf{q} = \frac{1 - 9}{12} \times \frac{5}{3.6.9.2^7} \times \frac{1}{2^3} = -\frac{5.8}{3.6.9.12.2^{10}} = \mathbf{E},$$

$$\frac{m - 4n}{5n} \mathbf{e} \mathbf{q} = \frac{1 - 12}{15} \times -\frac{5.8}{3.6.9.12.2^{10}} \times \frac{1}{2^3} = \frac{5.8.11}{3.6.9.12.15.2^{13}}$$
&c. &c.

$$2 + \frac{1}{3.2^{2}} - \frac{1}{3.6.2^{4}} + \frac{5}{3.6.9.2^{7}} + \frac{5.8}{3.6.9.12} + \frac{5.8.11}{3.6.9.12} + \frac{5.8.11}{3.6.9.12} + \frac{3.6.9.12}{3.6.9.12} + \frac{3$$

5. It is required to convert $\sqrt{2}$, or its equal $\sqrt{(1+1)}$, into an infinite series.

Ans.
$$1 + \frac{1}{2} - \frac{1}{2.4} + \frac{3}{2.4.6} - \frac{3.5}{2.4.6.8} + \frac{3.5.7}{2.4.6.8.10} &c.$$

6. It is required to convert $\sqrt[3]{7}$, or its equal $(8-1)^{\frac{1}{3}}$, into an infinite series.

Ans.
$$2 - \frac{1}{3.2^2} - \frac{1}{3.6.2^4} - \frac{5}{3.6.9.2^7} - \frac{5.8}{3.6.9.12.2^{10}} - &c.$$

7. It is required to convert $\sqrt[5]{240}$, or its equal $(243-3)^{\frac{1}{5}}$, into an infinite series.

Ans.
$$3 - \frac{1}{5.3^3} - \frac{4}{5.10.3^7} - \frac{4.9}{5.10.15.20.3^{15}} &c.$$

8. It is required to convert $(a\pm x)^{\frac{1}{2}}$ into an infinite series. Ans. $a^{\frac{1}{2}}\left\{1\pm \frac{x}{2a} - \frac{x^2}{2\cdot 4a^2} \pm \frac{3x^3}{2\cdot 4\cdot 6a^3} - \frac{3\cdot 5x^4}{2\cdot 4\cdot 6\cdot 8a^4} \pm &c.\right\}$ 9. It is required to convert $(a\pm b)^{\frac{1}{3}}$ into an infinite series.

Ans.
$$a^{\frac{1}{3}} \left\{ 1 \pm \frac{b}{3a} - \frac{2b^2}{3.6a^2} \pm \frac{2.5b^3}{3.6.9.a^3} - \frac{2.5.8b^4}{3.6.9.12a^4} \pm &c. \right\}$$

10. It is required to convert $(a-b)^{\frac{1}{4}}$ into an infinite series.

ries.
Ans.
$$a^{\frac{1}{4}} \left\{ 1 - \frac{b}{4a} - \frac{3b^2}{4.8a^2} - \frac{3.7b^3}{4.8.12a^3} - \frac{3.7.11b^4}{4.8.12.16a^4} - &c. \right\}$$

11. It is required to convert $(a+x)^{\frac{2}{3}}$ into an infinite series.

Ans.
$$a^{\frac{2}{3}}$$
 $\left\{1 + \frac{2x}{3a} - \frac{x^2}{9a^2} + \frac{4x^3}{9^2a^3} - \frac{4.7x^3}{9^212a^4} + \frac{4.7.10x^5}{9^2.12.15a^5} - \&c.\right\}$

12. It is required to convert $(1-x)^{\frac{2}{5}}$ into an infinite series.

Ans.
$$1 - \frac{2x}{5} - \frac{2.3x^3}{5.10} - \frac{2.3.8x^3}{5.10.15} - \frac{2.3.8.13x^4}{5.10.15.20} - &c.$$

13. It is required to convert $\frac{1}{(a\pm x)^{\frac{1}{2}}}$, or its equal

 $(a\pm x)^{-\frac{1}{2}}$ into an infinite series.

Ans.
$$\frac{1}{a^{\frac{1}{2}}} \left\{ 1 \mp \frac{x}{2a} + \frac{3x^2}{2.4a^2} \mp \frac{3.5x^3}{2.4.6a^3} + \frac{3.5.7x^4}{2.4.6.8a^4} \mp &c. \right\}$$

14. It is required to convert $\frac{a}{(a\pm x)^{\frac{1}{3}}}$, or its equal

 $(a\pm x)^{-\frac{1}{3}}$ into an infinite series.

Ans.
$$a^{\frac{2}{3}} \left\{ 1 \mp \frac{x}{3a} + \frac{4x^2}{3.6a^2} \mp \frac{4.7x^3}{3.69a^3} + \frac{4.7.10x^4}{3.6.9.12n^4} \mp &c. \right\}$$

15. It is required to convert $\frac{1}{(1+x)^{\frac{1}{5}}}$, or its equal

 $(1+x)^{\frac{1}{5}}$ into an infinite series.

Ans.
$$1 - \frac{x}{5} + \frac{6x^3}{5.10} - \frac{6.9x^3}{5.10.15} + \frac{6.11.16x^4}{5.10.15.20} - &c.$$

16. It is required to convert $\left(\frac{a+x}{a-x}\right)^{\frac{1}{2}}$, or its equal $(a+x)\left(a^2-x^2\right)^{-\frac{1}{2}}$, into an infinite series.

Ans. $1+\frac{x}{a}+\frac{x^2}{2u^2}+\frac{x^3}{2a^3}+\frac{3x^4}{8a^4}+\frac{3x^5}{8a^5}+\frac{5x^6}{16x^6}+\frac{5x^7}{16a^7}$ &c.

OF THE

INDETERMINATE ANALYSIS.

In the common rules of Algebra, such questions are usually proposed as require some certain or definite answer; in which case, it is necessary that there should be as many independent equations, expressing their conditions, as there are unknown quantities to be determined; or

otherwise the problem would not be limited.

But in other branches of the science, questions frequently arise that involve a greater number of unknown quantities than there are equations to express them; in which instances they are called indeterminate or unlimited problems; being such as usually admit of an indefinite number of solutions; although, when the question is proposed in integers, and the answers are required only in whole positive numbers, they are, in some cases, confined within certain limits, and in others, the problem may become impossible.

PROPLEM 1.

To find the integral values of the unknown quantities x and y in the equation

 $ax-by=\pm c$, or ax+by=c.

Where a and b are supposed to be given whole numbers, which admit of no common divisor, except when it is also a divisor of c.

BULE.

1. Let wh denote a whole, or integral number; and reduce the equation to the form

$$x = \frac{by + c}{a}wh$$
, or $x = \frac{c - by}{a}wh$.

2. Throw all whole numbers out of that of these two expressions, to which the question belongs, so that the numbers d and e in the remaining parts, may be each less than a; then

$$\frac{dy \pm e}{a} = wh$$
, or $\frac{e - dy}{a} = wh$.

3. Take such a multiple of one of these last formulæ, corresponding with that above mentioned, as will make the coefficient of y nearly equal to a, and throw the whole numbers out of it as before.

Or find the sum or difference of $\frac{ay}{a}$, and the expression above used, or any multiple of it that comes near $\frac{ay}{a}$, and the result, in either of these cases, will still be =wh, a whole number.

4. Proceed in the same manner with this last result; and so on, till the coefficient of y becomes = 1, and the remainder = some number r; then

$$\frac{y\pm r}{a} = wh. = p$$
, and $y=ap\mp r$,

Where p may be o, or any integral number whatever, that makes y positive; and, as the value of y is now known, that of x may be found from the given equation, when the question is possible*.

Note. Any indeterminate equation of the form

$$ax-by=\pm c$$
,

^{*} This rule is founded on the sornous principle, that the sum, difference, or product of any two whole numbers, is a whole number; and that, if a number divides the whole of any other number and a part of it, it will slee divide the remaining part.

in which a and b are prime to each other, is always possible, and will admit of an infinite number of answers in whole numbers.

But if the proposed equation be of the form ax+by=c,

the number of answers will always be limited; and, in some cases, the question is impossible; both of which circumstances may be readily discovered, from the mode of solution above given*.

EXAMPLES.

1. Given 19x-14y=11, to find x and y in whole numbers.

Here
$$x = \frac{14y+11}{19} = wh$$
, and also $\frac{19y}{19} = wh$.
Whence, by subtraction, $\frac{19y}{19} = \frac{14y+11}{19} = \frac{5y-11}{19} = wh$.
Also, $\frac{5y-11}{19} \times 4 = \frac{20y-44}{19} = y-2 + \frac{y-6}{19} = wh$.
And by rejecting $y=2$, which is a whole number, $\frac{y-6}{19} = wh = p$.

Whence we have y=19p+6. And $x = \frac{14y+11}{19} = \frac{14(19p+6)+11}{19} = \frac{266p+95}{19} = \frac{19}{19}$

^{*} That the coefficients a and b, when these two formulæ are possible, should have no common divisor, which is not at the same time, a divisor of c, is evident; for if $a \in md$, and b = me, we shall have $ax \pm by = mdx \pm mey = c$; and consequently $dx + \epsilon y = \frac{c}{m}$. But d, e, x, y, being supposed to be whole

numbers $\frac{c}{m}$ must also be a whole number, which it cannot be, except when m

is a divisor of c. Hence, if it were required to pay 100L in guineas and moidores only, the question would be impossible; since, in the equation 21x + 27y = 2000 which represents the conditions of the problem, the coefficients, 21 and 27, are each divisible by 3, whilst the absolute term 2000 is not divisible by it. See my Treaties of Algebra, for the method of resolving questions of this kind, by means of Continued Fractions.

14p + 5.

Whence, if p be taken =0 we shall have x=5 and y =6, for their least values; the number of solutions being obviously indefinite.

2. Given 2x+3y=25, to determine x and y in whole

positive numbers.

Here
$$x = \frac{25 - 3y}{2} = 12 - y + \frac{1 - y}{2}$$

Hence, since x must be a whole number, it follows that $\frac{1-y}{a}$ must also be a whole number.

Let therefore
$$\frac{1-y}{2} = wh = p$$
;
Then $1-y=2p$, or $y=1-2p$.

And since
$$x = 12 - y + \frac{1 - y}{2} = 12 - (1 - 2p) + p = 12 + 3p - 1$$
,

We shall have x=11+3p, and y=1-2p; Where p may be any whole number whatever, that will render the values of x and y in these two equations positive:

But it is evident, from the value of y, that p must be either 0 or negative; and consequently, from that of x, that it must be 0, -1, -2, or -3.

Then $\begin{cases} x=11, x=8, x=5, x=2, \\ y=1, y=3, y=5, y=7, \end{cases}$ Which are all the answers in whole positive numbers that the question admits of.

3. Given 3x=8y-16 to find the values of x and y in whole numbers.

Here
$$x = \frac{8y - 16}{3} = 2y - 5 + \frac{2y - 1}{3} = wh$$
; or $\frac{2y - 1}{3} = wh$.
Also $\frac{2y - 1}{3} \times 2 = \frac{4y - 2}{3} = y + \frac{y - 2}{3} = wh$.

Or, by rejecting y, which is a whole number, there will remain $\frac{y-2}{2} = wh = p$.

And
$$x = \frac{\text{Therefore } y = 3p + 2,}{3} = \frac{8y - 16}{3} = \frac{8(3p + 2) - 16}{3} = \frac{24p}{3} = 8p.$$

Where, if p be put=1, we shall have x=8 and y=5, for their least values; the number of answers being, as in the first question, indefinite.

4. Given 21x+17y=2000, to find all the possible val-

ues of x and y in whole numbers.

Here
$$x = \frac{2000 - 17y}{21} = 95 + \frac{5 - 17y}{21} = \pi h$$
.;
Or, omitting the 95, $\frac{5 - 17y}{21} = \pi h$.;

Consequently, by addition, $\frac{21y}{21} + \frac{5 - 17y}{21} = \frac{4y + 5}{21} = wh$.; Also, $\frac{4y + 5}{21} \times 5 = \frac{20y + 25}{21} = 1 + \frac{4 + 20y}{21} = wh$.;

Or, by rejecting the whole number 1, $\frac{4+20y}{21} = wh$.

And, by subtraction, $\frac{21y}{21} - \frac{4+20y}{21} = \frac{y-4}{21} = wh = p$;

Whence
$$y=21p+4$$
,
And $a=\frac{2000-17y}{21}=\frac{2000-17(21p+4)}{21}=92-17p$.

Where if p be put=0, we shall have the least value of y=4, and the corresponding, or greatest value of x=92.

And the rest of the answers will be found by adding 21 continually to the least value of y, and subtracting 17 from the greatest value of x; which being done we shall obtain the six following results:

$$x = 92 \mid 75 \mid 58 \mid 41 \mid 24 \mid 7$$

 $y = 4 \mid 25 \mid 46 \mid 67 \mid 88 \mid 109$

These being all the solutions, in whole numbers, that the

question admits of.

Note. 1. When there are three or more unknown quantities, and only one equation by which they can be determined, as

$$ax+by+cz=d$$
.

it will be proper first to find the limit of the quantity that has the greatest coefficient, and then to ascertain the different values of the rest, by separate substitutions of the several values of the former, from 1 up to that extent, as in the following question.

5. Given 3x+5y+7z=100, to find all the different va-

lues of x, y, and z, in whole numbers*.

Here each of the least integer values of x and y are 1, by the question; whence it follows, that

 $z = \frac{100 - 5 - 3}{7} = \frac{100 - 3}{7} = \frac{92}{7} = 13\frac{1}{7}.$

Consequently z cannot be greater than 13, which is also the limit of the number of answers; though they may be considerably less.

By proceeding, therefore, as in the former rule, we shall

bave

have
$$x = \frac{100 - 5y - 7z}{3} = 33 - y - 2z + \frac{1 - 2y - z}{3} = wh.;$$
And, by rejecting $33 - y - 2z$,
$$\frac{1 - 2y - z}{3} = wh.; \text{ or } \frac{3y}{3} + \frac{1 - 2y - z}{3} = \frac{y + 1 - z}{3} = wh.;$$
Whence $\frac{y + 1 - z}{3} = p$.
And $y = 3p + z - 1$;

And consequently, putting p=0, we shall have the least value of y=z-1; where z may be any number, from 1 up to 13, that will answer the conditions of the question.

When, therefore,
$$z=2$$
 we have $y=1$,
And $x=\frac{100-19}{3}=27$.

^{*} If any indeterminate equation, of the kind above given, has one or more of its coefficients, as c, negative, the equation may be put under the form

av + by = d + cz, in which case it is evident that an indefinite number of values may be given to the second side of the equation by means of the indefinite quantity z; and consequently, also, to x and y in the first. And if the coefficients a, b, c, in any such equation, have a common divisor, while d has not, the question, as in the first case, becomes impossible.

Hence, by taking z=2, 3, 4, 5, &c. the corresponding values of x and y, together with those of z, will be found to be as below.

Which are all the integral values of x, y, and z, that can be obtained from the given equation.

Note 2. If there be three unknown quantities, and only

ax+by+cz=d, and ex+fy+gz=h,

exterminate one of these quantities in the usual way, and find the values of the other two from the resulting equation, as before.

Then, if the values, thus found, be separately substituted, in either of the given equations, the corresponding values of the remaining quantities will likewise be determined: thus,

6. Let there be given x-2y+z=5, and 2x+y-z=7,

to find the values of x, y, and z.

Here, by multiplying the first of these equations by 2, and subtracting the second from the product, we shall have

$$3z - 5y = 3$$
, or $z = \frac{3+5y}{3} = 1+y+\frac{2y}{3} = wh$.;

And consequently $\frac{2y}{3}$, or $\frac{3y}{3} - \frac{2y}{3} = \frac{y}{3} = wh = p$.

Whence y=3p.

And, by taking p=1, 2, 3, 4, &c. we shall have y=3, 6, 9, 12, 15, &c. and z=6, 11, 16, 21, 26, &c.

But from the first of the two given equations

x=5+2y-z; whence, by substituting the above values for y and z, the results will give x=5, 6, 7, 8, 9, &c.

And therefore the first six values of x, y, and z, are as

below:

Where the law by which they can be continued is sufficiently obvious.

EXAMPLES FOR PRACTICE.

1. Given 3x=8y-16, to find the least values of x and y in whole numbers. Ans. x=8, y=5

2. Given 14x=5y+7, to find the least values of x and y in whole numbers

Ans. x=3, y=7

3. Given 27x=1600-16y, to find the least values of and y in whole numbers. Ans. x=48, y=19

-4. It is required to divide 100 into two such parts, that one of them may be divisible by 7, and the other by 11.

Ans. The only parts are 56 and 44

- 5. Given 9x+13y=2000, to find the greatest value of x and the least value of y in whole numbers.
- Ans. x=215, y=5 6. Given 11x+5y=254, to find all the possible values of x and y in whole numbers.

Ans. x=19, 14, 9, 4; y=9, 20, 31, 42

7. Given 17x+19y+21z=400, to find all the answers in whole numbers which the question admits of.

Ans. 10 different answers

Ans. 10 different answers

8. Given 5x+7y+11z=224, to find all the possible values of x, y, and z, in whole positive numbers.

Ans. The number of answers is 59

9. It is required to find in how many different ways it is possible to pay 20*l*. in half-guineas and half-crowns, without using any other sort of coin?

Ans. 7 different ways

10. I owe my friend a shilling, and have nothing about me but guineas, and he has nothing but louis-d'ors; how must I contrive to acquit myself of the debt, the louis being valued at 17s. a piece, and the guineas at 21s.?

Ans. I must give him 13 guineas, and he must

give me 16 louis

11. How many gallons of British spirits, at 12s., 15s., and 12s. a gallon, must a rectifier of compounds take to make a mixture of 1000 gallons, that shall be worth 17s. a gallon?

Ans. $111\frac{1}{6}$, at 12s., $111\frac{7}{6}$ at 15s .: and $777\frac{7}{2}$ at 18ϵ

PROBLEM II.

To find such a whole number, as, being divided by other given numbers, shall leave given remainders.

RULE.

1. Call the number that is to be determined x, the numbers by which it is to be divided a, b, c, &c. and the given remainders f, g, h, &c.

2. Subtract each of the remainders from x, and divide the differences by a, b, c, &c. and there will arise

$$\frac{x-f}{a}$$
, $\frac{x-g}{b}$, $\frac{x-h}{c}$, &c. = whole numbers.

3. Put the first of these fractions $\frac{x-f}{a} = p$, and substi-

tute the value of x, as found in terms of p, from this equation, in the place of x in the second fraction.

4. Find the least value of p in this second fraction, by the last problem, which put =r, and substitute the value of x, as found in terms of r, in the place of x in the third fraction.

Find, in like manner, the least value of r, in this third fraction, which put =s, and substitute the value of x, as found in terms of s, in the fourth fraction as before.

Proceed in the same way with the next following fraction, and so on, to the last; when the value of x, thus determined, will give the whole number required.

EXAMPLES.

1. It is required to find the least whole number, which, being divided by 17, shall leave a remainder of 7, and when divided by 26, shall leave a remainder of 13.

Let
$$x =$$
 the number required.
Then $\frac{x-7}{17}$ and $\frac{x-13}{26}$ = whole numbers.

And, putting $\frac{x-7}{17} = p$, we shall have x = 17p + 7.

Which value of x, being substituted in the second fraction, gives $\frac{17p+7-13}{26} = \frac{17p-6}{26} = wh.$

But it is obvious that $\frac{26p}{96}$ is also = wh.

And consequently $\frac{26p}{26} - \frac{17p - 6}{26} = \frac{9p + 6}{26} = wh$.

$$\operatorname{Or} \frac{9p+6}{26} \times 3 = \frac{27p+18}{26} = p + \frac{p+18}{26} = wh.$$

Where, by rejecting p, there remains $\frac{p+18}{26} = wh$. = r.

Therefore p=26r-18;

Whence, if r be taken = 1, we shall have p=8.

And consequently $x=17p+7=17\times 3+7=143$, the number sought.

2. It is required to find the least whole number, which, being divided by 11, 19, and 29, shall leave the remainders 3, 5, and 10, respectively.

Let x= the number required.

Then $\frac{x-3}{11}$, $\frac{x-5}{19}$ and $\frac{x-10}{29}$ = whole numbers.

And, putting $\frac{x-3}{11} = p$, we shall have x = 11p+3.

Which value of x, being substituted in the second fraction, gives $\frac{11p-2}{19}$ wh.

Or
$$\frac{11p-2}{19} \times 2 = \frac{22p-4}{19} = p + \frac{3p-4}{19} = wh$$
.

And, by rejecting p, there will remain $\frac{3p-4}{19} = \pi h$.

Also by multiplication $\frac{3p-4}{19} \times 6 = \frac{18p-24}{19} = \frac{18p-5}{19} = \frac{18p-5}{19}$

=wh. ;

Or, by rejecting the 1,
$$\frac{18p-5}{19}$$
 wh.

But $\frac{19p}{19}$ is likewise =wh.

Whence $\frac{19p}{19} - \frac{18p-5}{19} - \frac{p+5}{19} - \infty h$, which put =r.

Then we shall have p=19r-5, and x=11(19r-5)+3=209r-52.

And if this value be substituted for x in the third fraction, there will arise

 $\frac{209r-62}{29} = 7r-2 + \frac{6r-4}{29} = wh.;$

Or, by neglecting 7r-2, we shall have the remaining part of the expression $\frac{6r-4}{29}=wh$.;

But by multiplication, $\frac{6r-4}{29} \times 5 = \frac{30r-20}{29} = r + \frac{r-20}{29} = wh.;$

Or, by rejecting r, there will remain $\frac{r-20}{29} = wh$. which

put =s.

Then r=29s+20; where, by taking s=0, we shall have r=20.

And consequently $x = 209r - 52 = 209 \times 20 - 52 = 4128$,

the number required.
3. To find a number, which, being divided by 6, shall leave the remainder 2, and when divided by 13, shall leave

the remainder 3.

Ans. 68

4. It is required to find a number, which being divided by 7, shall leave 5 for a remainder, and if divided by 9,

the remainder shall be 2. Ans. 110
5. It is required to find the least whole number, which,

being divided by 39, shall leave the remainder 16, and when divided by 56, the remainder shall be 27.

Ans. 1147

6. It is required to find the least whole number, which,

being divided by 7, 8, and 9, respectively, shall leave the remainders 5, 7, and 8. Ans. 1727

7. It is required to find the least whole number, which, being divided by each of the nine digits, 1, 2, 3, 4, 5, 6, Ans. 2520

7, 8, 9, shall leave no remainders.

8. A person receiving a box of oranges, observed, that, when he told them out by 2, 3, 4, 5, and 6 at a time, he had none remaining; but when he told them out by 7 at a time, there remained 5; how many oranges were there Ans. 180 in the box?

OF THE

DIOPHANTINE ANALYSIS.

THIS branch of Algebra, which is so called from its inventor, Diophantus, a Greek mathematician of Alexandria in Egypt, who flourished in or about the third century after Christ, relates chiefly to the finding of square and cube numbers, or to the rendering certain compound expressions free from surds; the method of doing which is by making such substitutions for the unknown quantity, as will reduce the resulting equation to a simple one, and then finding the value of that quantity in terms of the rest.

It is to be observed, however, that questions of this kind do not always admit of answers in rational numbers, and that, when they are resolvable in this way, no rule can be given that will apply in all the cases that may occur; but as far as respects a particular class of these problems, relating to squares, they may generally be determined by means of some of the rules derived from the following

formulæ.

PROBLEM I.

To find such values of x as will make $\sqrt{(ax^2+bx+c)}$ rational, or $ax^2 + bx + c = a$ square.

RULE.

1. When the first term of the formula is wanting, or a = 0, put he side of the square sought = n; then $bx + c = n^2$.

And, consequently, by transposing c, and dividing by the coefficient b, we shall have $x = \frac{n^2 - c}{b}$; where n may be any number taken at pleasure.

2. When the last term is wanting, or c=0, put the side of the square sought=nx, or, for the sake of greater generality,= $\frac{mx}{n}$; then, in this case, we shall have $ax^2+bx=m^2x^2$

 $\frac{1}{n^2}$.

And, consequently, by multiplying by n^2 , and dividing by x, there will arise $an^2x+bn^2=m^2x$, and $x=\frac{bn^2}{m^2-an^2}$, where m and n, both in this and the following cases, may be any whole numbers whatever, that will give positive answers.

3. When the coefficient a, of the first term, is a square number, put it $=d^2$, and assume the side of the square sought $=dx+\frac{m}{n}$; then, $d^2x^2+bx+c=d^2x^2+\frac{2dm}{n}x+\frac{m^2}{n^2}$.

And, consequently, by cancelling d^2x^2 , and multiplying by n^2 , we shall have $bn^2x+cn^2=2dmnx+m^2$, and $x=m^2-cn^2$

 bn^2-2dmn

4. When the last term c is a square number, put it = e^2 , and assume the side of the square sought = $\frac{mx}{n} + e$; then, $ax^2 + bx + e^2 = \frac{m^2x^2}{n^2} + \frac{2em}{n}x + e^2$. And, consequently, by cancelling e^2 , and dividing by x, we shall have $ax + b = \frac{m^2x}{n^2} + \frac{2em}{n}$, and $x = \frac{bn^2 - 2emn}{m^2 - an^2}$.

5. When the given formula, or general expression, ax^2+bx+c

can be divided into two factors of the form fx+g and hx+k, which it always can when b^2-4ac is a square, let there

be taken $(fx+g)\times(hx+k)=\frac{m^2}{n^2}(fx+g)^2$; then, by re-

duction, we shall have $x = \frac{gm^2 - kn^2}{hn^2 - fm^2}$; where it may be observed that if the square root of $h^2 - 4mc$ when rational

observed, that if the square root of $b^2 - 4ac$, when rational, be put $= \delta$, the two factors above mentioned, will be

$$ax + \frac{b-\delta}{2}$$
, and $x + \frac{b+\delta}{2a}$

And, consequently by substituting them in the place of the former, we shall have

 $x = \frac{am^2(b-b) - n^2(b+b)}{2a(n^2 - am^2)}.$ 6. When the formula, last mentioned, can be separated

into two parts, one of which is a square, and the other the product of two factors, its solution may be obtained by putting the sum of the square and the product so formed,

equal to the square of the sum of its roots, and $\frac{m}{n}$ times one of the factors, and then finding the values of x as in the former instances.

7. These being all the cases of the general formula that are resolvable by any direct rule, it only remains to observe, that, either in these, or other instances of a different kind, if we can find, by trials, any one simple value of the unknown quantity which satisfies the condition of the question, an expression may be derived from this that will furnish as many other values of it as we please.

Thus, let p, in the given formula $ax^2 + bx + c$, be a value

of x so found, and make $ap^2 + bp + c = q^2$.

Then, by putting x=y+p, we shall have $ax^2+bx+c=a(y+p)^2+b(y+p)+c=ay^2+(2ap+b)y+ap^2+bp+c$, or $ax^2+bx+c=ay^2+(2ap+b)y+q^2$.

From which latter expression the values of y, and con-

sequently those of x may be found, as in Case 4.

Or, because $c = q^2 - bp - ap^2$, if this value be substituted for c in the original formula $ax^2 + bx + c$, it will become $a(x^2 - p^2) + b(x - p) + q^2$, or

 $g^2 + (x-p) \times (ax+ap+b) = a \text{ square };$ which last expression can be resolved by Case 6.

It may here, also, be farther observed, that by putting the given formula $ax^2 + bx + c = \frac{y^2}{4}$, and taking $x = \frac{z - b}{2a}$;

we shall have, by substituting this value for x in the former of these expressions, and then multiplying by 4a, and transposing the terms $ay^2 + (b-4ac) = z^2$; or putting, for the sake of greater simplicity, $b^2 - 4ac = b'$, this last expression may then be exhibited under the form $ay^2 + b' = z^2$, where it is obvious, that if $ay^2 + (b^2 - 4ac)$, or its equal $ay^2 + b'$, can be made a square, $ax^2 + bx + c$, will also be a square.

And as the proposed formula can always be reduced to one of this kind, which consists only of two terms, the possibility or impossibility of resolving the question, in this

state of it, can be more easily perceived.

EXAMPLES.

1. It is required to find a number, such that if it be multiplied by 5, and then added to 19, the result shall be a square.

Let x = the required number; then, as in Case 1, 5x + $19 = n^2$, or $x = \frac{n^2 - 19}{5}$; where it is evident that n may be

any number whatever greater than 19.

Whence, if n be taken =5, 6, 7, respectively, we shall have $x = \frac{25-19}{5} = 1\frac{1}{5}$, or $\frac{36-19}{5} = 3\frac{2}{5}$, or $\frac{49-19}{5} = 6$; the

latter of which is the least value of x, in whole numbers, that will answer the conditions of the question; and consequently $5x + 19 = 5 \times 6 + 19 = 30 + 19 = 49$, a square number as was required.

2. It is required to find an integral number, such that it

shall be both a triangular number and a square.

It is here to be observed, that all triangular numbers are of the form $\frac{x^2+x}{2}$; and therefore the question is reduced

to the making $\frac{x^2+x}{2}$, or its equal $\frac{2x^2+2x}{4}$ a square.

Where, since the divisor 4 is a square number, it is the same as if it were required to make $2x^2+2x$ a square.

Let therefore $2x^2 + 2x = \left(\frac{mx}{n}\right)^2 = \frac{m^2x^2}{n^2}$, agreeably to the method laid down in Case 2.

Then, by dividing by x, and multiplying the result by n^2 , the equation will become $2n^2x+2n^2=m^2x$, or $(m^2-2n^2)x=2n^2$; and consequently $x=\frac{2n^2}{m^2-2n^2}$; where, if n be taken =2, and m=3, we shall have x=8, and $\frac{x^2+x}{2}=\frac{64+8}{2}=\frac{72}{2}=36$, which is the least integral triangular

number that is at the same time a square.

3. It is required to find the least integral number, such that if 4 times its square be added to 29, the result shall

be a square. Here it is evident, that this is the same as to make $4x^2 + 29$ a square.

And, as the first term in the expression is a square, let $4x^2+29=\left(2x+\frac{m}{n}\right)^2=4x^2+\frac{4m}{n}x+\frac{m^2}{n^2}$; agreeably to Case 3.

Then, $\frac{4m}{n}x + \frac{m^2}{n^2} = 29$, or $\frac{4m}{n}x = 29 - \frac{m^2}{n^2}$; and consequently $x = \frac{29n^2 - in^2}{4mn}$; where, if m and n be each taken =1, we shall have $x = \frac{29-1}{4} = 7$, and $4x^2 + 29 = 4 \times 49 + 29 = 225 = (15)^2$, which is a square number, as was re-

quired.

4. It is required to find such a value of x as will make $7x^2 - 5x + 1$ a square.

Here the last term 1 being a square, let these be taken, according to Case 4.

$$7x^{2} - 5x + 1 = \left(\frac{m}{n}x - 1\right)^{2} = \frac{m^{2}}{n^{2}}x^{2} - \frac{2m}{n}x + 1.$$

Then, by rejecting the 1 on each side of the equation, and dividing by x, we shall have $7x-5=\frac{m^2}{n^2}x-\frac{2m}{n}$, and

consequently $x = \frac{2mn - 5n^2}{m^2 - 7n^2}$; where, if m and n be each

taken = 1, the result will give $x = \frac{2-5}{1-7} \cdot \frac{3}{6} = \frac{1}{2}$, or by tak-

ing n=3, and m=8, we shall have $x=\frac{48-45}{64-63}=3$, which makes $7\times3^2-5\times3+1=49=7^2$, as required.

5. It is required to find such a value of x as will make $8x^2 + 14x + 6$ a square.

Here, by comparing this expression with the general formula $ax^2 + bx + c$, we shall have a=8, b=14, and c=6.

And as neither a nor c, in the present instance, are squares, but $b^2 - 4ac = 196 - 192 = 4$ is a square, the given expression can be resolved, by Case 5, into the two following factors 8x + 6, and x + 1.

Let, therefore, $8x^2+14x+6=(8x+6)(x+1)=\frac{m^2}{n^2}(x+1)^2$, agreeably to the rule there laid down.

Then there will arise, by dividing each side by x+1, $3x+6=\frac{nx^2}{x^2}(x+1)$.

And, consequently, by multiplication and reduction, we shall have, in this case, $x = \frac{m^3 - 6n^2}{8n^2 - m^2}$; where it appears,

that, in order to obtain a rational answer, $\frac{m^2}{n^2}$ must be less than 8, and greater than 6.

Whence, by taking m=5, and n=2, we shall have $x=\frac{25-24}{32-25}=\frac{1}{7}$, which makes $\frac{8}{49}+\frac{14}{7}+6=\frac{400}{49}=\left(\frac{20}{7}\right)^2$, as required.

-6. It is required to find such a value of x as will make $2x^2 - 2$ a square.

Here, by comparing this with the general formula $ax^2 + bx + c$, as before, we shall have a = 2, b = c, and c = -2.

And as neither u nor c are squares, but $b^2 - 4ac = -4ac$ = $-4(2\times -2)=16$ is a square, the root of which is 4, the given expression can be resolved, by Case 5, into the two factors 2x-2, and x+1, or 2(x-1), and (x+1), which is evident indeed, in this case, from inspection.

Let, therefore, $2x^2-2=2(x-1)\times(x+1)=\frac{m^2}{n^2}(x+1)^2$, agreeably to the rule; and there will arise by division $2x-2=\frac{m^2}{n^2}(x+1)$. And, consequently, by multiplication,

and reducing the result, we shall have $x = \frac{2n^2 + m^2}{2n^2 - m^2}$; where, by taking n=1, and m=1, we shall have x=3, and $2x^2 - 2=18-2=16=(4)^2$, or taking n=2, and m=3, the result will give x=-17.

But as x enters the problem only in its second, +17 may be taken instead of -17; since either of them give $2x^2-2=576=(24)^2$.

7. It is required to find such a value of x as will make

 $5x^2 + 36x + 7$ a square.

Here, by comparing the expression with the general formula, we shall have a=5, b=36, and c=7.

And as neither a nor c are squares, but $b^2 - 4ac = 1296 - 140 = 1156 = (34)^2$, is a square, it can be resolved, as in the last example, into the two factors 5x + 1, and x + 7.

Whence, putting $5x^2 + 36x + 7 = (\bar{\epsilon}x + 1) \times (x + 7) = \frac{m^2}{n^4}$ $(x+7)^2$, there will arise, by dividing by x+7, $5x+1 = \frac{n^2}{n^2}(x+7)$.

And, consequently, by multiplication, and reducing the resulting expression, we shall have $x = \frac{7m^2 - n^2}{5n^2 - m^2}$; where,

taking m=2, and n=1, the substitution will give x= $\frac{7 \times 4 - 1}{5 \times 1 - 4} = 27$, which makes $5 \times (27)^2 + 36 \times 27 + 7 = 4624$ = (68)2, as required.

8. It is required to find such a value of x as will make

 $6x^2 + 13x + 10$ a square.

Here, by comparing the given expression with the general formula $ax^2 + bx + c$, we have a=6, b=12, and c=10. And as neither a, c, nor b? - 4ac, are squares, the question, if possible, can only be resolved by the method pointed out in Case 6.

In order, therefore, to try it in this way, let the first simple square 4, be subtracted from it, and there will re-

main, in that case, $6x^2 + 13x + 6$.

Then, since $(13)^2 - 4(6 \times 6) = 169 - 144 = 25$, is now a square, this part of the formula can be resolved by Case 5, into the two factors :

3x+2, and 2x+3.

Whence, by assuming, according to the rule, $6x^2 + 13x$ $+10=4+(3x+2)\times(2x+3)=\left\{2+\frac{m}{n}(3x+2)\right\}^2=4+$ $\frac{4m}{m}(3x+2)+\frac{m^2}{m^2}(3x+2)^2$, we shall have, by cancelling the 4, on each side, and dividing by 3x+2; $2x+3=\frac{4m}{m}$

$$+\frac{m^2}{n^2}(3x+2).$$

And, consequently, by multiplying by n2, and transposing the terms, we have $2n^2x-3m^2x=4mn+2m^2-3n^2$, or $4mn + 2m^2 - 3n^2$

 $2n^2 - 3m^3$

Where putting m=2, and n=3, the result will give x=24+8-27 5 $\frac{18-12}{6} = \frac{1}{6}$, or if m be taken = 13, and n=17, we shall

have
$$x = \frac{4 \times 17 \times 13 + 2 \times (13)^2 - 3 \times (17)^2}{9(17)^2 - 3(13)^2} = \frac{355}{71} = 5.$$

Which makes $6 \times (5)^2 + 13 \times 5 + 10 = 225 = (15)^2$, as required.

9. It is required to find such a value of x as will make

 $13x^2 + 15x + 7$ a square.

Here, by comparing this with the general formula, as before, we have a=13, b=15, and c=7. And as neither $a, b, \text{ nor } b^2 - 4ac, \text{ are squares, the answer to the question}$ if it be resolvable, can only be obtained by Case 6. In order, therefore, to try it, in that way, let $(1-x)^2$ or 1-x $2x+x^2$ be subtracted, from the given expression, and there will remain $12x^2 + 17x + 6$.

And as $(17)^2 - 4(6 \times 12)$, which is = 1, is now a square, this part of the formula can be resolved by Case 5, into the two factors 4x+3 and 3x+2. Whence, assuming 13

The two factors
$$4x + 3$$
 and $3x + 2$. Whence, assuming to $x^2 + 15x + 7 = (1-x)^2 + (4x+3) \times (3x+2) = \left\{ (1-x) + \frac{m}{n}(3x+2) \right\}^2 = (1-x)^2 + \frac{2m}{n}(1-x) \times (3x+2) + \frac{m^2}{n^2}(3x+2)^2$, we shall have, by cancelling $(1-x)^2$, and dividing by $5x+2$; $4x+3 = \frac{2m}{n}(1-x) + \frac{m^2}{n^2}(3x+2)$; and consequently, by multiplying by n^2 , and transposing the terms, there will arise $4n^2x + 2mnx - 3m^2x = 2mn + 2m^2 - 3n^2$, or $x = 2mn + 2m^2 - 3n^2$

 $4n^2 + 2mn - 3m^2$

Where putting m and n each =1, we shall have x= $\frac{2+2-3}{4+2-3} = \frac{1}{3}$, which makes $\frac{13}{9} + \frac{15}{3} + 7 = \frac{13}{9} + \frac{45}{9} + \frac{63}{9} = \frac{121}{9}$

 $=\left(\frac{11}{3}\right)^2$, as required.

10. It is required to find such a value of x as will make $7x^2 + 2$ a square.

Here it is easy to perceive that neither of the former

rules will apply.

But as the expression evidently becomes a square wher x=1, let, therefore, x=1+y, according to case 7, and we shall have

 $7x^2 + 2 = 9 + 14y + 7y^2$,

Or, putting $9+14y+7y^2=\left(3+\frac{m}{n}y\right)^2$, according to the rule, and squaring the right hand side, $9+14y+7y^2=9+\frac{6m}{n}y+\frac{m^2}{n^2}y^2$.

Hence rejecting the 9's and dividing the remaining terms by y, we have $7n^2y + 14n^2 = 6mn + m^2y$; and, consequently, $y = \frac{6mn - 14n^2}{7n^2 - m^2}$, and $x = 1 + \frac{6mn - 14n^2}{7n^2 - m^2}$; where it is evident that m and n may be any positive or negative numbers whatever.

If, for instance, m and n be each taken = 1, we shall have $y = -\frac{4}{3}$ and $x = -\frac{1}{3}$. Or, since the second power of x only enters the formula, we may take, as in a former instance, $x = \frac{1}{3}$, which value makes $7x^2 + 2 = \frac{7}{3} + 2 = \frac{7}{3} + \frac{1}{3} = \frac{7}{3} + \frac$

Or, if n=3 and n=-1, we shall have x=17, and $7x^2 + 2 = 7 \times (17)^2 + 2 = 2025 = (45)^2$, a square as before.

And by proceeding in this manner, we may obtain as many other values of x as we please.

PROBLEM II.

To find such values of x as will make $\sqrt{(ax^3+bx^2+cx+d)}$ rational, or $ax^3+bx^2+cx+d=a$ square. This problem is much more limited and difficult to be resolved, than the former; as there are but a few cases of it that admits of answers in rational numbers; and in these the rules for obtaining them are of a very confined nature; being mostly such as are subject to certain limitations, or that admit only of a few simple answers, which in the instances here mentioned, may be found as follows.

RULE.

1. When the third and fourth terms of the formula are wanting, or c and d are each =0, put the side of the square sought =nx, then $ax^3+bx^2=n^2x^2$.

And, consequently, by dividing each side of the equation by x^2 , we shall have $ax+b=n^2$, or $x=\frac{n^3-b}{a}$, where n may be any integral or fractional number whatever.

2. When the last term d is a square, put it = e, and assume the side of the required square = $e + \frac{c}{2e}x$; and the

reversed formula is $e^2 + cx + bx^2 + ax^3 = e^2 + cx + \frac{c^2}{4e^2}x^2$.

Whence, by expunging the terms $e^2 + cx$, which are common, and dividing by x^2 , we shall have, $4ac^2x + 4be^2 = c^2$; and, consequently, $x = \frac{c^2 - 4be^2}{4ac^2}$.

Or, if, in the same case, there be put $e + \frac{c}{2e}x + \frac{4be^2 - c^2}{8e^3}x^2$ for the side of the required square, we shall have, by square.

ing, $e^2 + cx + bx^2 + ax^3 = e^2 + cx + bx^2 + \frac{c(4be^2 - c^2)}{8e^4}x^3 + \frac{c(4be^2 - c^2)}{8e^4}x$

 $\frac{(4be^2-c^2)^2}{64e^6}x^4$. And as the first three terms $(e^2+cx+bx^2)$

are now common, there will arise, by expunging them, and then multiplying by $64e^{a}$, $64ae^{6}x^{3} = 8ce^{2}(4be^{2} - c^{2})x^{3} + (4be^{2} - c^{2})^{2}x^{4}$.

Whence, by dividing each side of this last equation by x^3 , and reducing the result, we shall have

$$x = \frac{64ae^6 - 8ce^3(4be^2 - c^3)}{(4be^2 - c^2)^2},$$

which last method gives a new value of x, different from that before obtained.

3. When neither of the above rules can be applied to the question, the formula can be resolved, by first finding, by trial, as in the former problem, some value of the unknown quantity that makes the given expression a square; in which case, other values of it may be determined from this, when they are possible, as follows:

Thus, let p be a value of x so found, and make $ap^3+bp^2+cp+d=q^2$;

Then, by putting x=y+p, we shall have ap^3+bp^2+cp $+d=a(y+p)^3+b(y+p)^2+c(y+p)+d=ay^2+(3ap+b)y^2+(3p^2+2p+c)y+up^3+bp^2+ep+d, \text{ or } ax^3+bx^2+cx+$ $d = ay^{2} + (3ap + b)y^{2} + (3p^{2} + 2p + c)y + q^{2}.$

From which latter form, the value of y, and consequently that of x, may be found by either of the methods given

in Case 2.

It may also be further remarked, that, if the given formula, in any case of this kind, can be resolved into factors, such that one of them shall be a square, it will be sufficient to make the remaining factor a square, in order to render the whole expression so; since a square, multiplied or divided by a square, is still a square.

EXAMPLES.

1. It is required to find such a value of x as will make $11x^3 + 3x^2$ a square.

Let the given expression $11x^3 + 3x^2 = n^2x^2$: agreeably

to Case 1.

Then, by dividing by x^2 , we shall have $11x+3=n^2$; and, consequently, $x = \frac{n^2 - 3}{11}$; where n may be any num-

Taking, therefore, n=2, 3, 4, 5, &c. respectively, we shall have, in this case, $x=\frac{1}{11}, \frac{6}{11}, \frac{13}{11}$, or 2, the last of which is the least integral answer that the question admits

2. It is required to find such values of x as will make

 $x^3 - 2x^2 + 2x + 1$ a square.

Here, the last term 1, being a square, let $1+2x-2x^2+$ $x^3 = (1+x)^2 = 1+2x+x^2$, agreeably to the first part of Case 2.

Then, since the first two terms, on both sides of the equation, destroy each other, we shall have $x^3 - 2x^2 = x^2$, or $x^3 = 3x^2$, and consequently x = 3; which, by substitution, makes $1+2x-2x^2+x^3=1+6-18+27=16$, a square as required.

Again, by putting x=y+3, according to Case 3, we shall have $1+2x-2x^2+x^3=1+2(y+3)-2(y+3)^2+(y+3)^3$

 $=16+17y+7y^2+y^3$.

And, consequently, by making $16+17y+7y^2+y^3=(4+\frac{17}{8}y)^2=16+17y+\frac{289}{64}y^2$, agreeably to the first part of Case 2, by cancelling 16+17y, there will arise $7y^2+y^3=\frac{289}{64}y^2$, or $y+7=\frac{289}{64}$.

Whence $y = \frac{289}{64} - 7 = \frac{289 - 446}{64} = \frac{159}{64}$, and $x = 3 - \frac{159}{64} = \frac{192 - 159}{64} = \frac{33}{64}$, for another value of x.

Which number, being substituted in the original formula, make $1+2x-2x^2+x^3=\frac{429025}{262144}=\left(\frac{555}{162}\right)^2$, a square, as before.

before.

3. It is required to find such values of x as will make $3x^3 - 5x^2 + 6x + 4$ a square.

Here, 4 being a square, let $4+6x-5x^2+3x^3=(2+$

 $(\frac{3}{2}x)^2 = 4 + 6x + \frac{9}{4}x^3$, as in the first part of Case 2.

Then, since the two first terms on each side of the equation destroy each other, we shall have $3x^3 - 5x^2 = \frac{a}{4}$ x^2 , or $3x - 5 = \frac{a}{4}$; and, consequently, in this case $x = \frac{5 + \frac{a}{4}}{3} = \frac{29}{12}$.

Whence $(2+\frac{3}{2}\times\frac{29}{12})^2 = (2+\frac{29}{8})^2(\frac{45}{6})^2$ a square, as was

required.

Or, by the second method of the same Case, let 4+6x $-5x^2+3x^3=(2+\frac{3}{2}x-\frac{29}{16}x^2)^2=4+6x-5x^2-\frac{87}{16}x^3+\frac{841}{256}x^4$; then, as the three first terms on each side of this equation destroy each other, we shall have $\frac{841}{256}x^4-\frac{87}{16}x^3$

=3 \mathfrak{e}^3 , or $\frac{841}{256}x - \frac{87}{16}$ =3, or 841x - 1392 = 768; and con-

sequently, $x = \frac{1392 + 768}{841} = \frac{2160}{841}$, which is another value of x, that, being substituted in the original formula, will make it a square.

4. It is required to find such values of x as will make x3.

+3 a square.

Here, it is evident, that the expression is a square when x=1. Let therefore x=1+y, and we shall have $3+x^3$

 $=4+3y+3y^2+y^3$.

And as the first part of this is a square, make according to the first part of Case 2, $4+3y+3y^2+y^3=(2+\frac{9}{4}y)^2=4+3y+\frac{9}{16}y^2$. Then, because the first two terms on each side of the equation destroy each other, we shall have $y^3+3y^2=\frac{9}{16}y^2$, or $y+3=\frac{9}{16}y$.

Whence
$$y = \frac{9}{16} - 3 = \frac{9 - 48}{16} = \frac{39}{16}$$
, and $x = 1 - \frac{39}{16} = \frac{16 - 39}{16} = \frac{23}{16}$; which is a second value of x .

Again, let
$$4+3y+3y^2+y^3=(2+\frac{3}{4}y+\frac{39}{64}y^2)^2=4+3y+3y^2+\frac{117}{123}y^3+\frac{1521}{4096}y^4$$
, according to the second part of

Case 2.

Then as the first three terms on each side of the equation destroy each other, we shall have $\frac{1521}{4090}y^4 + \frac{117}{128}y^3 = y^2$,

or
$$\frac{1521}{4096}y + \frac{117}{128} = 1$$
.

Whence, also, $y = \frac{352}{1521}$, and $x = 1 + \frac{352}{1521} = \frac{1873}{1521}$, which

is a third value of x.

And by proceeding in the same way with either of these new values of x as with the first, other values of it may be obtained; but the resulting fraction will become continually more complicated in each operation.

PROBLEM III.

 $cx^2 + dx + e$) rational, or $ax^4 + bx^3 + cx^2 + dx + e = a$ square.

The resolution of expressions of this kind, in which the indeterminate, or unknown quantity, rises to the fourth power, is the utmost limit of the researches that have hitherto been made on formulæ affected by the sign of the square root; and in this Problem, as well as in that last given, there are only a few particular cases that admit of answers in rational numbers; the rest being either impossible, or such as afford one or two simple solutions; which may generally be found as follows:

BULE.

1. When the last term e, of the given formula, is a square, put it = f^2 , and make $f^2 + dx + cx^2 + bx^3 + ax^4 = (f + \frac{d}{2f}x + \frac{4cf^2 - d^2}{8f^3}x^2)^2 = f^2 + dx + cx^2 + \frac{d(4cf^2 - d^2)}{8f^4}x^3 + \frac{d(4cf^2 - d^2)}{8f^4}x^4 + \frac{d(4cf^2$

Then, by expunging the first three terms, which are common to each side of the equation, there will remain $bx^3 + ax^4 = \frac{d(4cf^2 - d^2)}{8f^4}x^3 + \frac{(4cf^2 - d^2)^2}{64f^6}x^4$. And consequently, by dividing by x^3 , and reducing the result, we shall

have $x = \frac{64bf^8 - 8df^2(4cf^2 - d^2)}{(4cf^2 - d^2)^2 - 64af^6}$. 2. When the coefficient, a, of the first term of the formula, is a square, put it $=g^2$, and make $g^2x^4 + bx^3 + cx^2 + dx + e = (gx^2 + \frac{b}{2g}x + \frac{4cg^2 - b^2}{8g^3})^2 = g^4x^4 + bx^3 + cx^2 + \frac{4cg^2 - b^2}{8g^3}$ $\frac{b(4cg^2-b^2)^2}{8g^4}x + \frac{(4cg^2-b^2)^2}{6^4g^6}.$ Then, $dx + e = \frac{b(4cg^2-b^2)}{8g^4}x + \frac{(4cg^2-b^2)^2}{6^4g^6}$; and con-

sequently, $x = \frac{(4cg^2 - b^2)^2 - 64cg^6}{64dg^6 - 8bg^2(4eg^2 - b^2)}$; which form likewise fails under similar circumstances with the former.

3. When the first and last terms of the formula are both squares, put $a=g^2$, and $e=f^2$, and make $f^2+dx+cx^2+bx^3+g^2x^4=(f+\frac{d}{2f}x+gx^2)^2=f^2+dx+(2fg+\frac{d^2}{4f^2})x^2+\frac{dg}{f}x^3+g^2x^4$. Then $cx^2+bx^3=(2fg+\frac{d^2}{4f^2})x^2+\frac{dg}{f}x$. And, consequently, $x=\frac{f^2(2fg-c)+\frac{1}{4}d^2}{f(bf-dg)}$.

Or, because g enters the given formula only in its second power, it may be taken either negatively or positively; and, consequently, we shall have $x = \frac{1}{4} \frac{d^2 - f^2(2fg + c)}{f(bf + dg)}$. So that this mode of solution furnishes two different answers.

Also, if there be taken for another supposition $f^2+dx+cx^2+bx^3+g^2x^4=(f+\frac{b}{2g}x+gx^2)^2=f^2+\frac{b}{g}fx+(2fg+\frac{b^2}{4g^2})x^2+bx^3+g^2x^4$, hence by cancelling, $dx+cx^2=\frac{bf}{g}x+(2fg+\frac{b^2}{4g^2})x^2$; and consequently, $x=\frac{g(bg-bf)}{\frac{1}{4}b^2+g^2(2fg-c)}$.

And because f enters the given formula only in the second power, it may be taken either negatively or positively; and, consequently, we shall also have $x = \frac{g(bg + bf)}{\frac{1}{4}b^2 - g^2(2fg + c)}$.

So that this solution likewise furnishes two values of x, which are each different from the former.

But these forms all fail under similar circumstances with

those of the second Case.

4. When neither the first nor the last terms are squares, the formula cannot be resolved in any other way, than by first endeavouring to discover by trials, some simple value of the unknown quantity, that will answer the conditions of the question; and then finding other values of it, according to the methods pointed out in the two last problems.

Thus, let p be a value of x so found, and make ap^4 +

 $bp^3 + cp^2 + dp + e = q^2$.

Then by putting $\dot{x}=y+p$, we shall have $ap^4+bp^3+cp^2+dp+e=a(y+p)^4+b(y+p)^3+c(y+p)^2+d(y+p)+e=ay^4+(ap+b)y^3+(6ap^2+3bp+c)y^2+(4ap^3+3bp^2+2cp+d)y+ap^4+bp^3+cp^2+dp+e, or <math>ax^4+bx^3+cx^2+dx+e=ay^4+(ap+b)y^3+(6ap^2+3bp+c)y^2+(4ap^3+3bp^2+2cp+d)y+q^2$. From which last formula the value of y, and consequently that of x, may be found by Case 1.

EXAMPLES.

1. It is required to find such a value of x, as will make $1-2x+3x^2-4x^3+5x^4$ a square.

Here, the first term 1, being a square, let $1-2x+3x^2-4x^3+5x^4=(1-x+x^2)^2=1-2x+3x^2-2x^3+x^4$, agreeably to the method in Case 1.

Then we shall have $5x^4 - 4x^3 = x^4 - 2x^3$.

And, consequently, 5x - 4 = x - 2; whence $x = \frac{2}{4} = \frac{1}{2}$. And consequently, $1 - 2x + 3x^2 - 4x^3 + 5x^4 = 1 - 1 + \frac{3}{4} - \frac{1}{2}$. $+ \frac{5}{16} = \frac{9}{16}$; which is a square number, as was required.

2. It is required to find such a value of x, as will make

 $4x^4 - 2x^3 - x^2 + 3x - 2$ a square.

Here the first term being a square, let $4x^4 - 2x^3 - x^2 + 3x - 2 = (2x^2 - \frac{1}{2}x - \frac{5}{16})^2 = 4x^4 - 2x^3 - x^2 + \frac{5}{16}x + \frac{25}{256}$, according to the method in Case 2.

Then we shall have $3x-2=\frac{5}{16}x+\frac{25}{256}$, or $3x-\frac{5}{6}$ 16

 $+\frac{25}{256}$. Whence, 768x-80x=512+25; and, consequently, $x=\frac{512+25}{768-80}=\frac{537}{688}$.

Or, if we put $x = \frac{1}{y}$, the formula in that case will become

$$\frac{4}{y^4} - \frac{2}{y^3} - \frac{1}{y^2} + \frac{3}{y} - 2.$$

And, therefore, multiplying this by y^4 , which is a square, it will be $4-2y-y^2+3y^3-2y^4$. Where the first term being now a square, if the expression, so transformed, be resolved by Case 1, we shall have $y=\frac{688}{537}$; and $x=\frac{1}{y}$

 $\frac{537}{688}$, as before.

3. It is required to find such values of x, as will make $1+3x+7x^2-2x^3+4x^4$ a square.

Here, both the first and last terms being squares, let $1+3x+7x^2-2x^3+4x^4=(1+\frac{3}{2}x+2x^2)^2=1+3x+\frac{25}{4}x^2+6$ x^3+4x^4 , according to the method in Case 3.

Then, we shall have $6x^3 + \frac{25}{4}x^2 = 7x^2 - 2x^3$; or 6x + 2

$$x=7-\frac{25}{4}$$
; and, by reduction, $x=\frac{3}{32}$.

And, if we put the same formula $1+3x+7x^2-2x^3+4x^4=(1+\frac{2}{3}x-2x^2)^2$, $=1+3x-\frac{7}{4}x^2-6x^3+4x^4$, we shall have, by cancelling, $7x^2-2x^2=-\frac{7}{4}x^2-6x^3$; whence 6x

$$-2x = -\frac{7}{4} - 7 = -\frac{35}{4}$$
; or $x = -\frac{35}{16}$.

And, in a similar manner, other values of x may be found, by employing the method of substitution pointed out in the latter part of Case 3.

4. It is required to find such values of x as will make

 $2x^4 - 1$ a square.

Here, 1 being an obvious value of x, let, according to

Case 4, x = 1 + y.

Then $2x^4 - 1 = 2(1+y)^4 - 1 = 2(1+4y+6y^2+4^3+y^4) - 1 = 1+8y+12y^2+8y^3+2y^4$. And since the first term of this last expression is now a square, we shall have, by Case 1, $1+8y+12y^2+8y^3+2y^4=(1+4y-2y^2)^2=1+8y+12y^2-16y^3+4y^4$.

Whence, as the three first terms of the two numbers of this equation destroy each other, there will remain $4y^4 - 16y^3 = 2y^4 + 8y^3$; or y = 12; and, consequently, x = 1 + y = 13; which value being substituted for x, makes $2x^4 - 1$

=57121=(239)², as required. And if 13 be now taken, as the known value of x, and the operation be repeated as before, we shall obtain, for another value of x, the complicated fraction $\frac{10607469769}{2447192159}$.

PROBLEM IV.

To find such values of x as will make $\sqrt[3]{(ax^3+bx^2+cx+d)}$ rational, or $ax^3+bx^2+cx+d=a$ cube. This formula, like the two latter of those relating to squares, cannot be resolved by any direct method, except in the cases where the first or last terms of the expression are cubes; it being necessary, in all the rest, that some simple number answering the conditions of the question, should be first found by trial, before we can hope to obtain others; but when this can be done, the problem, in each of the cases here mentioned, may be resolved as follows.

RULE.

1. When the last term d of the given formula is a cube, put it = e^3 , and make $e^3 + cx + bx^2 + ax^3 = (e + \frac{c}{3e^2}x)^3 =$

$$e^3 + cx + \frac{c^2}{3e^3}x^2 + \frac{c^3}{27e^6}x^3$$
.

Then, by expunging the two first terms on each side of the equation, which are common, there will remain $ax^2 + bx^2 = \frac{c^3}{27e^6}x^3 + \frac{c^2}{3e^3}x^2$; whence, by division and reduction, we shall have $27ae^6x + 27be^6 = c^3x + 9c^2e^3$, and consequently $x = \frac{9e^3(3be^3 - c^2)}{c^3 - 27ae^6}$; which form fails when the coefficients b and c, or a and c, are each equal 0.

2. When the coefficient a of the first term is a cube, put it = f^3 , and make $f^3x^3 + bx^2 + cx + d = \left(fx + \frac{b}{3f^3}\right)^3 = f^3x^3 + bx^2 + \frac{b^3}{3f^3}x + \frac{b^3}{27f^5}$.

Then, by expunging the two first terms on each side of the equation, as before, there will remain $cx+d=\frac{b^2}{3f^2}x+\frac{b^3}{27f^6}$; whence, by multiplying by $27f^6$, we shall have 27

 $f^{*}cx+27df^{6}=9b^{2}f^{3}x+b^{3}$, and consequently $x=\frac{b^{3}-27df^{6}}{2f^{3}(3cf^{3}-b^{2})}$; which form likewise fails, when b and c, or b and d, are each =0.

3. When the first and last terms are both cubes, put $a = f^3$ and $d = e^3$, and make $e^3 + cx + bx^2 + f^3x^3 = (e + fx)^3 =$

 $e^3 + 3fe^2x + 3f^2ex^2 + f^3x^3$.

Then, $cx+bx^2=3fe^2x+3f^2ex^2$:

Whence, we shall have $bx-3f^2ex=3fe^2-c$; and, consequently $x=\frac{3fe^2-c}{b-3f^2e}$; which formula may be resolved by either of the two first cases,

4. When neither the first nor the last terms are cubes, let p be a value of x, found by inspection, or by trials, and

make $ap^3+bp^2+cp+d=q^3$.

Then, by putting x=y+p, we shall have $ap^3+bp^2+rp+d=a(y+p)^3+b(y+p)^2+c(y+p)+d=ay^3+(3ap+b)$ $y^2+(3ap^2+2bp+c)y+ap^3+bp^2+cp+d$, or $ax^3+bx^2+cx+d=ay^3+(3ap+b)y^2+(3ap^2+2bp+c)y+q^3$.

From which latter form, the value of y, and conse-

quently that of x, may be found, as in Case 1.

EXAMPLES.

1. It is required to find such a value of x as will make x^2+x+1 a cube.

Here, the last term being a cube, let the root of the

cube sought = $1 + \frac{1}{3}x$, according to Case 1.

Then, by cubing, we shall have $1+x+x^2=1+x+\frac{1}{3}x^2+\frac{1}{27}x^3$;

And, since the two first terms on each side of this equation destroy each other, there will remain $x^2 = \frac{1}{3}x^2 + \frac{1}{27}x^3$.

Whence, dividing by x^2 , we shall have $\frac{1}{27}x + \frac{1}{3} = 1$, or x + 9 = 27; and consequently x = 27 - 9 = 18; which num-

ber, by substitution, makes $1+x+x^2=1+18+324=343$ =7° a cube number, as was required.

And if we now take this value of x, and proceed according to the method employed in Case 4, we shall obtain x = -

137826 50653; which last number will also lead in like manner, to other new values.

2. It is required to find such a value of x as will make x^3+3x^2+133 a cube.

Here, the first term being a cube, let its root =1+x, according to Case 2.

Then, by cubing, we shall have $133+3x^2+x^3=(1+r)^3=1+3x+3x^2+x^3$.

And since the two last terms of this equation destroy each other, there will remain 1+3x=13.5, or 3x=133-13.5

1=132; whence $x = \frac{132}{3} = 44$, and $x^3 + 3x^2 + 133 = 91125$

==(45)3, a cube number, as was required.

And if 45 be now taken as a known value of x, other values of it may be found, as in the last example.

3. It is required to find such a value of x, as will make $8+28x+89x^2-125x^3$ a cube.

Here, let the root sought =2-5x according to Case 3. Then, by cubing we shall have $8+28x+89x^2-125x^3$

 $= (2-5x)^3 = 8 - 60x + 150x^2 - 125x^3.$

And, since the first and last terms of this equation destroy each other, there will remain $28x + 89x^2 = -60x + 150x^2$.

Whence, by dividing by x, and transposing the terms we shall have 150x - 89x = 28 + 60, or 64x = 88; and consegretations.

quently $x = \frac{88}{61}$

And as this formula can also be resolved either by the first or second case, other values of x, may be obtained, that will equally answer the conditions of the question.

4. It is required to find such a value of x, as will make

 $2x^3 - 3x + 7 \text{ a cube.}$

Here, -1 being a value of x that is readily found, by inspection, let x=y-1, agreeably to Case 4.

Then, by substitution, we shall have $2x^3-3x+7=2(y-1)^3-3(y-1)+7=2y^3-6y^2+3y+8$.

 $-1)^3 - 3(y-1) + 7 = 2y^3 - 6y^2 + 3y + 8$. And as the last term of this expression is a cube, let $8 + 3y - 6y^2 + 2y^3 = (2 + \frac{1}{4}y)^3 = 8 + 3y + \frac{3}{8}y^2 + \frac{1}{64}y^3$ according to Case 1. Then, by expunging the equal terms on each side, there will remain $2y^3 - 6y^2 = \frac{3}{8}y^2 + \frac{1}{64}y^3$.

Whence, dividing by y^2 , and reducing the terms, we shall have 128y - 384 = y + 24, or 127y = 408; and consequently $y = \frac{408}{127}$, and $x = \frac{408}{127} - 1 = \frac{281}{127}$.

Which number, by substitution, makes $2x^3-3x+7=\frac{2\times(281)^3}{(127)^3}-3\times\frac{281}{127}+7=\frac{45118016}{2048383}=\left(\frac{356}{127}\right)^3$, as required. And, by taking this last as a new value of x, others may be determined by the same method.

PROBLEM V.

Of the resolution of double and triple equalities.

When a single formula, containing one or more unknown quantities, is to be transformed into a perfect power, such as a square or a cube, this is called, in the Diophantine Analysis, a simple equality; and when two formulæ, containing the same unknown quantity, or quantities, are to be each transformed to some perfect power, it is then called a double equality, and so on; the methods of resolving which, in such cases as admit of any direct rule are as follows:

RULE.

1. In the case where the unknown quantity does not exceed the first degree, as in the double equality

 $ax+b=\square$, and $cx+d=\square$,

let the first of these formulæ $ax+b=z^2$, and the second $cx+d=w^2$.

Then, by equating the two values of x, as found from these equations, we shall have $cz^2 + ad - bc = aw^2$, or acz^2

 $+a(ad-bc)=a^2w^2$.

And since the quantity on the right hand of this equation is now a square, it only remains to find such a value of z as will make, when the question is resolvable, $acz^2 + a(ad -bc) = \Box$; which being done, according to the method $z^2 - b$

pointed out in Problem 1, we shall have $x = \frac{z^2 - b}{a}$.

When the unknown quantity does not exceed the second degree, and is found in each of the terms of the two formulæ; as in the double equality

 $ax^2 + bx = \square$, and $cx^2 + dx = \square$.

Let $x = \frac{1}{y}$, then, by substitution, and multiplying each of the resulting expressions by y^2 , we shall have

 $a+by=\Box$, and $c+dy=\Box$,

from which last formula, the value of y, when the question is possible, and consequently that of x, may be determined as in Case 1.

But if it were required to make the two general expres-

sions

 $ax^2+bx+c=\square$, and $dx^2+ex+f=\square$, the solution could only be obtained in a few particular cases, as the resulting equality would rise to the fourth power.

3. In the case of a triple equality, where it is required

to make

 $ax + by = \square$, $cx + dy = \square$, and $ex + fy = \square$,

let the first of them $ax+by=u^2$, the second $cx+dy=v^2$,

and the third $ex+fy=w^2$.

Then, by first eliminating x in each of these equations, and afterwards y, in the two resulting equations, we shall have $(af-be)v^2-(cf-de)u^2=(ad-bc)w^2$ or, putting v=uz, and reducing the terms, the result will

give the simple equality $\frac{af-be}{ad-bc}z^2 - \frac{cf-de}{ad-bc} = \frac{w^2}{u^2}$; where

the right hand member being a square, it only remains to find a value of z that will make the left hand member a square; which, when possible, may be done by Problem 1. Hence, having z, we have as above, v=uz; and the first

two equations will give $x = \frac{d - bz^2}{ad - bc}u^2$, and $y = \frac{az^2 - c}{ad - bc}u^2$, where u may be any whole or fractional number whatever.

But if the three formulæ, here proposed, contained only one variable quantity, the simple equality, to which it would be necessary to reduce them, would rise, as in the last case, to the fourth power; and be equally limited

with respect to its solution.

4. In other cases of this kind, all that can be done is to find successively by the former rules, several answers. when one is known; and, if neither this nor any of the above mentioned modes of solution are, found to succeed, the Problem under consideration can only be determined by adopting some artifice of substitution that will fulfil one or more of the required conditions, and then resolving the remaining formulæ, when they are possible, by the methods already delivered for that purpose; but as no general precepts can be given, for obtaining the solution in this way, the proper mode of proceeding, in such cases, must chiefly depend upon the skill and sagacity of the learner.

EXAMPLES.

1. It is required to find a number x, such that x+128and x+192 shall be both squares.

Here, according to Case 1, let $x+128=w^2$, and x+192

 $=z^2$.

Then, by eliminating x, and equating the result, we shall have $w^2 - 128 = z^2 - 192$, or $w^2 + 64 = z^2$.

And, as the quantity on the right hand side of the equation is now a square, it only remains to make w2 +64 a square.

For which purpose, put its root = w + n; then $w^2 + 64$ $=w^2+2nw+n^2$, or $2nw+n^2=64$; and consequently w= $\frac{1}{2n}$; where taking n, which is arbitrary, =2, we shall have $w = \frac{64-4}{4} = \frac{60}{4} = 15$; and consequently $x = w^2 - 128$ = $15^2 - 128 = 225 - 128 = 97$, the answer.

2. It is required to find a number x, such that $x^2 + x$

and $x^2 - x$ shall be both squares.

Here, according to Case 2, of the last Problem, let $x=\frac{1}{y}$; then we shall have to make $\frac{1}{y^2}+\frac{1}{y}$, and $\frac{1}{y^2}-\frac{1}{y}$ squares;

or, by reduction, $\frac{1}{y^2}(1+y) = \square$, and $\frac{1}{y^2}(1-y) = \square$.

Or, since a square number, when divided by a square number, is still a square, it is the same as to make

 $1+y=\square$ and $1-y=\square$,

for this purpose, therefore, let $1+y=z^2$, or $y=z^2-1$; then $1-y=2-z^2$; which is also to be made a square.

But as neither the first nor last terms of this formula, are squares, we must, in order to succeed, find some simple number, that will answer the condition required; which, it is evident, from inspection, will be the case when z=1.

Let, therefore, z=1-w, agreeably to Problem 1. Case 7, and we shall have $1-y=2-z^2=2-(1-w)^2=1+2w$

 $-w^2$; or $y=w^2-2w$;

or, putting 1-nw for the root of the former of these expressions, there will arise, by squaring, $1+2w-w^2=1-2nw+n^2w^2$.

Whence, expunging the one on each side, and dividing by w, we shall have $2-w=-2n+n^2w$; and consequently

 $w = \frac{2n+2}{n^2+1}$, and $x = \frac{1}{y} = \frac{1}{w^2 - 2w} = \frac{(n^2+1)^2}{4n - 4n^3}$,

where, in order to render the value of x positive, n may be taken equal to any proper fraction whatever.

Or, if for the sake of greater generality, $\frac{m}{n}$ be substi-

tuted for n, we shall then have

$$x = \frac{(m^2 + n^2)^2}{4mn(n^2 - m^2)}$$

where m and n may now be taken equal to any integral numbers whatever, provided n be made greater than m.

If, for instance, n=2 and m=1, we shall have $x=\frac{25}{24}$;

and if n=3 and m=2, $x=\frac{169}{120}$; and so on, for any other number.

3. It is required to find three whole numbers in arithmetical progression, such, that the sum of every two of them should be a square.

Let x, x+y, and x+2y, be the three numbers sought; and put $2x+y=u^2$, $2x+2y=v^2$, and $2x+3y=w^2$, agreeably to Case 3.

Then, by eliminating x and y from each of these equations, we shall have $v^2 - u^2 = w^2 - v^2$, or $2v^2 - u^2 = w^2$.

And, if we now put v=uz, there will arise $2u^2z^2-u^2=w^2$; or by dividing by u^2 , $2z^2-1=\frac{w^2}{u^2}$; where the right

hand member being a square, it only remains to make $2z^2 - 1$ a square, which it evidently is when z=1.

But as this value would be found not to answer the con-

ditions of the question, let z=1-p; then $2z^2-1=2(1-p)^2-1=1-4p+2p^2$. And, consequently, if this last expression be put $=(1-p)^2-1=1$

And, consequently, if this last expression be put = $(1 - np)^2$, we shall have, by squaring, $1 - 4p + 2p^2 = 1 - 2np + n^2p^2$, or $-4 + 2p = -2n + n^2p$; whence

$$p = \frac{2n-4}{n^2-2}$$
 and $z = 1 - \frac{2n-4}{n^2-2} = \frac{n^2-2n+2}{n^2-2}$

or, if, for the sake of greater generality, $\frac{m}{n}$ be substituted for n in this last expression, we shall have

$$z = \frac{m^2 - 2mn + 2n^2}{m^2 - 2n^2}.$$

And since, by the two first equations, $y=v^2-u^2=u^2z^2-u^2=u^2=(z^2-1)u^2$, and $x=\frac{1}{2}(u^2-y)=\frac{1}{2}(2-z^2)u^2$, it is evident, that z must be some number greater than 1, and less than $\sqrt{2}$.

If, therefore, m=9 and n=5 we shall have

$$z = \frac{81 - 90 + 50}{81 - 50} = \frac{41}{31}, x = \frac{241}{31^2} \times \frac{u^2}{2} \text{ and } y = \frac{720}{31^2} \times u^2.$$

Or, taking $u=2\times31$, x=482, and y=2880, we have x=482, x+y=3362, and x+2y=6242, which are the numbers required.

4. It is required to divide a given square number into we such parts, that each of them shall be a square.*

Let a^2 = given square number, and x^2 and $a^2 - x^2$ its wo parts. Then since x^2 is a square, if only remains to make $a^2 - x^2$ a square.

For which purpose let its root =nx-a and we shall have $x^2-x^2=n^2x^2-2anx+a^2$, or $-x^2=n^2x^2-2anx$; whence,

y reduction, $x = \frac{2an}{n^2 + 1}$, the root of the first part; and $zx - a = \frac{2an^2}{n^2 + 1} - a = \frac{an^2 - a}{n^2 + 1}$ the root of the second.

Therefore $\left(\frac{2an}{n^2+1}\right)^2$ and $\left(\frac{an^2-a}{n^2+1}\right)^2$ are the parts re-

uired; where a and n may be any numbers taken at easure, provided n be greater than 1.

5. It is required to divide a given number, consisting of to known square numbers, into two other square numbers.

Let a^2+b^2 be the given numbers, and x^2 , y^2 the two quired numbers, whose sum, x^2+y^2 , is to be equal to $+b^2$.

Then it is evident, that if x be either greater or less an a, y will be accordingly less or greater than b. Let erefore x=a+mz, and y=b-nz, and we shall have a^z $2anz+m^2z^2+b^2-2bnz+n^2z^2=a^2+b^2$.

Or, by transposition and rejecting the terms which are

To this we may add the following useful property.

f s and r be any two unequal numbers, of which s is the greater, it can
a be readily shown, from the nature of the problem, that 2rs, s^2-r^2 and s^2+r^2

be the perpendicular base, and hypothemise of a right-angled triangle. From which expressions, two square numbers may be found, whose sum difference shall be square numbers; for $(2rs)^2 + (s^2 - rs)^2 = (s^2 + s^2)$, and $(s^2 + r^2)^2 - (2rs)^2 = (s^2 - r^2)^2$, or $(s^2 + r^2)^2 - (s^2 - rs)^2$.

common to each side of the equation, $m^2z^2 + n^2z^2 = 2bnz$ -2amz, or $m^2z+n^2z=2bn-2am$; whence

 $\frac{2bn-2am}{m^2+n^2}, x = \frac{2bmn+a(n^2-m^2)}{m^2+n^2}, y = \frac{2amn+b(m^2-n^2)}{m^2+n^2}$

where m and n may be any numbers, taken at pleasure, provided their assumed values be such as will render the values of x, y, and z, in the above expressions, all positive. 6. It is required to find two square numbers, such that

their difference shall be equal to a given number.

Let d= the given difference; which resolve into two

factors a, b; making a the greater and b the less.

Then, putting x= the side of the less square, and x+ $b = \text{side of the greater, we shall have } (x+b)^2 - x^2 = x^2$ $+2bx+b^2-x^2=d(ab)$ or $2bx+b^2=d(ab)$.

Whence, dividing each side of this equation by b, we shall have $x = \frac{a-b}{a}$ = the side of the less square sought,

and $x+b=\frac{a-b}{a}+b=\frac{a+b}{a}=$ the side of the greater.

If, for instance, d=60, take $a \times b = 30 \times 2$, and we shall have $x = \frac{30 - 2}{2} = 14$, and $x + 2 = \frac{30 + 2}{2} = 16$, or $16^2 - 14^2$

=256 - 196=50 the given difference.

7. As an instance of the great use of resolving formulæ of this kind into factors, let it be proposed in addition to what has been before said, to find two numbers x and y, such that the difference of their squares, x^2-y^2 , shall be an integral square.

Here the factors of x^2-y^2 , being x+y and x-y, we shall have $(x+y) \times (x-y) = x^2 - y^2$. And, since this product is to be a square, it will evidently become so, by making each of its factors a square, or the same multiple of a

square.

Let there be taken, therefore, for this purpose, $x+y=mz^2, x-y=ms^2.$

Then, by the question, we shall have $(x+y)\times(x-y)$ or its equal $x^2 - y^2 = m^2 r^2 s^2$; which is evidently a square. whatever may be the values of m, r, s.

But, by addition and subtraction, the above equations give, when properly reduced

 $x = \frac{m(r^2 + s^2)}{9}, y = \frac{m(r^2 - s^2)}{9},$

where, as above said, m, r, and s, may be assumed at pleasure. Thus, if we take m=2, we shall have $x=r^2+s^2$, and $y=r^2-s^2$, which expressions will obviously give integral values of x and y, if r and s be taken = any integral numbers.

8. It is required to find two numbers, such that, if either of them be added to the square of the other, the

sums shall be squares.

Let x and y be the numbers sought; and consequently $x^{2}+y$ and $y^{2}+x$ the expressions that are to be transformed into squares. Then if r-x be assumed for the side of the first square, we shall have $x^2 + y = r^2 - 2rx + x^2$, or $y = r^2$

$$-2rx$$
; and consequently $x = \frac{r^2 - y}{2r}$

And if s+y be taken for the side of the second square,

we shall have $y^2 + \frac{r^2 - y}{\varphi_r} = s^2 + 2sy + y^2$; or, by reducing the equation, $r^2-y=4rsy+2rs^2$, and consequently, by reduction, $y=\frac{r^2=2rs^2}{4rs+1}$, and $x=\frac{2r^2s+s^2}{4rs+1}$; where r and smay be any numbers, taken at pleasure, provided r be greater than 2s2.

9. It is required to find two numbers, such that their sum

and difference shall be both squares.

Let x and $x^2 - x$ be the two numbers sought; then, since their sum is evidently a square, it only remains to make their difference, $x^2 - 2x$, a square.

For this purpose, therefore, put its root =x-r and we

shall have $x^2 - 2x = x^2 - 2rx + r^2$;

Or, by transposition, and cancelling x2 on each side of the equation, $2rx-2x=r^2$; whence

$$x = \frac{r^2}{2r - 2}$$
, and $x^2 - 2x = \frac{1}{4} \left(\frac{r^2}{r - 1}\right) - \frac{r^3}{r - 1}$;

where r may be any number, taken at pleasure, provided

it be greater than 2.

10. It is required to find three numbers, such that not only the sum of all three of them, but also the sum of

every two shall be a square number.

Let 4x, x^2-4x and 2x+1, be the three numbers sought, then $4x+(x^2-4x)=x^2$, $(x^2-4x)+(2x+1)=x^2-2x+1$, and $4x+(x^2-4x)+(2x+1)=x+2x+1$, being all squares, it only remains to make 4x+(2x+1), or its equal, 6x+1, a square. For which purpose, let $6x+1=n^2$, and we shall

have, by transposition and division, $x = \frac{n^2 - 1}{6}$, whence,

$$\frac{4n^2-4}{6}, \frac{(n^2-1)}{36}, \frac{-4n^2-4}{6}, \text{ and } \frac{2n^2-2}{6}+1, \text{ or their equals } \frac{2n^2-2}{3}, \frac{n^4-26n^2+25}{36}, \text{ and } \frac{n^2+2}{3} \text{ are the numbers required.}$$

Where n may be any number, taken at pleasure, pro-

vided it be greater than 5.

QUESTIONS FOR PRACTICE.

1. It is required to find a number x, such that x+1 and x-1 shall be both squares. Ans. $x=\frac{s}{2}$.

2. It is required to find a number x, such that x+4 and x+7 shall be both squares. Ans. $\frac{51}{16}$.

3. It is required to find a number x, such that 10+x and 10-x shall be both squares. Ans. x=6.

4. It is required to find a number x, such that $x^2 + 1$

and x+1 shall be both squares. Ans. $\frac{4}{9}$.

5. It is required to find three integral square numbers, such that the sum of every two of them shall be squares.

Ans. 528, 5796, and 6325. 6. It is required to find two numbers x and y, such that $x^2 + y$ and $y^2 + x$ shall be both squares.

Ans. $x = \frac{3}{20}$, and $y = \frac{7}{10}$. 7. It is required to find three integral square numbers, that shall be in harmonical proportion.

Ans. 25, 49, and 1225.

8. It is required to find three integral cube numbers, x^3 , y^3 , and z^3 , whose sum may be equal to a cube.

Ans. 3, 4, and 5.

9. It is required to divide a given square number (100) into two such parts that each of them may be a square number.

Ans. 64, and 36.

10. It is required to find two numbers, such that their difference may be equal to the difference of their squares, and that the sum of their squares shall be a square number.

Ans. \(\frac{4}{3} \) and \(\frac{3}{3} \).

11. To find two numbers such, that if each of them be added to their product, the sums shall be both squares.

Ans. 3 and 5.

12. To find three square numbers in arithmetical progression.

Ans. 1, 25, and 49

13. To find three numbers in arithmetical progression such that the sum of every two of them shall be a square number.

Ans. 120½, 340½, and 1560½.

14. To find three numbers such, that if to the square of each the sum of the other two be added, the three sums shall be all squares.

Ans. 1, \(\frac{3}{4} \) and \(\frac{1}{4} \)

15. To find two numbers in proportion as 8 is to 15, and such that the sum of their squares shall be a square number.

Ans. 576 and 1080

46. To find two numbers such, that if the square of

each be added to their product the sums shall be both squares.

Ans. 9 and 16.

17. To find two whole numbers such, that the sum or difference of their squares, when diminished by unity shall be a square.

Ans. 8 and 9

13. It is required to resolve 4225, which is the square

of 65, into two other integral squares.

Ans. 2704 and 1521

19. To find three numbers in geometrical proportion such that each of them, when increased by a given number (19), shall be square numbers. Ans. $81, \frac{5}{4}$ and $\frac{25}{12} \frac{5}{96}$ 20. To find two numbers such, that if their product be

20. To find two numbers such, that if their product be added to the sum of their squares, the result shall be a square number.

Ans. 5, and 3, 8 and 7, 16 and 5, &c.

21. To find three whole numbers such, that if to the square of each the product of the other two be added, the three sums shall be all squares. . Ans. 9, 73, and 328

22. To find three square numbers such that their sum when added to each of their three sides, shall be all square

numbers.

Ans. $\frac{4418}{62926}$, $\frac{13254}{62926}$, and $\frac{19831}{62926}$ = roots required 23. To find three numbers in geometrical progression such, that if the mean be added to each of the extremes, the sums, in both cases shall be squares.

Ans. 5, 20, and 80 24. To find two numbers such, that not only each of them, but also their sum and their difference, when in-

creased by unity, shall be all square numbers.

Ans. 3024 and 5624 25. To find three numbers such, that whether their sum be added to, or subtracted from, the square of each of them, the numbers thence arising shall be all squares.

Ans. 406, 518, and 761 26. To find three square numbers such, that the sum of

their squares shall also be a square number.

Ans. 9, 16, and 144

27. To find three square numbers such, that the difference of every two of them shall be a square number. Ans. 485809, 34225, and 23409

28. To divide any given cube number (8) into three Ans. 1, 64 and 12,5 other cube numbers.

29. To find three square numbers such, that the difference between every two of them and the third shall be a square number. Ans. 1492, 2412, and 2692

30. To find three cube numbers such, that if from each of them a given number (1) be subtracted, the sum of the

remainders shall be a square number.

Ans. 4913, 21952, and 8.

OF THE

SUMMATION AND INTERPOLATION OF INFINITE SERIES.

The doctrine of Infinite Series is a subject which has engaged the attention of the greatest mathematicians, both of ancient and modern times; and, when taken in its whole extent, is, perhaps, one of the most abstruse and

difficult branches of abstract mathematics.

To find the sum of a series, the number of the terms of which is inexhaustible, or infinite, has been regarded by some, as a paradox, or a thing impossible to be done; but this difficulty will be easily removed, by considering that every finite magnitude whatever is divisible in infinitum, or consists of an indefinite number of parts, the aggregate, or sum, of which, is equal to the quantity first proposed.

A number actually infinite is, indeed, a plain contradiction to all our ideas; for any number that we can possibly conceive, or of which we have any notion, must always be determinate and finite; so that a greater may still be assigned, and a greater after this; and so on, without a possibility of ever soming to an end of the increase or ad-

dition.

This inexhaustibility, therefore, in the nature of numbers, is all that we can distinctly comprehend by their infinity; for though we can easily conceive that a finite quantity may become greater and greater without end, yet we are not, by that means, enabled to form any notion of the ultimatum, or last magnitude, which is incapable of farther augmentation.

Hence, we cannot apply to an infinite series the common notion of a sum, or of a collection of several particular numbers, which are joined and added together, one after another; as this supposes that each of the numbers composing that sum, is known and determined. But as every series generally observes some regular law, and continually approaches towards a term, or limit, we can easily conceive it to be a whole of its own kind, and that it must have a certain real value, whether that value be determinable or not.

Thus in many series, a number is assignable, beyond which no number of its terms can ever reach, or indeed, be ever perfectly equal to it; but yet may approach towards it in such a manner, as to differ from it by less than any quantity that can be named. So that we may justly call this the value or sum of the series; not as being a number found by the common method of addition, but such a limitation of the value of the series, taken in all its infinite capacity, that, if it were possible to add all the terms together, one after another, the sum would be equal to that number.

In other series, on the contrary, the aggregate, or value of the several terms, taken collectively, has no limitation; which state of it may be expressed by saying, that the sum of the series is infinitely great; or, that it has no determinate or assignable value, but may be carried on to such a length, that its sum shall exceed any given number whatever.

Thus, as an illustration of the first of these cases, it may be observed, that if r be the ratio, g the greatest term, and l the least, of any decreasing geometric series, the sum, according to the common rule, will be (rg-l), (r-1): and if we suppose the less extreme, l, to be diminished till it becomes =0, the sum of the whole series will be $rg \div (r-1)$: for it is demonstrable, that the sum of no assignable number of terms of the series can ever be equal to that quotient; and yet no number less than it will ever be equal to the value of the series.

Whatever consequences, therefore, follow from the supposition of $rg \div (r-1)$ being the true and adequate value of the series, taken in all its infinite capacity, as if all the parts were actually determined, and added together, no assignable error can possibly arise from them, in any operation or demonstration where the sum is used in that sense; because, if it should be said that the series exceeds that

value; it can be proved, that this excess must be less than any assignable difference; which is, in effect, no differance at all; whence the supposed error cannot exist, and consequently $rg \div (r-1)$ may be looked upon as expressing the true value of the series, continued to infinity.

We are, also, farther satisfied of the reasonableness of this doctrine, by finding, in fact, that a finite quantity is frequently convertible into an infinite series, as appears in the case of circulating decimals. Thus two-thirds expressed decimally is $\frac{2}{3}$ =.66666, &c. = $\frac{6}{10}$ + $\frac{1}{10}$ +

With respect to the processes by which the summation of various kinds of infinite series are usually obtained, one of the principal is by the method of differences pointed out and illustrated in Prob. IV. next following.

Another method is that first employed by James and John Bernoulli, which consists in resolving the given series into several others of which the summation is known; or by subtracting from an assumed series, when put == s, the same series, deprived of some of its first terms; in which case a new series will arise, the sum of which will be known.

A third method which is that of Demoivre, consists in putting the sum of the series =s, and multiplying each side of the equation by some binomial or trinomial expression, which involves the powers of the unknown quantity x, and certain known co-efficients; then taking x, after the process is performed, of such a value that the assumed binomial, &c. shall become =0, and transposing some of the first terms, a series will arise, the sum of which will be known, as before.

Each of which methods, modified so as to render it more commodious in practice, together with several other artifices for the same purpose, will be found sufficiently elucidated in the miscellaneous questions succeeding the following problems.

PROBLEM I.

Any series being given to find its several orders of differences.

RULE.

1. Take the first term from the second, the second from the third, the third from the fourth, &c. and the remainders will form a new series, called the first order of differences.

2. Take the first term of this last series from the second, the second from the third, the third from the fourth, &c. and the remainders will form another new series, called the second order of differences.

3. Proceed in the same manner, for the third, fourth, fifth, &c. orders of differences; and so on till they terminate, or are carried as far as may be thought necessary*.

EXAMPLES.

1. Required the several orders of differences of the series 1, 22, 32, 42, 52, 62, &c.

1, 4, 9, 16, 25, 36, &c.

3, 5, 7, 9, 11, &c. 1st diff.

2, 2, &c. 2d diff.

0, 0, 0, &c. 3d diff.

2. Required the several orders of differences of the series 1, 23, 33, 43, 53, 63, &c.

1, 8, 27, 64, 125, 216, &c.

7, 19, 37, 61, 91, &c. 1st diff.

12, 18, 24, 30, &c. 2d diff.

6, 6, &c. 3d diff. 0, 0, &c. 4th diff.

^{*} When the several terms of the series continually increase, the differences will be all positive; but when they decrease, the differences will be negative and positive alternately.

3. Required the several orders of differences of the series 1, 3, 6, 10, 15, 21, &c.

Ans. 1st, 2, 3, 4, 5, &c.; 2d, 1, 1, 1, &c.

4. Required the several orders of differences of the series 1, 6, 20, 50, 105, 196, &c.

Ans. 1st, 5, 14, 30, 55, 91, &c.; 2d, 9, 16, 25, 36, &c.; 3d, 7, 9, 11, &c.; 4th, 2, 2, &c.

5. Required the several orders of differences of the series $\frac{1}{2}$, $\frac{1}{4}$, $\frac{1}{8}$, $\frac{1}{16}$, $\frac{1}{32}$, &c.

PROBLEM II.

Any series a, b, c, d, e, &c. being given, to find the first term of the nth order of differences.

RULE.

Let I stand for the first term of the nth differences.

Then will
$$a - nb + n \cdot \frac{n-1}{2}c - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3}d + n \cdot \frac{n-1}{2}$$

 $\frac{n-2}{3} \cdot \frac{n-3}{4}e$, &c. to $n+1$ terms $=\delta$, when n is an even

number. • And $-c + nb - n \cdot \frac{n-1}{2}c + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3}d - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3}$.

 $\frac{n-3}{3}e$ &c. to n+1 terms $=\delta$, when n is an odd number.*

ENAMPLES.

1. Required the first term of the third order of differences of the series 1, 5, 15, 35, 70, &c.

Here a, b, c, d, e, &c. = 1, 5, 15, 35, 70, &c. and <math>n=3.

Whence
$$-\alpha + nb - n \cdot \frac{n-1}{2}c + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3}d = -\alpha$$

^{*} When the terms of the several orders of differences happen to be very great, it will be more convenient to take the logarithms of the quantities concerned whose differences will be smaller; and when the operation is this hed, the quantity answering to the last logarithm may be easily found.

+3b-3c+d=-1+15-45+35=4= the first term required.

2. Required the first term of the fourth order of differences of the series 1, 8, 27, 64, 125, &c.

Here a, b, c, d, e, &c. = 1, 8, 27, 64, 125, &c. and n=4. Whence $a-nb+n.\frac{n-1}{2}c-n.\frac{n-1}{2}.\frac{n-2}{3}d+n.\frac{n-1}{2}$

$$\frac{n-2}{3} \cdot \frac{n-3}{4} e = a-4b+6c-4d+e=1-32+162-256+125$$

=0; so that the first term of the fourth order is 0.

3. Required the first term of the eighth order of differences of the series, 1, 3, 9, 27, 81, &c.* Ans. 256.

4. Required the first term of the fifth order of differences of the series, $1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}, \frac{1}{64}$, &c.

Ans. $-\frac{1}{36}$.

PROBLEM III.

To find the *n*th term of the series, a, b, c, d, e, &c. when the differences of any order become at last equal to each other.

RULE.

Let d', d'', d^{iv} , d^{iv} , &c. be the first of the several orders of differences, found as in the last problem.

Then will $a + \frac{n-1}{1}d' + \frac{n-1}{1} \cdot \frac{n-2}{2}d'' + \frac{n-1}{1} \cdot \frac{n-2}{2} \cdot \frac{n-3}{3}$ $d''' + \frac{n-1}{1} \cdot \frac{n-2}{2} \cdot \frac{n-3}{3} \cdot \frac{n-4}{4} \cdot d^{1v}$ &c. =nth term required.

EXAMPLES.

1. It is required to find the twelfth term of the series 2, 6, 12, 20, 30, &c.

^{*} The labour, in questions of this kind may be often abridged, by putting ciphers for some of the terms at the beginning of the series; by which means several of the differences will be equal to 0, and the answer on that account, obtained in fewer terms.

Here 4 and 2 are the first terms of the differences.

Let, therefore, 4=d', 2=d'', and n=12. Then $a+\frac{n-1}{1}d'+\frac{n-1}{1}\frac{n-2}{2}d'=2+11d'+55d''=2$

+44+110=156=12th term, or the answer required.

2. Required the twentieth term of the series, 1, 3, 6, 10, 15, 21, &c.

Here 2 and 1 are the first terms of the differences. Let, therefore, 2=d', 1=d', and n=20.

Then $a + \frac{n-1}{1}d' + \frac{n-1}{1} \cdot \frac{n-2}{2}d' = 1 + 19d' + 171d'' = 1 + 171d'' = 171$

38+171=210=20th term required.

3. Required the fifteenth term of the series, 1, 4, 9, 16, 25, 36, &c.

Ans. 225.

4. Required the twentieth term of the series, 1, 8, 27, 64, 125, &c.

5. Required the thirtieth term of the series, $1, \frac{1}{3}, \frac{1}{6}$,

 $\frac{1}{10}$, $\frac{1}{15}$, $\frac{1}{21}$, &c.

PROBLEM IV.*

To find the sum of n terms of the series, a, b, c, d, e, &c. when the differences of any order become at last equal to each other.

^{*}When the differences in this or the former rule are finally =0, any term, or the sum of any number of the terms, may be accurately the termined; but if the differences do not vanish, the result is only an approximation; which, however, may be often very usefully applied in resolving various questions that may occur in this branch of the subject, and which will become continually nearer the truth as the difference diminish.

RULE.

Let d', d'', d''', d^{iv} , &c. be the first of the several orders of differences.

Then will
$$na+n$$
. $\frac{n-1}{2}d'+n$. $\frac{n-1}{2}$. $\frac{n-2}{3}d''+n$. $\frac{n-1}{2}$. $\frac{n-2}{3}d''+n$. $\frac{n-1}{2}$. $\frac{n-2}{3}\frac{n-3}{4}d'''+n$. $\frac{n-1}{2}\frac{n-2}{3}\frac{n-3}{4}\frac{n-4}{5}d^{lv}$, &c. = to the sum of n terms of the series.

EXAMPLES.

1. Required the sum of n terms of the series, 1, 2, 3, 4, 5, 6, &c.

Where 1 and 0 are the first terms of the differences, Let, therefore, a=1, d=1, and d'=0.

Then will
$$na+n.\frac{n-1}{2}d'=n+\frac{n^2-n}{2}=\frac{n^2+n}{2}=\text{sum of } n$$
 terms, as required.

2. Required the sum of n terms of the series, 1^2 , 2^2 , 3^2 , 4^2 , 5^2 , &c., or 1, 4, 9, 16, 25, &c.

Where 3 and 2 are the first terms of the differences, Let therefore, a=1, d=3, and d'=2.

Then will
$$na+n$$
. $\frac{n-1}{2}d+n$. $\frac{n-1}{2}$. $\frac{n-2}{3}d''=n+3n$. $\frac{n-1}{2}+2n$. $\frac{n-1}{2}$. $\frac{n-2}{3}=\frac{3n^2-3n}{2}+\frac{n^3-3n^2+2n}{3}=\frac{3n^2-3n}{3}$

$$\frac{n \times (n+1) \times (2n+1)}{s}$$
 = sum of π terms as required.

3. Required the sum of n terms of the series, 13, 23, 33, 43, 53, &c., or 1, 8, 27, 64, 125, &c.

&c.

Here 1, 8, 27, 64, 7, 19, 37,

12, 18, 24, 6,

&c.

Where the first terms of the differences are 7, 12, and 6.

Let, therefore, a=1, d'=7, d'=12, and d''=6.

Then will $na + n \cdot \frac{n-1}{2}d + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3}d + n \cdot \frac{n-1}{2}$

$$\frac{n-2}{3} \cdot \frac{n-3}{4} d''' = n + 7n \cdot \frac{n-1}{2} + 12n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} + 6n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \frac{n-3}{4} = n + \frac{7n^2 - 7n}{2} + 2n^3 - 6n^2 + 4n + \frac{n-2}{2} + \frac{n-3}{3} + \frac{n-3}{4} = n + \frac{7n^2 - 7n}{2} + 2n^3 - 6n^2 + 4n + \frac{n-2}{2} + \frac{n-3}{3} + \frac{n-3}{4} = n + \frac{n-3}{2} + \frac{n-3}{4} + \frac$$

$$\frac{n-1}{2} \cdot \frac{n}{3} \cdot \frac{n}{4} = n + \frac{n}{2} + 2n^3 - 6n^2 + 4n + \frac{n^4 - 6n^3 + 11n^2 - 6n}{4} = \frac{4n}{4} + \frac{14n^2 - 14n}{4} + \frac{8n^3 - 24n^2 + 16n}{4}$$

$$+\frac{n^4-6n^3+11n^2-6n}{4} = \frac{n^4+2n^3+n^2}{4} = \text{sum of } n \text{ terms as required.}$$

4. Required the sum of n terms of the series, 2, 6, 12, Ans. $\frac{n\times(n+1)\times(n+2)}{n+2}$ 20, 30, &c.

5. Required the sum of n terms of the series, 1, 3, 6, Ans. $\frac{n}{1} \cdot \frac{n+1}{2} \cdot \frac{n+2}{3}$ 10, 15, &c.

6. Required the sum of n terms of the series, 1, 4, 10, 20. 35, &c.

7. Required the sum of n terms of the series 14, 24, 34, 44, &c., or 1, 16, 81, 256, &c.

Ans. $\frac{n^5}{5} + \frac{n^4}{9} + \frac{n^3}{3} -$

8. Required the sum of *n* terms of the series 1^5 , 2^5 , 3^5 , 5^5 , &c. Ans. $\frac{n^6}{6} + \frac{n^5}{2} + \frac{5n^4}{12} - \frac{n^2}{12}$ 45, 55, &c.

PROBLEM V.

The series a, b, c, d, ϵ , &c. being given, whose terms are an unit's distance from each other, to find any intermediate term by interpolation.

RULE.

Let x be the distance of any term y, that is to be interpolated. from the first term, and d', d', d', &c. the terms of the differences.

Then will
$$a + xd' + x \cdot \frac{x-1}{2}d' + x \cdot \frac{x-1}{2} \cdot \frac{x-2}{3}d'' + x \cdot \frac{x-1}{2} \cdot \frac{x-3}{3}d''' + x \cdot \frac{x-1}{2} \cdot \frac{x-3}{3}d'' + x \cdot \frac{x-1}{2} \cdot \frac{x-1}{3}d'' + x \cdot \frac{x-1}{2}d' + x \cdot \frac{x-1}{3}d'' + x \cdot$$

EXAMPLES.

1. Given the logarithmic sines of 1° 0′, 1° 1′, 1° 2′ and 1° 3′, to find the log. sine of 1° 1′ 40″.

Here 1°0′ 1°1′ 1°2′ 1°3′ Sines 8.2418553 8.2490332 8.2560943 8.2630424 71779 70611 69481 —1168 —1130

Whence the first terms of the differences are 71779,—1168, and 38.

Let, therefore, $x=1^{\circ}$ 1' $40''-1^{\circ}$ 0'=1' $40''=1^{\circ}$ = 1 distance of y, the term to be interpolated; and d'=71779, d''=1168, and d''=38.

Then will $y=a+xd'+x.\frac{x-1}{2}d'+x.\frac{x-1}{2}.\frac{x-2}{3}d''=a+\frac{5}{3}d'+\frac{5}{9}d'-\frac{5}{81}d'''=3.2418553+.0119631+0000694-$

.0000002=8.2538876=sine of 1° 1′ 40″, as was required. 2. Given the series $\frac{1}{50'51'52'53'54}$, &c. to find the term which stands in the middle between the two terms $\frac{1}{52}$ and

 $\frac{1}{53}$. Ans. $\frac{1}{105}$

3. Given the natural targents of 88° 54', 88° 55', 88° 56', 88° 57', 88° 58', 88° 59', to find the tangent of 88° 56' 11". Ans. 55.7-11144

PROBLEM VI.

Having given a series of equidistant terms, a, b, c, d, e, &c. whose first differences are small; to find any intermediate term by interpolation.

RULE.

Find the value of the unknown quantity in the equation which, stands against the given number of terms, in the following table, and it will give the term required.*

1. a-b=0

2.
$$a-2b+c=0$$

3. $a-3b+3c-d=0$
4. $a-4b+6c-4d+e=0$
5. $a-5b+10c-10d+5e-f=0$
6. $a-6b+15c-20d+15e-6f+g=0$
or $\begin{vmatrix} a-nb+n & \frac{n-1}{2}c-n & \frac{n-1}{2} & \frac{n-2}{3}d+1 \\ n & \frac{n-1}{2} & \frac{n-2}{3} & \frac{n-3}{4}e, &c. = 0. \end{vmatrix}$

EXAMPLES.

1. Given the logarithms of 101, 102, 104, and 105, to find the logarithm of 103.

^{*} The more terms are given, in any series of this kind, the more accurately will the equation that is to be used approximate towards the true result, or answers required.

Here the number of terms are 4. And against 4, in the table, we have a - 4b + 6c - 4d + e=0; or $c = \frac{4 \times (b+d) - (a+e)}{}$ = value of the unknown quantity, or term to be found.

a=2.0043214Where, taking the logs of b = 2.0086002101, 102, 104, and 105 d=2.0170333e = 2.0211893

> And consequently $4 \times (b+d) = 16.1025340$ a+e=4.0255107

6)12.0770233

2.0128372= log. of 103, as

required.

2. Given the cube roots of 45, 46, 47, 48, and 49, to find the cube root of 50. Ans. 3.684031.

3. Given the logarithms of 50, 51, 52, 54, 55, and 56, to find the logarithm of 53. Ans. 1.7242758695.

PROMISCUOUS EXAMPLES RELATING TO SERIES.

1. To find the sum (s) of n terms of the series, 1, 2, 3, 4, 5, 6, &c.

And consequently n(n+1)=2s; or $s=\frac{1}{2}=sum re$ quired.

2. To find the sum (s) of n terms of the series, 1, 3, 5, 7, 9, 11, &c.

First,
$$1+3+5+7+9$$
 &c. . . . $(2n-1)=s$.
And $(2n-1)+(2n-3)+(2n-5)+...+1=S$,

Therefore by addition, 2n+2n+2n+2n+2n+&c....2n=2s.And consequently $2n \times n = 2s$;

Or
$$s = \frac{2n^2}{2} = n^2 = \text{sum required.}$$

3. Required the sum (s) of n terms of the series, a+ (a+d)+(a+2d)+(a+3d)+(a+4d) &c.

First, a + (a+d) + (a+2d) + (a+3d) &c. ... +

 $\{a+(n-1)d\}=s.$

And a+(nd-d)+a+(nd-2d)+a+(nd-3d)+a+(nd-4d) &c. ... a=s.

Therefore, by addition, 2a+(nd-d)+2a+(nd-d)+2a + (nd - d) &c. + 2a + (nd - d) = 2s.

And consequently $(2a+nd-d) \times n=2s$;

Or
$$s = (2a + nd - d) \times \frac{n}{2} = sum required.$$

Or the same may be done in a different manner, as follows:

$$a+(a+d)+(a+2d)+(a+3d)+(a+4d) &c.$$

$$=\begin{vmatrix} (+1+1+1+1+1&c.)\times a\\ (+0+1+2+3+4&c.)\times d \end{vmatrix} = s$$
But n terms of $1+1+1+1+1 &c. = n$.

And *n* terms of $0+1+2+3+4 &c. \frac{n \times (n-1)}{9}$.

Whence
$$s=na+\frac{n\times(n-1)d}{2}=\left\{2a+d(n-1)\right\}\times\frac{n}{2}$$

which is the same answer as before.

4. To find the sum (s) of n terms of the series 1, x, x2, x3, x4, &c.

First, $1 + x + x^2 + x^3 + x^4$, &c. . . . $x^{n-1} = s$. And $x+x^2+x^3+x^4+x^5$, &c. . . . $x^n=sx$. Whence, by subtraction, $x^n - 1 = sx - s_n$

Or
$$s = \frac{x^n - 1}{x - 1} = \text{sum required}$$
.

And, when x is a proper fraction, the sum of the series, continued ad infinitum may be found in the same man-

Thus, putting
$$1+x+x^2+x^3+x^4+x^5$$
, &c.=s,

We shall have $x+x^2+x^3+x^4+x^5$, &c.=sx, And consequently -1=sx-s; or s-sx=1,

Whence $s = \frac{1}{1-x} = \text{sum of an infinite number of terms,}$ as was to be found.

5. Required the sum (s) of the circulating decimal .999999 &c. continued ad infinitum.

First, .999999 &c. =
$$\frac{9}{10} + \frac{9}{100} + \frac{9}{1000}$$
 &c. = $9(\frac{1}{10} + \frac{1}{100} + \frac{1}{1000} + \frac{1}{10000} + \frac{1}{6}$.)=s. Or, $\frac{1}{10} + \frac{1}{100} + \frac{1}{1000} + \frac{1}{10000} + \frac{1}{6}$. Therefore, $1 + \frac{1}{10} + \frac{1}{100} + \frac{1}{1000} + \frac{1}{6}$. $= \frac{10s}{9}$.

And consequently $1 = \frac{10s}{9} - \frac{s}{9} = \frac{9s}{9} = s$;

Whence s=1= sum of the series. 6. Required the sum (s) of the series $a^2+(a+d)^2+(a+2d)^2+(a+3d)^2+(a+4d)^2$, &c. continued to n terms.

Hen

First,
$$a^2 = a^2$$

 $(a+d)^2 = a^2 + 2 \times 1ad + 1d^2$
 $(a+2d)^2 = a^2 + 2 \times 2ad + 4d^2$
 $(a+3d)^2 = a^2 + 2 \times 3ad + 9d^2$
 $(a+4d)^2 = a^2 + 2 \times 4ad + 16d^2$

Whence

whence
$$s = \begin{vmatrix} \text{Sum of } n \text{ terms of } (1+1+1+1+&c.)a^2 \\ + \dots & \text{ditto of } (1+2+3+4+&c.)ad \\ + \dots & \text{ditto of } (1+4+9+16+&c.)d \\ \text{But } n \text{ terms of } 1+1+1+1+&c.=n. \end{vmatrix}$$

And of
$$1+2+3+4$$
 &c. $=\frac{n(n-1)}{1\cdot 2}$
Also of $1+4+9+$ &c. $=\frac{n(n-1)(2n-1)}{1\cdot 2\cdot 3}$

SUMMATION OF INFINITE SERIES.

Therefore $s=na^2+n(n-1)$ $ad+\frac{n(n-1)(2n-1)}{1}d^2=$

the whole sum of the series to n terms.

7. Required the sum (s) of the series $a^3+(a+d)^3+(a+d)^3$ $+2d)^3+(\alpha+3d)^3+(\alpha+4d)^3$ &c. continued to n terms.

First, $a^3 = a^3$

$$\begin{array}{l} (a+d)^3 = a^3 + 3 \times 1a^2 d + 3 \times 1ad^2 + 1d^2 \\ (a+2d)^3 = a^3 + 3 \times 2a^2 d + 3 \times 4ad^2 + 8d^3 \\ (a+3d)^3 = a^3 + 3 \times 3a^2 d + 3 \times 9ad^2 + 27d^3 \\ (a+4d)^3 = a^3 + 3 \times 4a^2 d + 3 \times 16ad^2 + 64d^3 \end{array}$$

 $(a+5d)^3=a^3+3\times5a^2d+3\times25ad^2+125d^3$

&c. Whence

whence
$$s = \begin{vmatrix}
\text{Sum of } n \text{ terms of } (1+1+1+1&c.)a^3 \\
+ ... \text{ ditto of } (1+2+3+4&c.)3a^2d \\
+ ... \text{ ditto of } (1+4+9+16&c.)3ad^2 \\
+ ... \text{ ditto of } (1+8+27+64&c.)d^3$$
But $n \text{ terms of } 1+1+1+1+1&c.=n$.

Ditto . . . of
$$1+2+3+4$$
 &c. = $\frac{n(n-1)}{1}$

Ditto . . . of
$$1+4+9+16$$
 &c. = $\frac{n(n-1)(2n-1)}{1 \cdot 2 \cdot 3}$

Ditto . . . of
$$1+8+27+64$$
 &c. $=\frac{n^4-2n^3+n^2}{2\times 2}$
Therefore, $s=na^3+\frac{n(n-1)3a^2d}{2}+\frac{n(n-1)(2n-1)3ad^2}{6}$

Therefore,
$$= na^3 + \frac{n(n-1)3a^2d}{2} + \frac{n(n-1)(2n-1)3ad^2}{6} + \frac{(n^4 - 2n^3 + n^2)d^3}{4} = \text{sum of } n \text{ terms, as was to be found.}$$

8. Required the sum (s) of n terms of the series 1+3 +7+15+31, &c.

The terms of this series are evidently equal to 1, (1+ 2), (1+2+4), (1+2+4+8), &c. or to the successive sums of the geometrical series, 1, 2, 4, 8, 16, &c.

Let, therefore, a=1 and r=2, and we shall have

 $a+ar+ar^2+ar^3+ar^4$ &c. =1+2+4+8+16, &c. But the successive sums of 1, 2, 3, 4, &c. terms of this series are,

1.
$$\frac{ar-a}{r-1} = (r-1) \times \frac{a}{r-1}$$

2. $\frac{ar^2-a}{r-1} = (r^2-1) \times \frac{a}{r-1}$
3. $\frac{ar^3-a}{r-1} = (r^3-1) \times \frac{a}{r-1}$
4. $\frac{ar^4-a}{r-1} = (r^4-1) \times \frac{a}{r-1}$

Therefore $s = \frac{a}{r-1} \times \begin{vmatrix} n \text{ terms of } r + r^2 + r^3 + r^4 & \&c. \\ -n \text{ terms of } 1 + 1 + 1 + 1 & \&c. \\ \text{But } 1 + 1 + 1 + 1 + 1 + 1 & \&c. \\ = n \end{vmatrix}$

And
$$r+r^2+r^3+r^4+&c.=(r^n-1)\times \frac{r}{r-1}$$

Whence $s = \frac{r(r^n - 1)}{r - 1} \times \frac{a}{r - 1} - n \times \frac{a}{r - 1} = 2(2^n - 1) - n =$ whole sum required.

9. It is required to find the sum of *n* terms of the series. $\frac{1}{1} + \frac{3}{2} + \frac{7}{4} + \frac{15}{8} + \frac{31}{61} + \frac{63}{39}$, &c.

Here the terms of this series are the successive sums of the geometrical progression $\frac{1}{1} + \frac{1}{9} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} \&c$

Let, therefore, a=1 and r=2, then will $\frac{1}{1} + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} \&c. = a + \frac{a}{r} + \frac{a}{r^2} + \frac{a}{r^3} + \frac{a}{r^4} + \frac{a}{r^5} + \frac{a}{r^6} \&c.$

But the successive sums of 1, 2, 3, 4, &c. terms of this series are.

1.
$$\frac{(r-1)\times a}{(r-1)\times 1} = (r-1)\times \frac{a}{r-1}$$
2. $\frac{(r^2-1)\times a}{(r^2-1)\times r} = (r-\frac{1}{r})\times \frac{a}{r-1}$
3. $\frac{(r^3-1)\times a}{(r-1)\times r^2} = (r-\frac{1}{r^2})\times \frac{a}{r-1}$
4. $\frac{(r^4-1)\times a}{(r-1)\times r^3} = (r-\frac{1}{r^3})\times \frac{a}{r-1}$
\$c.

Therefore

$$s = \frac{a}{r-1} \times \begin{vmatrix} n \text{ terms of } r + r + r + r + r & \&c. \\ -n \text{ terms of } \frac{1}{1} + \frac{1}{r} + \frac{1}{r^2} + \frac{1}{r^3} \&c. \end{vmatrix}$$

These being the two series derived from the above expressions,

But
$$r+r+r+r+r+r+4$$
 &c. $=nr$.
 $\frac{1}{1}+\frac{1}{r}+\frac{1}{r^2}+\frac{1}{r^3}$ &c. $=\frac{r^n-1}{(r-1)r^{-1}}$.

 $s = \frac{a}{r-1} \times (nr - \frac{r^n - 1}{(r-1)r^{n-1}}) = \frac{(n-1)2^n + 1}{2^{n-2}} = \text{sum required}.$

10. Required the sum (s) of the infinite series of the reciprocals of the triangular number $\frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \frac{1}{10} + \frac{1}{15}$ &c.

Let
$$\frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \frac{1}{10} \&c.$$
 ad infinitum =s.
Or $\frac{1}{1.1} + \frac{1}{1.3} + \frac{1}{2.3} + \frac{1}{2.5} \&c.$... =s.
Then $\frac{1}{1.2} + \frac{1}{2.3} + \frac{1}{3.4} + \frac{1}{4.5} \&c.$... = $\frac{s}{2}$.
That is, $(\frac{1}{1} - \frac{1}{2}) + (\frac{1}{2} - \frac{1}{3}) + (\frac{1}{3} - \frac{1}{4}) + (\frac{1}{4} - \frac{1}{5}) \&c. = \frac{s}{2}$.
Or, $\begin{vmatrix} \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} \&c. \\ \frac{1}{2} + \frac{1}{3} - \frac{1}{4} - \frac{1}{5} - \frac{1}{6} - \frac{1}{7} \&c. \end{vmatrix} = \frac{s}{2}$

Whence $\frac{s}{2} = \frac{1}{1}$; or s = 2 = sum required.

11. And if it be required to find the sum of n terms of the same series, $\frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \frac{1}{10} + \frac{1}{15} &c.$

Let
$$z = \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} \&c.$$
 to $\frac{1}{n}$.

Then
$$z - \frac{1}{1} = \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} &c.$$
 to $\frac{1}{n}$.

And $z - \frac{1}{1} + \frac{1}{n+1} = \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} &c.$ to $\frac{1}{n+1}$.

Therefore $\frac{1}{1} - \frac{1}{n+1} = \frac{1}{2} + \frac{1}{6} + \frac{1}{12} + \frac{1}{20} &c.$ to $\frac{1}{n-1}$.

Or $\frac{n}{n+1} = \frac{1}{2} + \frac{1}{6} + \frac{1}{12} + \frac{1}{20} &c.$ to $\frac{1}{n(n+1)}$.

Whence $\frac{2n}{n+1} = \frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \frac{1}{10} &c.$ to $\frac{2}{n(n+1)}$.

Or $\frac{1}{1} + \frac{1}{3} + \frac{1}{6} + \frac{1}{10} + \frac{1}{15} &c.$ to $\frac{2}{n(n+1)} = \frac{2n}{n+1} = \text{sum of } n \text{ terms}$ of the series, as was required.

12. Required the sum of the infinite series, $\frac{1}{1 \cdot 2 \cdot 3} + \frac{1}{2 \cdot 3 \cdot 4} + \frac{1}{3 \cdot 4 \cdot 5} + \frac{1}{4 \cdot 5 \cdot 6} &c.$

Let $z = \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} &c.$ ad infinitum.

Then $z - \frac{1}{1} = \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} &c.$ by transposition.

And $1 = \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \frac{1}{4 \cdot 5} &c.$ by subtraction.

Or $1 - \frac{1}{2} = \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \frac{1}{4 \cdot 5} + \frac{1}{5 \cdot 6} &c.$ by transposition.

Whence $\frac{1}{2} = \frac{4}{1 \cdot 4 \cdot 3} + \frac{6}{2 \cdot 9 \cdot 4} + \frac{3}{3 \cdot 16 \cdot 5} &c.$ by subtraction.

Or $\frac{1}{2} = \frac{2}{1 \cdot 2 \cdot 3} + \frac{2}{2 \cdot 3 \cdot 4} + \frac{2}{3 \cdot 4 \cdot 5 \cdot 6} &c.$

And $\frac{1}{2} \div 2 = \frac{1}{1 \cdot 2 \cdot 3} + \frac{2}{2 \cdot 3 \cdot 4} + \frac{2}{3 \cdot 4 \cdot 5 \cdot 6} &c.$

And $\frac{1}{2} \div 2 = \frac{1}{1 \cdot 2 \cdot 3} + \frac{2}{2 \cdot 3 \cdot 4} + \frac{2}{3 \cdot 4 \cdot 5 \cdot 6} &c.$

But $\frac{1}{2} \div 2 = \frac{1}{4}$; therefore $\frac{1}{1.2.3} + \frac{1}{2.3.4} + \frac{1}{3.4.5} + \frac{1}{4.5.6}$ as

 $infinitum = \frac{1}{4}$, which is the sum required.

13. And if it were required to find the sum of n terms of the same series $\frac{1}{1.2.3} + \frac{1}{2.3.4} + \frac{1}{3.4.5} + \frac{1}{4.5.6}$ &c.

Let
$$z = \frac{1}{1.2} + \frac{1}{2.3} + \frac{1}{3.4} + &c. \text{ to } \frac{1}{n(n+1)}$$

Then $z = \frac{1}{2} = \frac{1}{2.3} + \frac{1}{3.4} + \frac{1}{4.5} &c. \text{ to } \frac{1}{n(n+1)}$.
And $z = \frac{1}{2} + \frac{1}{(n+1)(n+2)} = \frac{1}{2.3} + \frac{1}{3.4} + \frac{1}{4.5} + \frac{1}{5.6} + \frac{1}{6.7} + \frac{1}{$

 $\frac{1}{7.8}$ &c. continued to $\frac{1}{(n+1)(n+2)}$ terms.

Therefore $\frac{1}{2} - \frac{1}{(n+1)(n+2)} = \frac{2}{1.2.3} + \frac{2}{2.3.4} + \frac{2}{3.4.5}$ &c. to *n* terms, by subtraction.

Whence $\frac{1}{1} - \frac{1}{2 \cdot (n+1)(n+2)} = \frac{1}{1 \cdot 2 \cdot 3} + \frac{1}{2 \cdot 3 \cdot 4} + \frac{1}{3 \cdot 4 \cdot 5}$ &c. to n terms, by division.

And consequently $\frac{1}{1.2.3} + \frac{1}{2.3.4} + \frac{1}{3.4.5}$ &c. continued to n

terms =
$$\frac{1}{4} - \frac{1}{2 \cdot (n+1)(n+2)}$$
 = sum required.

14. Required the sum (s) of the series $\frac{1}{2} - \frac{1}{4} + \frac{1}{8} - \frac{1}{16} + \frac{1}{16}$

$$\frac{1}{32}$$
 - &c. continued ad infinitum.

Let
$$x = \frac{1}{2}$$
 and $s = \frac{z}{1+x}$.
Then $\frac{z}{1+x} = x(1-x+x^2-x^3+x^4 \& c.)$
And $z = (1+x) \times (x-x^2+x^3-x^4+x^5 \& c.$
Whence, by multiplication,
 $x-x^2+x^3-x^4+x^5 \& c.$
 $\frac{1+x}{1+x}$
 $\frac{1+x}{1+x^2-x^3+x^4+x^5 \& c.}$

Whose sum is
$$=x+0+0+0+0&c$$
.

252

Therefore z=x, and $x-x^2+x^3-x^4+x^5$ &c. $=\frac{x}{1+x}$ Or $\frac{1}{o}-\frac{1}{4}+\frac{1}{3}-\frac{1}{16}+\frac{1}{30}$ &c. $=\frac{\frac{1}{2}}{1+1}=\frac{1}{3}=$ sum required.

15. Required the sum of the series $\frac{1}{2} + \frac{2}{4} + \frac{3}{8} + \frac{4}{16} + \&c.$ continued ad infinitum.

Let
$$x = \frac{1}{2}$$
 and $s = \frac{z}{(1-x)^2}$.
Then $\frac{z}{(1-x)^2} = x + 2x^2 + 3x^3 + 4x^4 + 5x^5 \&c$.
And $z = (1-x)^2 \times (x + 2x^2 + 3x^3 + 4x^4 + 5x^5 \&c.)$
Whence, by multiplication,
$$x + 2x^2 + 3x^3 + 4x^4 \&c.$$

$$1 - 2x + x^2$$

$$x + 2x^2 + 3x^3 + 4x^4 \&c.$$

$$-2x^2 - 4x^3 - 6x^3 \&c.$$

$$+ x^3 + 2x^4 \&c.$$

Whose sum is =x+0+0+0+0+0 &c. 'Therefore z=x.

And
$$x+2x^2+3x^3+4x^4+5x^5$$
 &c. = $\frac{x}{(1-x)^2}$.

Or $\frac{1}{2} + \frac{2}{4} + \frac{3}{8} + \frac{4}{16} + \frac{5}{32} + \frac{6}{64} \&c. = \frac{\frac{1}{2}}{(1 - \frac{1}{2})^2} = 2 = \text{sum of the infinite series required.}$

16. It is required to find the sum (s) of the series $\frac{1}{3}$ + $\frac{4}{9}$ + $\frac{9}{27}$ + $\frac{16}{81}$ + $\frac{25}{243}$ &c. continued ad infinitum.

Let
$$x = \frac{1}{3}$$
 and $\frac{z}{(1-x)^3} = s$.

Then
$$\frac{1}{(1-x)^3} = x + 4x^2 + 9x^3 + 16x^4 + 25x^5 &c.$$

And $z = (1-x)^3 \times (x + 1x^2 + 9x^3 + 16x^4 &c.) = x + x^2$, as will be found by actual multiplication.

Therefore
$$x+x^2=z$$
,
And $x+4x^2+9x^3+16x^3\&c.+\frac{x(1+x)}{(1-x)^3}$.

 $\frac{1}{3} + \frac{4}{9} + \frac{9}{27} + \frac{16}{81} &c. = \frac{\frac{1}{3}(1 + \frac{1}{3})}{(1 - \frac{1}{3})^3} = \frac{3}{2} = 1\frac{1}{2} = \text{sum required.}$

17. Required the sum (s) of the series $\frac{a}{m} + \frac{a+d}{mr} + \frac{a+2d}{mr^2} + \frac{a+3d}{mr^3}$ &c. continued ad infinitum.

Let
$$x = \frac{1}{r}$$
, and $s = \frac{z}{m(1-x)^2}$
Then $\frac{z}{m(1-x)^2} = \frac{a}{m} + \frac{a+d}{mr} + \frac{a+2d}{mr^2} + \frac{a+3d}{mr^2} &c.$
Or, $\frac{a+d}{(1-x^2)} = a + \frac{a+d}{r} + \frac{a+2d}{r^2} + \frac{a+3d}{r^3} &c.$
That is, $\frac{z}{(1-s^2)^2} = a + (a+d)x + (a+2d)x^2 + (a+3d)x^3 + (a+4d)x^4 &c.$

 $u+(a+d)x+(a+2d)x^2+(a+3d)x^3+(a+4d)x^4 &c.$ And $z=(1-x)^2 \times \{a+(a+d)x+(a+2d)x^2+(a+3d)x^3 &c.\} = (1-x)a+dx,$

as will appear by actually multiplying by $(1-x)^2$

Therefore z = (1-x)a+dx; and consequently $\frac{a}{m}+\frac{a+d}{mr}+\frac{a+2d}{mr^2}\&c.=\frac{r}{m}\left\{\frac{a(r-1)+d}{(r-1)^2}\right\}=$ sum of the infinite series required.

EXAMPLES FOR PRACTICE.

1. Required the sum of 100 terms of the series 2, 5, 8, 11, 14, &2. Ans. 15050

2. Required the sum of 50 terms of the series $1+2^2+3^2+4^2+5^2$ &c. Ans. 42925

3. It is required to find the sum of the series 1+3x+

254

 $6x^2 + 10x^3 + 15x^4$ continued ad infinitum, &c. when x is less than 1.

Ans. $\frac{1}{(1-x)^3}$

4. It is required to find the sum of the series $1+4x + 10x^2 + 20x^3 + 35x^4$, &c. continued ad infinitum, when x is less than 1.

Ans. $\frac{1}{(1-x)^4}$

5. It is required to find the sum of the infinite series $\frac{1}{13} + \frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} &c.$ Ans. $\frac{5}{10}$, or $\frac{1}{2}$

6. Required the sum of 40 terms of the series (1×2) + (3×4) + (5×6) + (7×8) &c. Ans. 22960

7. Required the sum of n terms of the series $\frac{2x-1}{2x}$

 $-\frac{2x-3}{2x} + \frac{2x-5}{2x} + \frac{2x-7}{2x} &c.$ Ans. $n(\frac{2x-n}{2x})$

3. Required the sum of the infinite series $\frac{1}{1.2.3.4}$

 $\frac{1}{2.6.4.5} + \frac{1}{3.4.5.6} + \frac{1}{4.5.6.7} &c. Ans. \frac{1}{18}$

9. Required the sum of the series $\frac{1}{1} + \frac{1}{4} + \frac{1}{10} + \frac{1}{20} + \frac{1}{35}$

Sc. continued ad infinitum. Ans. $\frac{3}{2}$, or $1\frac{1}{2}$

10. It is required to find the sum of n terms of the series $1+8x+27x^2+64x^3+125x^4$, &c.

Ans. $\frac{1+4x+x^2}{(1-x)^4}$

11. Required the sum of n terms of the series $\frac{1}{r} + \frac{2}{r^2} + \frac{2}{r^2}$

 $\frac{3}{5^{13}} + \frac{4}{r^3} + \frac{5}{r^3} + \frac{6}{r^6} & \text{c.} \qquad \text{Ans. } \frac{1}{(r-1)} - \frac{1}{r^n} \left\{ \frac{nr + r - 1}{(r-1)^2} \right\}$

12. Required the sum of the series $\frac{1}{2.6} + \frac{1}{4.8} + \frac{1}{6.10}$

$$\frac{1}{3.12} \&c. \dots + \frac{1}{2n(4+2n)}.*$$
Ans. $\Sigma = \frac{3}{16}, s = \frac{5n+3n^2}{32+18n+16n^2}$

13. Required the sum of the series $\frac{1}{3.8} + \frac{1}{6.12} + \frac{1}{9.10}$

$$+\frac{1}{12.20} \&c. ... + \frac{1}{3n(4+4n)}.$$

Ans. $\Sigma = \frac{1}{12}$, $s = \frac{n}{12 + 12n}$

14. Required the sum of the series $\frac{6}{2.7} + \frac{6}{7.12} + \frac{6}{12.17}$

$$+\frac{6}{17.22}$$
 &c. . . . $+\frac{6}{(5n-3).(5n+2)}$.

Ans. $\Sigma = \frac{3}{5}$, $s = \frac{3n}{2+5n}$

15. Required the sum of the series $\frac{1}{3.6} - \frac{1}{6.8} + \frac{1}{9.10}$

$$\frac{1}{12.12} + &c. \cdot \cdot \pm \frac{1}{3n(4+2n)}.$$
Ans. $\Sigma = \frac{1}{24}$, $s = \frac{n}{2(3+6n)} = \frac{n}{4(6+6n)}$

16. Required the sum of the series $\frac{2}{3.5} - \frac{3}{5.7} + \frac{4}{7.9}$

$$\frac{5}{9(11} + &c. \quad .. \quad \pm \frac{1+n}{(1+2n)\cdot(3+2n)}.$$
Ans. $\Sigma = \frac{1}{12}$, $s = \frac{1}{12} - \frac{1}{4(3+4n)}$

^{*} The symbol Σ , made use of in these, and some of the following series, denotes the sum of an infinite number of terms, and S the sum of n terms.

17. Required the sum of the series
$$\frac{5}{1.2.3} + \frac{6}{2.3.4} + \frac{7}{3.4.5} + \frac{8}{4.5.6} & \dots + \frac{4+n}{n(1+n) \cdot (2+n)} \\ & \text{Ans. } \Sigma = \frac{3}{2}, s = \frac{3}{2} - \frac{2}{1+n} + \frac{1}{2+n} *$$

OF LOGARITHMS.

Logarithms are a set of numbers that have been computed and formed into tables, for the purpose of facilitating many difficult arithmetical calculations; being so contrived, that the addition and subtraction of them answers to the multiplication and division of natural numbers with which they are made to correspond.

^{*} The series here treated of are such as are usually called algebraical; which, of course, embrace only a small part of the whole doctrine. Those, therefore, who may wish for farther information on this abstruse but highly curious subject are referred to the Miscellanea Anniytica of Demoirre, Siering's Mithod Differ., James Bernoulli de Seri. Infin, Simpson's Math. Diss. sert., Waring's Medit. Anniyt., Clark's translation of Lorgna's Series, the various works of Euler, and Lacroix Traite du Calcul Diff. et Int., where they will find nearly all the materials that have been hitherto collected respecting this branch of analysis.

⁺ This mode of computation, which is one of the happiest and most useful discoveries of modern times, is due to Lord Napier, Berno of Merchiston, in Scotland, who first published a table of these numbers, in the year 1614, under the title of Comm Mirificum Logarithmorum; which performance was eagerly received by the learned throughout Europe, whose efforts were immediately directed to the improvement and extensions of the new calculus, that had so unexpectedly presented itself.

Mr. Henry Briggs, in particular, who was, at that time, professor of geometry in Gresham College, greatly contributed to the advancement of this doctaine, not only by the very advantageous alteration which he first introduced into the system of these numbers, by making 1 the legarithm of 10, instead of 2 3023552, as had been done by Napier; but also by the publication, in 1624 and 1633, of his two great works, the Arithmetica Logarithmica and the Trigonometria Britanies, both of which were formed upon the principle above mentioned; as are, likewise, all our common logarithmitic tables, at present in use

Or, when taken in a similar but more general sense, logarithms may be considered as the exponents of the powers to which a given, or invariable, number must be raised, in order to produce all the common, or natural numbers. Thus, if

 $a^{x}=y, a^{x'}=y', a^{x''}=y'', &c.$

then will the indices x, x', x'', &c. of the several powers of c, be the logarithms of the numbers y, y', y'', &c. in the scale, or system of which a is the base.

So, that, from either of these formulæ it appears, that the logarithm of any number, taken separately, is the index of that power of some other number, which when involved in the usual way, is equal to the given number.

And since the base σ , in the above expressions, can be assumed of any value, greater or less than 1, it is plain that there may be an endless variety of systems of logarithms, answering to the same natural number.

It is, likewise, farther evident, from the first of vese equations, that when y=1, x will be =0, whatever may be the value of a; and consequently the logarithm of 1 is always 0, in every system of logarithms.

And if x=1, it is manifest from the same equation that the base a will be =y; which base is, therefore the number whose proper logarithm, in the system to which it belongs, is 1.

Also, because ax = y, and ax' = y', it follows from the multiplication of powers, that $a^{x} \times a^{x'}$, or $a^{x^{x}} x' = yy'$; and consequently, by the definition of logarithms, given above, $x+x' = \log y$, or

 $\log y' = \log y + \log y'$.

And, for a like reason, if any number of the equations $a^x = y$, $a^x = y'$, $a^x = y''$, &c. be multiplied together we shall have $a^{x+x'+x''} e^x = yy'y'$ &c.; and consequently x + x' + x'' &c. = $\log_2 yy'y'$ &c.; or

 $\log yy'y''$ &c.; = $\log y + \log y' + \log y''$ &c.

See, for farther details on this part of the subject, the Introduction to my Treatise of Plane and Spherical Prigonometry, 8vo. 2d Edit. 1813; and for the construction and use of the tables consult those of Sherwin, Hutton, Taylor, Callet, and Borda, where every necessary information of this kind may be readily obtained.

From which it is evident, that the logarithm of the product of any number of factors is equal to the sum of the

logarithms of those factors.

Hence, if all the factors of a given number, in any case of this kind, be supposed equal to each other, and the sum of them be denoted by m, the preceding property will then become

 $\log y^m = m \log y$.

From which it appears, that the logarithm of the mth power of any number is equal to m times the logarithm of that number.

In like manner, if the equation $a^x = y$ be divided by $a^{x'} =$ y', we shall have, from the nature of powers, as before, $\frac{a^2}{a^2}$, or $a^2 = \frac{y}{y}$; and by the definition of logarithms, laid

down, in the first part of this article, $x-x'=\log \frac{y}{y'}$ or

$$\log \frac{y}{y} = \log y - \log y'$$
.

Hence the logarithm of a fraction, or of the quotient arising from dividing one number by another, is equal to the logarithm of the numerator minus the logarithm of the denominator.

And if each member of the common equation $a^x = y$ be raised to the fractional power denoted by $\frac{m}{r}$, we shall

have, in that case, $a^{-x} = y^{-x}$;

And, consequently, by taking the logarithms, as before,

 $\frac{m}{x} = \log_{x} \frac{m}{y^{n}}$, or $\log_{x} \frac{m}{y^{n}} = \frac{m}{n} \log_{x} y$.

Where it appears, that the logarithm of a mixed root, or gower, of any number, is found by multiplying the logarithm of the given number by the numerator of the index of that power, and dividing the result by the denomipator.

And if the numerator m, of the fractional index, be in this case, taken equal to 1, the above formula will then become

$$\log_{\cdot} y^{\frac{1}{n}} = \frac{1}{n} \log_{\cdot} y.$$

From which it follows, that the logarithm of the nth root of any number is equal to the nth part of the loga-

rithm of that number.

Hence, besides the use of logarithms in abridging the operations of multiplication and division, they are equally applicable to the raising of powers and extracting of roots; which are performed by simply multiplying the given logarithm by the index of the power, or dividing it by the number denoting the root.

But although the properties here mentioned are common to every system of logarithms, it was necessary, for practical purposes, to select some one of them from the rest, and to adapt the logarithms of all the natural num-

bers to that particular scale.

And as 10 is the base of our present system of arithmetic, the same number has accordingly been chosen for the base of the logarithmic system, now generally used.

So that, according to this scale, which is that of the

common logarithmic tables, the numbers . . . 10-4, 10-3, 10-2, 10-1, 100, 101, 102, 103, 104, &c.

Or
$$\frac{1}{10000}, \frac{1}{1000}, \frac{1}{100}, \frac{1}{10}, \frac{1}{10}, 1, 10, 100, 1000, 10000, &c.$$

10000'1000'100 10' have for their logarithms

 $\ldots -4, -3, -2, -1, 0, 1, 2, 3, 4, &c.$

Which are evidently a set of numbers in arithmetical progression, answering to another set in geometrical progression; as is the case in every system of logarithms.

And therefore, since the common, or tabular, logarithm of any number (n) is the index of that power of 10, which when involved, is equal to the given number, it is plain, from the following equation,

 $10^{x} = n$, or $10^{-x} = \frac{1}{n}$,

that the logarithms of all the intermediate numbers, in the above series, may be assigned by approximation, and made to occupy their proper places in the general scale.

It is also evident, that the logarithms of 1, 10, 100, 1000, &c. being 0, 1, 2, 3, &c. respectively, the logarithm of any number, falling between 0, and 1, will be 0 and some decimal parts; that of a number between 10 and 100, 1 and some decimal parts; of a number between 100 and 1000, 2 and some decimal parts; and so on, for other numbers of this kind.

And for a similar reason, the logarithms of $\frac{1}{10}$, $\frac{1}{100}$, $\frac{1}{1000}$, &c. or of their equals .1, .01, .001, &c. in the de-

scending part of the scale, being -1, -2, -3, &c. the logarithm of any number, falling between 0 and 1, will be -1, and some positive decimal parts; that of a number between .1 and .01, -2 and some positive decimal parts; of a number between .01 and .001, -3, and some positive decimal parts; &c.

Hence, likewise, as the multiplying or dividing of any number by 10, 100, 1000, &c. is performed by barely increasing or diminishing the integral part of i's logarithm by 1, 2, 3, &c. it is obvious that all numbers, which consist of the same figures, whether they be integral, fractional, or mixed, will have, for the decimal part of their logarithms, the same positive quantity.

So that, in this system, the integral part of any logarithm, which is usually called its index, or characteristic, is always less by I than the number of integers which the natural number consist of; and for decimals, it is the number which denotes the distance of the first significant figure

from the place of units.

Thus, according to the logarithmic tables in common use, we have

Logarithm
0.1361496 1.3021144
1.3021144
2.5253817
1.6676490
$\overline{2}.7891575$
&c.

Where the sign — is put over the index, instead of before it, when that part of the logarithm is negative, in order to distinguish it from the decimal part, which is always to be considered as +, or affirmative.

Also agreeably to what has been before observed, the logarithm of 38540 being 4.5859117, the logarithms of any other numbers, consisting of the same figures, will be as follows:

Numbers.	Logarithms
3854	3.5859117
385.4	2.5859117
38,54	1.5859117
3.854	0 5859117
.3854	1.5859117
.03854	2 5859117
.003854	3.5859117

Which logarithms, in this case, as well as in all others of a similar kind, whether the number contains ciphers or not differ only in their indices, the decimal, or positive part, being the same in them all.*

And as the indices, or integral parts, of the logarithms of any numbers whatever, in this system, can always be thus readily found from the simple consideration of the rule above mentioned, they are generally omitted in the tables, being left to be supplied by the operator, as occasion requires.

^{*} The great advantages attending the common, or Briggean system of logarithms, above all others, arise chiefly from the readiness with which we can always find the characteristic or integral part of any logarithm from the bare inspection of the natural number to which it belongs, and the circumstance, that multiplying or dividing any number by 10, 100, 1000, &c. only influences the characteristic of its logarithm, without allecting the decimal part. Thus, for instance, if i be made to denote the index or integral part of the logarithm of any number n, and d its decimal part, we shall have log n = 1 + d:

 $[\]log_{i} 10m \times n = (i+m) + d$; $\log_{i} \frac{n}{10m} = (i-m) + d$; where it is plain that the decimal part of the logarithm, in each of these cases, remains the same.

It may here, also, be farther added, that when the logarithm of a given number in any particular system, is known, it will be easy to find the logarithm of the same number in any other system, by means of the following equations,

ax = n, and ex = n; or $\log n = x$, and l. n = x. Where $\log n$ denotes the logarithm of n, in the system of which a is the base, and l, its logarithm in the system of which e is the base.

For, since $a^x = e^{x'}$, or $a^{\frac{x}{x'}} = e$, and $e^{\frac{x'}{x}} = a$, we shall have, for the base $a, \frac{x}{x} = \log_{1} e$, or $x = x' \log_{1} e$;

and for the base $e, \frac{x'}{n} = l$. a, or x' = x l. a.

Whence, by substitution, from the former equations,

$$\log n = l$$
, $n \times \log e$; or $\log n = l$, $n \times \frac{1}{l \cdot a}$,

Where the multiplier log. e, or its equal $\frac{1}{l.a}$, expresses the constant relation which the logarithms of n have to each other in the systems to which they belong.

But the only system of these numbers, deserving of notice, except that above described, is the one that furnishes what have been usually called hyperbolic or Neperian logarithms, the base e of which is 2.718281828459.

Hence, in comparing these with the common or tabular logarithms, we shall have, by putting a in the latter of the above formula=10, the expression.

log.
$$n=l$$
. $n \times \frac{1}{l \cdot n}$, or l . $n=\log n \times l$. 10.

Where log. in this case, denotes the common tabular logarithm of the number n, and l. its hyperbolic logarithm; the constant factor, or multiplier, $\frac{1}{l+10}$, which is

$$\frac{1}{2.3025850929}$$
, or its equal .4342944819,

being what is usually called the modulus of the common system of logarithms.*

PROBLEM I.

To compute the logarithm of any of the natural numbers 1, 2, 3, 4, 5, &c.

RULE, I

1. Take the geometrical series, 1, 10, 100, 1000, 10000, &c. and apply to it the arithmetical series, 0, 1, 2, 3, 4, &c. as logarithms.

2. Find a geometric mean between 1 and 10, 10 and 100, or any other two adjacent terms of the series, be-

twixt which the number proposed lies.

3. Also, between the mean, thus found, and the nearest extreme, find another geometrical mean, in the same manner; and so on, till you are arrived within the proposed limit of the number whose logarithm is sought.

4. Find, likewise, as many arithmetical means between the corresponding terms of the other series, 0, 1, 2, 3, 4, &c. in the same order as you found the geometrical ones, and the last of these will be the logarithm answering to the number required.

EXAMPLES.

Let it be required to find the logarithm of 3.

^{*} It may here be remarked, that, although the common logarithms have superseded the use of hyperbolic or Neperian logarithms, in all the ordinary operations to which these numbers are generally applied, yet the latter are not without some advantages peculiar to themselves; being of frequent occurrence in the application of the Fluxionary Calculus, to many analytical and physical problems, where they are required for the finding of certain fluents, which could not be so readily determined without their assistance; on which account great pains have been taken to calculate fables of hyperbolic logarithms, to a considerable extent, chiefy for this purpose. Mr. Barlow, in a Collection of Mathematical Tables lately published, has given them for the first 1000 numbers.

Here the proposed number lies between 1 and 10.

First, then, the log. of 10 is 1, and the log of 1 is 0.

Therefore $\sqrt{(10\times1)} = \sqrt{10} = 3$. 1622777 is the geometrical mean;

And $\frac{1}{2}(1+0)=\frac{1}{2}=.5$ is the arithmetical mean;

Hence the log. of 3.1622777 is .5.

Secondly the log. of 10 is 1, and the log of 3.1622777 is 5.

Therefore $\sqrt{(10 \times 3.1622777)} = 5.6234132$ is the geometrical mean:

And $\frac{1}{2}(1+.5)=.75$ is the arithmetical mean; Hence the log. of 5.6234132 is .75.

Thirdly, the log. of 10 is 1, and the log. of 5.6234132 is .75;

Therefore $\sqrt{(10 \times 5.6234132)} = 7.4989422$ is the geometrical mean:

And $\frac{1}{2}(1+.75)$ =.875 is the arithmetical mean ;

Hence the log. of 7.4989422 is .875.

Fourthly, the log. of 10 is 1, and the log. of 7.4989422 is .875;

Therefore $\sqrt{(10 \times 7.4989422)} = 8.6596431$ is the geometrical mean,

And $\frac{1}{2}(1+.875)=.9375$ is the arithmetical mean; Hence the log. of 8.6596431 is .9375.

Fifthly, the log. of 10 is 1, and the log. of 8.6596431 is .9375.

Therefore $\sqrt{(10 \times 3.6596431)} = 9.3057204$ is the geometrical mean,

And $\frac{1}{2}(1+.9375)=.96875$ is the arithmetical mean;

Hence the log. of 9.3057204 is .96875.

Sixthly, the log. of 8.6596431 is .9375, and the log. of 9.3057204 is .96875;

Therefore $\sqrt{(8.6596431 \times 9\ 3057204)} = 8.9768713$ is the geometrical mean,

And $\frac{1}{2}(.9375 + .96875) = .953125$ is the arithmetical mean;

Hence the log. of 8.9768713 is .953125.

And, by proceeding in this manner, it will be found, after 25 extractions, that the logarithm of 8.9999998 is .9542425;

which may be taken for the logarithm of 9, as it differs from it so little, that it may be considered as sufficiently exact for all practical purposes.

And in this manner were the logarithms of all the prime

numbers at first computed.

RULE II.

When the logarithm of any number (n) is known, the logarithm of the next greater number may be readily found from the following series, by calculating a sufficient number of its terms, and then adding the given logarithm to their sum.

to their sum.

$$\operatorname{Log.}(n+1) = \operatorname{log.} n + \operatorname{n'} \left\{ \frac{1}{2n+1} + \frac{1}{3(2n+1)^3} + \frac{1}{5(2n+1)^5} + \frac{1}{7(2n+1)^7} + \frac{1}{9(2n+1)^9} + \frac{1}{11(2n+1)^{11}} & \text{c.} \right\}$$

$$\operatorname{Or},$$

$$\operatorname{Log.}(n+1) = \operatorname{log.} n + \left\{ \frac{\operatorname{n'}}{2n+1} + \frac{\operatorname{A}}{3(2n+1)^2} + \frac{3\operatorname{B}}{5(2n+1)^2} + \frac{5\operatorname{C}}{7(2n+1)^2} + \frac{9\operatorname{E}}{9(2n+1)^2} & \text{c.} \right\}$$
Where new is a represent the terms immediately

Where A, B, C, &c represent the terms immediately preceding those in which they are first used, and n'= twice the modulus=.8685889638 *

EXAMPLES.

1. Let it be required to find the common logarithm of the number 2.

Here, because n+1=2, and consequently n=1 and 2n+1=3, we shall have

^{*} It may here be remarked, that the difference between the logarithms of any two consecutive numbers, is so much the less as the numbers are greater; and consequently the series which comprises the latter part of the above expression will in that case converge so much the faster. Thus $\log n$ and $\log (n+1)$, or its equal $\log n + \log (1+\frac{1}{n})$, will, obviously, differ but little from each other when n is a large number.

$$\frac{\frac{M}{2n+1}}{\frac{A}{3(2n+1)^2}} = \frac{.8685389638}{3} . = .289529654 (A)$$

$$\frac{A}{3(2n+1)^2} = \frac{.289529654}{3.3^2} . = .010723321 (B)$$

$$\frac{3B}{5(2n+1)^2} = \frac{3 \times .010723321}{5.3^2} = .000714888 (C)$$

$$\frac{5C}{7(2n+1)^2} = \frac{5 \times .000714888}{7.3^2} = .000056737 (D)$$

$$\frac{7D}{9(2n+1)^2} = \frac{7 \times .000056737}{9.3^2} = .000004903 (E)$$

$$\frac{9E}{11(2n+1)^2} = \frac{9 \times .000004903}{113(2n+1)^2} = .000000446 (F)$$

$$\frac{11S}{13(2n+1)^2} = \frac{11 \times .000000446}{15.3^2} = .000000042 (G)$$

$$\frac{13G}{15(2n+1)^2} = \frac{13 \times .000000042}{15.3^2} = .000000004 (H)$$
Sum of 8 terms . . .301029995
Add log. of 1 . . .000000000

Which logarithm is true to the last figure inclusively.

2. Let it be required to compute the logarithm of the number 3.

Here, since n+1=3, and consequently n=2, and 2n+1=5, we shall have

$$\frac{M'}{2n+1} = \frac{868588964}{5} . . = .173717793 \text{ (A)}$$

$$\frac{A}{3(2n+1)^2} = \frac{.173717793}{3.5^2} . = .002316237 \text{ (B)}$$

$$\frac{3}{5(2n+1)^2} = \frac{3+.002316237}{5.5^2} = .000055590 \text{ (c)}$$

$$\frac{5c}{7(2n+1)^2} = \frac{5 \times 000055590}{7.5^2} = .000001538 \text{ (b)}$$

7 _D	7×.00	000	158	38	υ,	000000050 (E
$\frac{9(2n+1)^2}{9\varepsilon}$	9 9×.00	.5 ²	000	50		000000000 (E
$11(2n+1)^3$	11	.52	-			00000002 (F
Sum of 6	terms					.176091260
Log. of 2		٠	٠	٠	٠	.301029995
Log. of 3						.477121255

Which logarithm is also correct to the nearest unit in the last figure.

And in the same way we may proceed to find the loga-

rithm of any prime number.

Also, because the sum of the logarithms of any two numbers gives the logarithm of their product, and the difference of the logarithms the logarithm of their quotient, &c.; we may readily compute, from the above two logarithms, and the logarithm of 10, which is 1, a great number of other logarithms, as in the following examples:

.301029995	Because 2×2=4, therefore log. 2	3.
2	Mult. by 2	
.602059990	gives log. 4	
.301029995	Because 2×3=6, therefore to log. 2	4.
.477121255		
.778151250	gives log. 6	
.301029995	Because 23=8, therefore log. 2 mult. by 3	5.
.903089985	gives log. 8	

6. Because 32=9, therefore log. 3 .477121255 mult. by 2 2
gives log. 9 .954242510
7. Because $\frac{1}{2}^0 = 5$, therefore from log. 10 1.000000000 take log. 2 .301029995
gives log. 5 .698970005
8. Because $3 \times 4 = 12$, therefore to log. 3 add log. 4 .602059991
gives log. 12 1.079181246

And, thus, by computing, according to the general formula, the logarithms of the next succeeding prime numbers 7, 11, 13, 17, 19, 23, &c. we can find, by means of the simple rules, before laid down for multiplication, division and the raising of powers, as many other logarithms as we please, or may speedily examine any logarithm in the table.

MULTIPLICATION

BY LOGARITHMS.

Take out the logarithms of the factors from the table, and add them together; then the natural number answer-

ing to the sum will be the product required.

Observing, in the addition, that what is to be carried from the decimal part of the logarithms is always affirmative, and must, therefore, be added to the indices, or integral parts, after the manner of positive and negative quantities in algebra.

Which method will be found much more convenient, to those who possess a slight knowledge of this science, than that of using the arithmetical complements.

EXAMPLES.

1. Multiply 37,153 by 4,086, by logarithms,

 $Nos. Logs. 37.153 \dots 1.5699939 + 0.086 \dots 0.6112984$

Prod. 151.8071 . 2.1812923

2. Multiply 112.246 by 13.958, by logarithms.

Nos. Logs.
112.246 . . . 2.0491709
13.958 . . . 1.1448232

Prod. 1563.128 . 3.1939941

3. Multiply 46.7512 by .3275, by logarithms.

 Nos.
 Logs.

 46.7512
 1.6697928

 .3275
 1.5152113

Prod. 15.31102 . 1.1850041

Here, the +1, that is to be carried from the decimals, cancels the -1, and consequently there remains 1 in the upper line to be set down.

4. Multiply .37816 by .04782, by logarithms.

Prod. .0180836 . 2.2572852

Here the + 1 that is to be carried from the decimals,

destroys the - 1, in the upper line, as before, and there remains the - 2 to be set down.

5. Multiply 3.768, 2.053, and .007693, together.

Nos.				Logs.
7.768				0.5761109
2 053				0.3123889
.007693		٠		3.8860997
ProdC	59	511	ı	2.7745995

Here the + 1, that is to be carried from the decimals, when added to -3, makes -2, to be set down.

6. Multiply 3.586, 2.1046, .8372, and .0294, together

Nos.					Logs.
3.586					0.554610
2.1046					0 323170
.8372					$\bar{1.922829}$
.0294					$\bar{2}.468347$
Prodi	.18	57	618		1.268956

Here the +2, that is to be carried, cancels the -2, and there remains the -1 to be set down.

7. Multiply 23.14 by 5.062 by logarithms.

Ans. 117.1347

8. Multiply 4.0763 by 9.8432, by logarithms.

Ans. 40.12383

9. Multiply 498.256 by 41.2467, by logarithms.

Ans. 20551.41 10. Multiply 4.026747 by .012345, by logarithms.

Ans. .0497102 379, together, by

11. Multiply 3.12567, .02868, and .12379, together, by logarithms.

Aps. .09109705

12. Multiply 2876.9, 10674, .098762, and .0031596, by logarithms. Ans. .0958299

DIVISION BY LOGARITHMS.

From the logarithm of the dividend, as found in the tables, subtract the logarithm of the divisor, and the natural number answering to the remainder, will be the quotient

required.

Observing, if the subtraction cannot be made in the usual way, to add, as in the former rule, the 1 that is to be carried from the decimal part, when it occurs, to the index of the logarithm of the divisor, and then this result, with its sign changed, to the remaining index, for the index of the logarithm of the quotient.

EXAMPLES.

1.	Divide	4768.2	by 3	6.9	54,	by	logarithms.
		Nos.				•	Logs.
		4768.2					3.6783545
		36.954					1.5676615
	-	Quot.	129.	032			2.1106930
2.	Divide	21.754	by s	2.46	78	, by	logarithms.
		Nos.					Logs.
		21.754				*	1.3375391
		2.4673					0.3923100
		Quot.	.815	18			0.9452291
3.	Divide	4.6257 Nos	by	.176	808	, b	y logarithms. Logs.
		4.625	7 .				0.6651725
		.17608	3 .			~	1.2457100
		Quot.	26.2	741		17	1.4194625

Here - 1, in the lower index, is changed into +1, which is then taken for the index of the result.

4. Divide .27684 by 5.1576, by logarithms.

Nos.					Logs.
.27684					1.4422988
5.1576				٠	0.7124477
Quot() 5 3	676	31		2.7297811

Here the 1 that is to be carried from the decimals, is taken as -1, and then added to -1, in the upper index, which gives -2 for the index of the result.

5. Divide 6.9875 by .075789, by logarithms.

Nos.	_			Logs.
6.9875	· .			0.8443218
.07578	9 .			2.8796062
Quot.	2,13	67		1.9647156

Here the 1, that is to be carried from the decimals, is added to -2, which makes -1, and this put down, with its sign changed, is +1.

6. Divide .19876 by .0012345, by logarithms.

No	5.					Logs.
.1987	76					1.2983290
.0019	234	5			••	3.0914911
Quot.	16	1.0	05	1		2.2068379

Here -3, in the lower index, is changed into +3, and this added to -1, the other index, gives +3 - 1 or 2.

7. Divide 125 by 1728, by logarithms.

Ans. .0723379 3. Divide 1728.95 by 1.10678, by logarithms.

Ans. 1562,144

3. Divide .1023674 by 4.96523, by logarithms.

Ans. 2.061685

10. Divide 19956.7 by .048235, by logarithms.

Ans. 413739

11. Divide .067859 by 1234.59, by logarithms.

Ans. .0000549648

THE RULE OF THREE,

OR PROPORTION, BY LOGARITHMS.

For any single proportion, add the logarithms of the second and third terms together, and subtract the logarithm of the first from their sum, according to the foregoing rules; then the natural number answering to the result will be the fourth term required.

But if the proportion be compound, add together the logarithms of all the terms that are to be multiplied. and from the result take the sum of the logarithms of the other terms, and the remainder will be the logarithm of

the term sought.

Or, the same may be performed more conveniently thus, Find the complement of the logarithm of the first term of the proportion, or what it wants of 10, by beginning at the left hand, and taking each of its figures from 9, except the last significant figure, on the right which must be taken from 10; then add this result and the logarithms of the other two terms together, and the sum, abating 10 in the index, will be the logarithm of the fourth term, as before.

And, if two or more logarithms are to be subtracted, as in the latter part of the above rule, add their complements and the logarithms, of the terms to be multiplied together, and the result, abating as many 10's in the index as there are logarithms to be subtracted, will be the logarithm of the term required; observing, when the index of the logarithm, whose complement is to be taken, is negative, to add it, as if it were affirmative, to 9; and then take the rest of the figures from 9, as before

EXAMPLES.

1. Find a fourth proportional to 37.125, 14.768, and 135.279, by logarithms.

Log. of 37.125 . . . 1.5696665

Gomplement . . . 8.4303335

Log. of 14.768 . . 1.1693217

Log. of 135.279 . . 2.1312304

Ans. 53.81099 . 1.7308856

2. Find a fourth proportion to .05764, .7186, and .34721, by logarithms.

3. Find a third proportional to 12.796 and 3.24718, by logarithms.

4. Find the interest of 2791. 5s. for 274 days, at 43 per cent. per annum, by logarithms.

Comp. log. of 100		8.0000000
Comp. log. of 365		7.4377071
Log. of 279.25 .		2.4459932
Log. of 274		2.4377506
Log. of 4.5		0.6532125
Ans. 9.433296	1	0.9746634

5. Find a fourth proportional to 12.678, 14.065, and 100.979, by logarithms.

Ans. 112.0263

6. Find a fourth proportional to 1.9864, .4678, and 50.4567, by logarithms. Ans. 11.88262

7. Find a fourth proportional to .09658, .24958, and .008967, by logarithms.

Ans. .02317234

8. Find a mean proportional between .498621 and 2.9587, and a third proportional to 12.796 and 3.24718 by logarithms.

Ans. 17.55623 and .8240216.

INVOLUTION,

OR THE RAISING OF POWERS BY LOGARITHMS.

Take out the logarithm of the given number from the tables, and multiply it by the index of the proposed power; then the natural number, answering to the result, will be the power required.

Observing, if the index of the logarithm be negative, that this part of the product will be negative; but as what is to be carried from the decimal part will be affirmative, the index of the result must be taken accordingly.

EXAMPLES.

t. Find the square of 2.7568, by logarithms.

2. Fi

Lcg. of 2.75	8		0.4402477
--------------	---	--	-----------

Square 7.599946	 0.8804954
ind the cube of 7. Log. of 7.0851	logarithms. 0.8503399 3
Cube 355.6475	2.5510197

3. Find the fifth power of .87451, by logarithms.

Log. of .87451 . . . 1.9417648

5

Fifth power .5114695 . 1.7088240

Where 5 times the negative index $-\overline{1}$, being -5, and +4 to carry, the index of the power is 1.

4. Find the 365th power of 1.0045, by logarithms

Log. 1.0045* . . 0.0019499

97495 116994

58497

Power 5.148888 . 0.7117135

5. Required the square of 6.05987, by logarithms.
Ans. 36.72203

6. Required the cube of .176546, by logarithms.
Ans. .005502674

^{*} This answer 5.143888 though found strictly according to the general rule, is not correct in the last four figures 8888; nor can the answers to such questions relating to very high powers be generally found true to 6 places of figures by the table of Log. commonly used; if any power above the hundred thousandth were required, not one figure of the answer here given could be dypended on. The Log. of 1.0045 is 00194994108 true to eleven places, which multiplied by 365 gives 7.117285 true to 7 places, and the corresponding number true to 7 places is 5.149067. See Doctor Adrain's addition of Hut. Math. Vol. 1. p. 169.

- 7. Required the 4th power of .076543, by logarithms. Ans. .0000343259
- 3. Required the 5th power of 2.97643 by logarithms.
 Ans. 233.6031
- 9. Required the 6th power of 21.0576 by logarithms. Ans. 87187340
- 10. Required the 7th power of 1.09684, by logarithms.

 Ans. 1.909864

EVOLUTION.

OR THE EXTRACTION OF ROOTS, BY LOGARITHMS.

Take out the logarithm of the given number from the table, and divide it by 2, for the square root, 3 for the cube root, &c. and the natural number answering to the result will be the root required.

But if it be a compound root, or one that consists both of a root and a power, multiply the logarithm of the given number by the numerator of the index, and divide the product by the denominator, for the logarithm of the root sought.

Observing, in either case, when the index of the logarithm is negative, and cannot be divided without a remainder, to increase it by such a number as will render it exactly divisible; and then carry the units borrowed, as so many tens, to the first figure of the decimal part, and divide the whole accordingly.

EXAMPLES.

1. Find the square root of 27.465, by logarithms. Log. of 27.465 . . 2)1.4387796

2.	Find the co	ube root of	35.6415, by logs	arithms.
	Log. of 3	5.6415 .	3)1.55195	60

3. Find the 5th root of 7.0825, by logarithms. Log. of 7.0825 . . . 5)0.8501866

4. Find the 365th root of 1.045, by logarithms. Log. of 1.045 . . . 365)0.0019499

Root 1.000121 . . . 0.0000534

5. Find the value of $(.001234)^{\frac{2}{3}}$ by logarithms. Log. of .00123430913152

3)6.1826304

Ans. .00115047 . . . 2.0608768

Here, the divisor 3 being contained exactly twice in the negative index -6, the index of the quotient, to be put down, will be -2.

Find the value of $(.024554)^{\frac{3}{2}}$ by logarithms.

Log. of .024554 . . . 2.3901223

2)5.1703669

Ans. .00384754 . . . 3.5851834

Here 2 not being contained exactly in -5, 1 is added to it, which gives -3 for the quotient; and the 1 that is bor-

rowed being carried to the next figure, makes 11, which, divided by 2, gives .58, &c.

7. Required the square root of 365.5674, by logarithms.
Ans. 19.11981

8. Required the cube root of 2.987635, by logarithms.
Ans. 1.440265

9. Required the 4th root of .967845, by logarithms.

Ans. .9918624

10. Required the 7th root of .098674, by logarithms.

Aus. .7183146

11. Required the value of $\left(\frac{21}{373}\right)^{\frac{2}{3}}$, by logarithms.

Ans. .146895

12. Required the value of $\left(\frac{112}{1727}\right)^{\frac{2}{5}}$, by logarithms.

MISCELLANEOUS EXAMPLES IN LOGARITHMS.

- 1. Required the square root of $\frac{2}{123}$, by logarithms.

 Ans. .1275153
- 2. Required the cube root of $\frac{1}{3,14159}$, by logarithms.
- Ans. .6827842 3. Required the .07 power of .00563, by logarithms.
- Ans. .6958821
 4. Required the value of $\frac{(\frac{2}{3})^{\frac{1}{2}} \times (\frac{3}{4})^{\frac{1}{3}}}{17!}$, by logarithms

Ans. .04279825

5. Required the value of $\frac{1}{7}\sqrt{\frac{5}{8}} \times .012\sqrt[3]{\frac{7}{11}}$, by logarithms. Aus. .001165713

6. Required the value of $\frac{\frac{1}{3}\sqrt{\frac{1}{2}\frac{1}{3}} \times .03\sqrt[3]{15\frac{1}{3}}}{7\frac{1}{3}\sqrt[3]{12\frac{1}{3}} \times .19\sqrt[3]{17\frac{1}{3}}}$ by logarithms.

Ans. .3009158638

7. Required the value of $\frac{127}{4} \left(\frac{5}{6} \sqrt{19 + \frac{4}{3}} \frac{3}{3} \cdot 35\frac{1}{3} \right)$, by logarithms. $\frac{127}{4} \left(\frac{127}{14} \frac{7}{19} - \frac{1}{17} \frac{3}{8} \cdot 26\frac{1}{3} \right)$

MISCELLANEOUS QUESTIONS.

1. A person being asked what o'clock it was, replied that it was between eight and nine, and that the hour and minute hands were exactly together; what was the time?

Ans 8h, 43min, 38f, sec.

2. A certain number, consisting of two places of figures, is equal to the difference of the squares of its digits, and if 36 be added to it the digits will be inverted; what

is the number?

3. What two numbers are those, whose difference, sum, and product, are to each other as the numbers 2, 3, and 5, respectively?

Ans. 2 and 10

4. A person, in a party at cards, betted three shillings to two upon every deal, and after twenty deals found he had gained five shillings; how many deals did he win?

Ans. 13
5. A person wishing to enclose a piece of ground with palisades, found, if he set them a foot asunder, that he should have too few by 150, but if he set them a yard asunder he should have too many by 70; how many had

he?

6. A cistern will be filled by two cocks, A and B, running together, in twelve hours, and by the cock A alone in twenty hours; in what time will it be filled by the cock

B alone? Ans. 30 hours 7. If three agents, A, B, c, can produce the effects a, b, c, in the times e, f, g, respectively; in what time would

they jointly produce the effect d.

Ans. $d \div (\frac{a}{e} + \frac{b}{f} + \frac{c}{g})$

8. What number is that, which being severally added to 3, 19, and 51, shall make the results in geometrical progression?

Ans. 13

9. It is required to find two geometrical mean proportionals between 3 and 24, and four geometrical means

between 3 and 96.

Ans. 6 and 12; and 6, 12, 24, and 48

10. It is required to find six numbers in geometrical progression such, that their sum shall be 315, and the sum of the two extremes 165.

Ans. 5, 10, 20, 40, 80, and 160

11. The sum of two numbers is a, and the sum of their reciprocals is b; required the numbers.

Ans. $\frac{a}{2} \pm \frac{1}{2} \checkmark \left\{ \frac{a}{b} (ab - 4) \right\}$

12. After a certain number of men had been employed on a piece of work for 24 days, and had half finished it, 16 men more were set on, by which the remaining half was completed in 16 days; how many man were employed at first; and what was the whole expence, at 1s. 6d. a day per man?

Ans. 32 the number of men; and the whole expence 1151.4s.

13. It is required to find two numbers such, that if the square of the first be added to the second, the sum shall

be 62, and if the square of the second be added to the first, it shall be 176.

Ans. 7 and 13

14. The fore wheel of a carriage makes six revolutions more than the hind wheel, in going 120 yards; but if the circumference of each wheel was increased by three feet, it would make only four revolutions more than the hind wheel in the same space; what is the circumference of each wheel?

Ans. 12 and 15 feet

15. It is required to divide a given number a into two such parts, x and y, that the sum of mx and ny shall be

equal to some other given number b.

Ans. $x = \frac{ba - n}{m - n}$ and $y = \frac{am - b}{m - n}$

16. Out of a pipe of wine, containing 84 gallons, 10 gallons were drawn off, and the vessel replenished with

10 gallons of water; after which, 10 gallons of the mixture were again drawn off, and then 10 gallons more of water poured in; and so on for a third and fourth time; which being done, it is required to find how much pure wine remained in the vessel supposing the two fluids to have been thoroughly mixed each time? Ans. 484 gallons

17. A sum of money is to be divided equally among a certain number of persons; now if there had been 3 claimants less, each would have had 150!. more, and if there had been 6 more, each would have had 120!. less; required

the number of persons, and the sum divided.

Ans. 9 persons, sum 2700l.

18. From each of sixteen pieces of gold, a person filed the worth of half a crown, and then offered them in payment for their original value, but the fraud being detected, and the pieces weighed, they were found to be worth in the whole, no more than eight guineas; what was the original value of each piece?

Ans. 13s.

19. A composition of tin and copper, containing 100 cubic inches, was found to weigh 505 ounces; how many ounces of each did it contain, supposing the weight of a cubic inch of copper to be 51 ounces, and that of a cubic

inch of tin 41 ounces.

Ans. 420 oz. of copper, and 85 oz. of tin

20. A privateer running at the rate of 10 miles an hour, discovers a vessel 18 miles a head of her, making way at the rate of 8 miles an hour; how many miles will the latter run before she is overtaken.

Ans. 72 miles

21. In how many different ways is it possible to pay 1001, with seven shilling pieces and dollars of 4s. 6d. each?

Ans. 31 different ways

22. Given the sum of two numbers = 2, and the sum of their ninth powers = 32, to find the numbers by a quadratic equation.

Aus. $1 \pm \frac{1}{4} \sqrt{(6\sqrt{34} - 33)}$.

23. Given $y^3 - xy = 666$, and $x^3 + xy = 406$; to find x and y.

Ans. x = 7, and y = 9.

24. The arithmetical mean of two numbers exceeds the geometrical mean by 13, and the geometrical mean exceeds the harmonical mean by 12; what are the numbers?

Ans. 234 and 104

25. Given $x^3y + y^3x = 3$, and $x^6y^2 + y^0x^2 = 7$, to find the values of x and y.

Ans. $x=\frac{1}{2}(\sqrt{5}+1), y=\frac{1}{2}(\sqrt{5}-1)$ 26. Given x+y+z=23, xy+xz+yz=167, and xyz=385, to find x, y, and z. Ans. x=5, y=7, z=11

27. To find four numbers, x, y, z, and w, having the product of every three of them given; viz. xyz=231. xyw = 420, yzw = 1540, and xzw = 660.

Ans. x=3, y=7, z=11, and w=2028. Given x+yz=384, y+xz=237, and z+xy=192. to find the values of x, y, and z.

Ans. x=10, y=17, and z=22

29. Given $x^2 + xy = 108$, $y^2 + yz = 69$, and $z^2 + xz = 580$, to find the values of x, y, and z.

Ans. x=9, y=3, and z=20

30. Given $x^2+xy+y^2=5$ and $x^4+x^2y^2+y^4=11$, to find the values of x and y by a quadratic.

Ans. $x = \frac{2}{5}\sqrt{10 + \frac{1}{5}}\sqrt{5}, y = \frac{2}{5}\sqrt{10 - \frac{1}{5}}\sqrt{5}$

Given the equation $x^4-6x^3+13x^2-12x=5$, to Ans. $\frac{3}{9} \pm \frac{1}{9} \sqrt{29}$ find the value of x by a quadratic.

32. It is required to find by what part of the population a people must increase annually, so that they may be double at the end of every century.

Ans. By 144th part nearly 33. Required the least number of weights, and the weight of each, that will weigh any number of pounds from one pound to a hundred weight.

Ans. 1, 3, 9, 27, 81

34. It is required to find four whole numbers such, that the square of the greatest may be equal to the sum of the squares of the other three. Ans. 3, 4, 12, and 13

35. It is required to find the least number, which being divided by 6, 5, 4, 3, and 2, shall leave the remainders 5, 4, 3, 2, and 1, respectively.

36. Given the cycle of the sun 18, the golden number 8, and the Roman indiction 10, to find the year.

Ans. 1717

37. Given 256x-87y=1, to find the least possible values of x and y in whole numbers. Ans. x=52, and y=153

38. It is required to find two different isosceles triangles such, that their perimeters and areas shall be both expressed by the same numbers.

Ans. Sides of the one 29, 29, 40; and of the other 37, 37, 24

39. It is required to find the sides of three right angled triangles, in whole numbers, such, that their areas shall be all equal to each other.

Ans. 58, 40, 42; 74, 24, 70; 113, 15, 112

40. Given x=1.2655, to find a near approximate va-Ans. 3.82013 lue of x.

41. Given xy = 5000, and yx = 3000, to find the values Ans. x=4.691445, and y=5.510132of x and y.

42. Given $x^{x} + y^{y} = 285$, and $y^{x} - x^{y} = 14$, to find the values of x and y. Ans. x=4.016698, and y=2.825716

43. To find two whole numbers such, that if unity be added to each of them, and also to their halves, the sums, Ans. 48 and 1680 in both cases, shall be squares.

44. Required the two least nonquadrate numbers x and y such, that $x^2 + y^2$ and $x^3 + y^3$ shall be both squares.

Ans. x=364 and y=273

45. It is required to find two whole numbers such that their sum shall be a cube, and their product and quotient Ans. 25 and 100, or 100 and 900, &c. squares.

46. It is required to find three biquadrate numbers such, that their sum shall be a square. Ans. 124, 154, and 204

47. It is required to find three numbers in continued geometrical progression such, that their three differences Ans. 567, 1008, and 1792 shall be all squares.

48. It is required to find three whole numbers such, that the sum or difference of any two of them shall be square Ans. 434657, 420968, and 150568

numbers.

49. It is required to find two whole numbers such, that their sum shall be a square, and the sum of their squares a Ans. 4565486027761 and 1061652293520

50. It is required to find four whole numbers such, that the difference of every two of them shall be a square num-Ans. 1873432, 2288168, 2399057, and 6560657

51. It is required to find the sum of the series $\frac{1}{3} + \frac{2}{9}$

 $+\frac{3}{27}+\frac{4}{81}+$ &c. continued to infinity. Ans. 3

52. It is required to find the sum of the infinite series $\frac{3}{4} - \frac{9}{16} + \frac{27}{64} - \frac{81}{256} + \frac{243}{1024} &c.$ Ans. $\frac{3}{4}$

53. Required the sum of the series 5+6+7+8+9+

&c. continued to n terms.

Ans. $\frac{n}{2}(n+9)$

54. It is required to find how many figures it would take to express the 25th term of the series 21+22+24+28+218&c.
Ans. 5050446 figures

55. It is required to find the sum of 100 terms of the series $(1\times2)+(3\times4)+(5\times6)+(7\times8)+(9\times10)$ &c.

Ans. 343800

56. Required the sum of $1^2+2^2+3^2+4^2+5^2$ &c. . . $+50^2$ which gives the number of shot in a square pile, the side of which is 50.

Ans. 42925

57. Required the sum of 25 terms of the series 35+36 ×2+37×3+38×4+39×5 &c. which gives the number of shot in a complete oblong pile, consisting of 25 tiers, the number of shot in the uppermost row being 35.

Ans. 16575

APPENDIX.

OF THE APPLICATION OF ALGEBRA TO GEOMETRY.

In the preceding part of the present performance, I have considered Algebra as an independent science, and confined myself chiefly to the treating on such of its most useful rules and operations as could be brought within a moderate compass; but as the numerous applications, of which it is susceptible, ought not to be wholly overlooked, I shall here show, in compliance with the wishes of many respectable teachers, its use in the resolution of geometrical problems; referring the reader to my larger work on this subject, for what relates more immediately to the general doctrine of curves.*

For this purpose it may be observed, that when any proposition of the kind here mentioned is required to be resolved algebraically, it will be necessary, in the first place, to draw a figure that shall represent the several parts, or conditions, of the problem under consideration, and to regard it as the true one.

Then, having properly considered the nature of the

And in such cases where it may be requisite to extend this mode of application to trigonometry, mechanics, or any other branch of mathematics, a previous knowledge of the nature and principles of these subjects will be equally necessary.

^{*} The learner before he can obtain a competent knowledge of the method of application above mentioned, must first make himself master of the principal propositions of Euclid, or of those contained in my Elements of Geometry; in which work he will find all the essential principles of the science comprised within a much shorter compass than in the former.

question, the figure so formed, must, if necessary, be still farther prepared for solution, by producing, or drawing, such lines in it as may appear, by their connexion or relations to each other, to be most conducive to the end proposed.

This being done, let the unknown line, or lines, which it is judged will be the easiest to find, together with those that are known, be denoted by the common algebraical symbols, or letters; then, by means of the proper geometrical theorems, make out as many independent equations as there are unknown quantities employed; and the resolution of these, in the usual manner, will give the solution of the problem.

But as no general rules can be laid down for drawing the lines here mentioned, and selecting the properest quantities to substitute for, so as to bring out the most simple conclusions, the best means of obtaining experience in these matters will be to try the solution of the same problem in different ways; and then to apply that which succeeds the best to other cases of the same kind, when they afterwards occur.

The following directions, however, which are extracted, with some alterations, from Newton's Universal Arithmetic. and Simpson's Algebra and Select Exercises, will often be found of considerable use to the learner, by showing him how to proceed in many cases of this kind, where he would

otherwise be left to his own judgment.

1st. In preparing the figure in the manner above mentioned, by producing or drawing certain lines, let them be either parallel or perpendicular to some other lines in it, or be so drawn as to form similar triangles; and, if an angle be given, let the perpendicular be drawn opposite to it, and so as to fall, if possible, from one end of a given line.

2d. In selecting the proper quantities to substitute for, let those be chosen, whether required or not, that are nearest to the known or given parts of the figure, and by means of which the next adjacent parts, may be obtained by addition or subtraction only, without using surds.

3d. When in any problem, there are two lines, or quantities, alike related to other parts of the figure or problem, the fibest way is not to make use of either of them separately but to substitute for their sum, difference, or rectangle, or the sum of their alternate quotients; or for some other line or lines in the figure, to which they have both the same relation.

4th. When the area or the perimeter, of a figure is given, or such parts of it as have only a remote relation to the parts that are to be found, it will sometimes be of use to assume another figure similar to the proposed one, that shall have one of its sides equal to unity, or to some other known quantity; as the other parts of the figure, in such cases, may then be determined by the known proportions of their like sides, or parts, and thence the resulting equation required.

These being the most general observations that have hitherto been collected upon this subject, I shall now proceed to elucidate them by proper examples; leaving such farther remarks as may arise out of the mode of proceeding here used, to be applied by the learner, as occasion requires to the solutions of the miscellaneous problems

given at the end of the present article.

PROBLEM T.

The base, and the sum of the hypothenuse and perpendicular of a right angled triangle being given, it is required to determine the triangle.



Let ABC, right angled at c, be the proposed triangle; and put BC = b, and AC = x.

Then, if the sum of AB and AC be represented by s, the hypothenuse AB will be expressed by s-x.

But, by the well known property of right angled triangles (Euc. 1. 47)

$$AC^2 + BC^3 = AB^2$$
, or $x^2 + b^2 = s^2 - 2sx + x^2$.

Whence, omitting x^2 , which is common to both sides of the equation, and transposing the other terms, we shall have $2sx=s^2-b^2$, or

$$x = \frac{s^2 - b^2}{\sigma_{\alpha}}, \dots, *$$

which is the value of the perpendicular ac; where s and b may be any numbers whatever, provided s be greater than b.

In like manner, if the base and the difference between the hypothenuse and perpendicular be given, we shall have, by putting x for the perpendicular and d+x for the hypothenuse.

$$x^{2}+2dx+d^{2}=t^{2}+x^{2}$$
, or $x=\frac{b^{2}-d^{2}}{2d}$.

Where the base (b) and the given difference (d) may be any numbers as before, provided b be greater than d.

PROBLEM II.

The difference between the diagonal of a square and one of its sides being given, to determine the square.



^{*} The edition of Euclid, referred to in this and all the following problems is that of Dr. Simson, London, 1801; which may also be used in the geometrical construction of these problems, should the student be inclined to exercise his talents upon this elegant but more difficult branch of the subject.

Let x be the proposed square, and put the side x, or x, x.

Then, if the difference of BD and BC be put =d, the hypothenuse BD will be =x+d.

But since, as in the former problem, BC² + CD², or 2BC²
=BD², we shall have

$$2x^2 = x^2 + 2dx + d^2$$
, or $x^2 - 2dx = d^2$.

Which equation being resolved according to the rule laid down for quadratics, in the preceding part of the work, gives

 $x=d+d\sqrt{2}$.

Which is the value of the side BC, as was required.

PROBLEM III.

The diagonal of a rectangle ABCD, and the perimeter, or sum of all its four sides, being given, to find the sides.



Let the diagonal ac=d, half the perimeter ab+bc=a, and the base bc=x; then will the altitude ab=a-x.

And since, as in the former problem, $AB^2 + BC^2 = AC^2$, we shall have

$$a^{2}-2ax+x^{2}+x^{2}=d^{2}$$
, or $x^{2}-ax=\frac{d^{2}-a^{2}}{2}$.

Which last equation, being resolved, as in the former instance, gives

 $x = \frac{1}{2}a \pm \frac{1}{2} \checkmark (2d^2 - a^2).$ Where a must be taken greater than d and less than $d \checkmark 2$.

PROBLEM IV.

The base and perpendicular of any plane triangle ABC being given, to find the side of its inscribed square.



Let Ee be the inscribed square; and put bc=b, Ab=p, and the side of the square EH or EF=x.

Then, because the triangles ABC, AEH, are similar, (Euc. vi. 4,) we shall have

AD : BC :: AI : EH, OF
$$p : b :: (p-x) : x$$
.

Whence, taking the products of the means and extremes, there will arise

px=bp-bx. Which by transposition and division, gives

$$x = \frac{bp}{b+p}$$
.

Where b and p may be any numbers whatever, either whole or fractional.

PROBLEM V.

Having the lengths of three perpendiculars, EF, EG, EH, drawn from a certain point E, within an equilateral triangle ABC, to its three sides, to determine the sides.



Draw the perpendicular AD, and having joined EA, EE, and EC, put EF=a, EG=b, EH=c, and ED (which is $\frac{1}{2}$ BC) = x.

Then, since AB, BC, or CA, are each =2x, we shall have, by Euc. 1, 47,

 $AD = \sqrt{(AE^2 - ED^2)} = \sqrt{(4x^2 - x^2)} = \sqrt{3x^2} = x\sqrt{3}$.

And because the area of any plane triangle is equal to half the rectangle of its base and perpendicular, it follows, that

$$\triangle ABC = \frac{1}{2}BC \times AD = x \times x \sqrt{3} = x^{2} \sqrt{3},$$

$$\triangle BEC = \frac{1}{2}BC \times EF = x \times a = ax,$$

$$\triangle AEC = \frac{1}{2}AC \times EG = x \times b = bx,$$

$$\triangle AEB = \frac{1}{2}AB \times EH = x \times c = cx.$$

But the last three triangles BEC, AEC, AEB, are together, equal to the whole triangle ABC, whence

 $x \sqrt{3} = ax + bx + cx$

And, consequently, if each side of this equation be divided by x, we shall have

$$x\sqrt{3}=a+b+c$$
, or $x=\frac{a+b+c}{\sqrt{3}}$.

Which is, therefore, half the length of either of the

three equal sides of the triangle.

Con. Since, from what is above shown, AD is $= x\sqrt{3}$, it follows, that the sum of all the perpendiculars, drawn from any point in an equilateral triangle to each of its sides, is equal to the whole persendicular of the triangle.

PROBLEM VI.

Through a given point P, in a given circle ACBD, to draw a cord cD, of a given length.



Draw the diameter APB; and put cD = a, AP = b, PB = c, and cP = x; then will PD = a - x.

But, by the property of the circle (Euc. 111. 35,) CP X PD=AP X PB; whence

$$x(a-x)=bc$$
, or $x^2-ax=-bc$.

Which equation, being resolved in the usual way gives $x = \frac{1}{2}a \pm \sqrt{(\frac{1}{4}a^2 - bc)}$;

Where x has two values, both of which are positive.

PROBLEM VII.

Through a given point P, without a given circle ABDC, to draw a right line so that the part cp, intercepted by the circumference, shall be of a given length.



Draw PAB through the centre o; and put cD=a, PA=b, PB=c, and PC=x; then will PD=x+a.

But, by the property of the circle, (Euc. 111, 36, cor.;)

$$x(x+a)=bc$$
, or $x^2+ax=bc$.

Which equation being resolved, as in the former prob-

 $x=-\frac{1}{2}a\pm\sqrt{(\frac{1}{4}a^2+bc)};$ Where one value of x is positive and the other negative.*

^{*} The two last problems, with a few slight alterations, may be readily employed for finding the roots of quadratic equations by construction; but this, as well as the methods frequently given for constructing cubic and some of the higher orders of equations is a matter of little importance in the present state of mathematical science; analysis, in these cases, being generally thought a more commodious instrument than geometry.

PROBLEM VIII.

The base BC, of any plane triangle ABC, the sum of the sides AB, AC, and the line AD, drawn from the vertex to the middle of the base, being given, to determine the triangle.



Put BD or DC=a, AD=b, AB+AC=s, and AB=x; then will AC=s-x.

But, by my Geometry, B. II, 19, $AB^2 + AC^2 = 2BD^2 + 2AD^2$; whence


$$x^2 + (s-x)^2 = 2a^2 + 2b^2$$
, or $x^2 - sx = a^2 + b^2 - \frac{1}{2}s^2$.

Which last equation, being resolved as in the former instances, gives $x=\frac{1}{a}s\pm\sqrt{(a^2+b^2-\frac{1}{a}s^2)}$,

for the values of the two sides AB and Ac of the triangle; taking the sign + for one of them, and - for the other and observing that a^2+b^2 must be greater than $\frac{1}{2}s^2$.

PROBLEM IX.

The two sides AB, AC, and the line AD, bisecting the vertical angle, of any plane triangle, ABC, being given to find the base BC.



Put AB=a, AC=b, AD=c, and BC=x; then, by Euc. vi. 3, we shall have

AB(a):AC(b)::BD:DC.

And, consequently, by the composition of ratios (Euc. v, 18,)

$$a+b:a::x:BD = \frac{ax}{a+b},$$

and
 $a+b:b::x:DC = \frac{bx}{a+b}$

$$a+b:b::x:DC=\frac{bx}{a+b}$$

But, by Euc. vi, 13, BD X DC + AD2 = AB X AC; wherefore, also,

$$\frac{abx^2}{(a+b)^2} + c^2 = ab$$
, or

 $abx^2 = (a+b)^2 \times (ab-c^2)$. From which last equation we have

$$x=(a+b)\sqrt{\frac{ab-c^2}{ab}};$$

Which is the value of the base BC, as required.

PROBLEM X.

Having given the lengths of two lines AD, BE, drawn from the acute angles of a right angled triangle ABC, to the middle of the opposite sides, it is required to determine the triangle.



Put AD = a, BE = b, CD or $\frac{1}{2}$ CB = x, and CE or $\frac{1}{2}$ CA = y; then, since (Euc. 1, 47) cd2+ca2=ad2, and cE2+cB2=BE2, we shall have

$$x^{2}+iy^{2}=a^{2},$$

 $y^{2}+4x^{2}=b^{2}.$

Whence, taking the second of these equations from four times the first, there will arise

$$15y^2 = 4a^2 - b^2$$
, or $y = \sqrt{\frac{4a^2 - b^2}{15}}$

And in like manner, taking the first of the same equations from four times the second there will arise

$$15x^2 = 4b^2 - a^2$$
, or $x = \sqrt{\frac{4b^2 - a^2}{15}}$

Which values of x and y are half the lengths of the base and perpendicular of the triangle; observing that b must be less than 2a, and greater than $\frac{1}{2}a$.

PROBLEM XI.

Having given the ratio of the two sides of a plane triangle ABC, and the segments of the base, made by a perpendicular falling from the verticle angle, to determine the friangle.



Put BD=a, DC=b, AB=x, AC=y, and the ratio of the sides as m to n.

Then, since by the question, AB: AC:: m: n, and by B. 11, 16, of my Elements of Geometry, AB²—AC²=BD²—DC², we shall have

$$x : y :: m : n, \text{ and } x^2 - y^2 = a^2 - b^2.$$

But, by the first of these expressions, nx = my, or $y = \frac{nx}{m}$; whence, if this be substituted for y in the second, there will arise

$$x^2 - \frac{n^2}{m^2}x^2 = a^{\frac{1}{2}} - b^2$$
, or

$$(m^2-n^2)x^2=m^2(a^2-b^2)$$

And consequently, by division and extracting the square root, we shall have

$$x=m\sqrt{\frac{a^2-b^2}{m^2-n^2}}$$
, and $y=n\sqrt{\frac{a^2-b^2}{m^2-n^2}}$;

which are the values of the two sides AB, Ac, of the triangle, as was required.

PROBLEM XII.

Given the hypothenuse of a right angled triangle ABC, and the side of its inscribed square Dc, to find the other two sides of the triangle.



Put AB=h, DE, OF DF=s, AC=x, and CE=y; then, by similar triangles, we shall have

$$AC(x) : CB(y) :: AF(x-s) : FD(s).$$

And, consequently, by multiplying the means and extremes,

$$xy-sy=sx$$
, or $xy=s(x+y)$, . . . (1)

But since, by Euc. I, 47, Ac2 + CB2 = AB2, we shall likewise have

$$x^2 + y^2 = h^2 \dots (2)$$

 $x^2+y^2=h^2$ (2) Whence, if twice equation (1) be added to equation (2), there will arise

$$x^2 + 2xy + y^2 = h^2 + 2s(x+y)$$
, or $(x+y)^2 - 2s(x+y) = h^2$.

Which equation, being resolved after the manner of a quadratic, gives

 $x+y=s+\sqrt{(h^2+s^2)}$, or

 $y=3-x\pm\sqrt{(h^2+s^2)}$. Hence, if this value be substituted for y in equation (1), there will arise

$$x \left\{ s - x \pm \sqrt{(h^2 + s^2)} \right\} = s \left\{ s \pm \sqrt{(h^2 + s^2)} \right\}, \text{ or }$$

$$x^2 - \left\{ s \pm \sqrt{(h^2 + s^2)} \right\} = s \left\{ s \pm \sqrt{(h^2 + s^2)} \right\}.$$

And, consequently, by resolving this last equation, we shall have

$$x = \frac{1}{2} \left\{ s \pm \sqrt{(h^2 + s^2)} \right\} \pm \sqrt{\left\{ \frac{1}{4} h^2 - \frac{1}{2} s^2 + \frac{s}{2} \sqrt{(h^2 + s^2)} \right\}}$$

$$y = \frac{1}{2} \left\{ s + \sqrt{(h^2 + s^2)} \right\} + \sqrt{\left\{ \frac{1}{4}h^2 - \frac{1}{2}s^2 + \frac{s}{2}\sqrt{(h^2 + s^2)} \right\}}$$

Which are the values of the perpendicular as and base ac, as was required.

PROBLEM XIII.

Having given the perimeter of a right angled triangle ABC, and the perpendicular co, falling from the right angle on the hypothenuse, to determine the triangle.



Put p=perimeter, cD=a, Ac=x, and Bc=y; then AB= p-(x+y.)

But, by right angled triangles (Euc. 1, 47) Ac2+Bc2= AB2; whence

$$x^2+y^2=p^2-2p(x+y)+x^2+2xy+y^2.$$
 Or, by transposing the terms and dividing by 2
$$p(x+y)-\frac{1}{2}p^2=xy. \qquad (1)$$

And since, by similar triangles, AB : BC :: AC: CD, we shall also have, by multiplying the means and extremes,

AB X CD=BC XAC, OF

ap-a(x+y)=xy. (2) Whence, by comparing equation (1) with equation (2) there will arise

$$(a+p) \times (x+y) = ap + \frac{1}{2}p^{2}.$$
Where
$$x+y = \frac{p(a+\frac{1}{2}p)}{a+p}, \text{ or }$$

$$y = \frac{p(a+\frac{1}{2}p)}{a+p} - x.$$

And if these values be now substituted for x+y and y in equation (2), the result, when simplified and reduced, will give

 $(a+p)x^2-p(a+\frac{1}{2}p)x=-\frac{1}{2}ap^2$. From which last equation and the value of y, above found, we shall have

$$x \text{ or } AC = \frac{p(a+\frac{1}{2}p)}{2(a+p)} + \frac{p}{2(a+p)} \checkmark \left\{ (a-\frac{1}{2}p)^2 - 2a^2 \right\}$$

$$y \text{ or } BC = \frac{p(a+\frac{1}{2}p)}{2(a+p)} + \frac{p}{2(a+p)} \checkmark \left\{ (a-\frac{1}{2}p)^2 - 2a^2 \right\}$$

And, if the sum of these two sides be taken from p, the result will give

$$AB = p - (x+y) = \frac{p^2}{2(a+p)}$$

Which expressions are, therefore, respectively equal to the values of the three sides of the triangle.

PROBLEM XIV.

Given the perpendicular, base, and sum of the sides of an obtuse angled plane triangle ABC, to determine the twe sides of the triangle.



Let the perpendicular AD=p, the base BC=b, the sum of AB and AC=s, and their difference =x.

Then, since half the difference of any two quantities added to half their sum gives the greater, and, when subtracted, the less, we shall have

$$AB = \frac{1}{2}(s+x)$$
, and $AC = \frac{1}{2}(s-x)$.

But, by Euc. 1, 47, $cD^2 = Ac^2 - AD^2$, or $cD = \sqrt{\frac{1}{4}(s-x)^2 - p^2}$; and, by B. II, 12, $AB^2 = BC^2 + AC^2 + 2BC \times CD$; whence

$$\frac{1}{4}(s+x)^2 = b^2 + \frac{1}{4}(s-x)^2 + 2b\sqrt{\frac{1}{4}(s-x)^2 - p^2}, \text{ or } sx - b^2 = 2b\sqrt{\frac{1}{4}(s-x)^2 - p^2}.$$

And if each of the sides of this last equation be squared, there will arise, by transposition and simplifying the result.

$$(s^2-b^2)x^2=b^2(s^3-b^2)-4b^2p^2$$
, or $x=b\sqrt{1-\frac{4p^2}{s^2-b^2}}$.

Whence, by addition and subtraction, we shall have


$$AB = \frac{s}{2} + \frac{b}{2} \sqrt{(1 - \frac{4p^2}{s^2 - b^2})}, \text{ and}$$

$$AC = \frac{s}{2} - \frac{b}{2} \sqrt{(1 - \frac{4p^2}{s^2 - b^2})}.$$

Whicheare the sides of the triangle, as was required.

PROBLEM XV.

It is required to draw a right line BFE from one of the angles B of a given square BD, so that the part FE, intercepted by DE and DC, shall be of a given length.



Bisect re in c, and put AB or BC=a, re or GE=b, and BG=x; then will BE=x+b and BF=x-b,

But since, by right angled triangles, AE2 = BE2 - AB2, we shall have

$$AE = \sqrt{\{(x+b)^2 - a^2\}}$$
.

And, because the triangles BCF, EAB, are similar, BF: BC: "BE: AE, OF

$$a(x+b)=(x-b)\sqrt{(x+b)^2-a^2}$$

Whence, by squaring each side of this equation, and arranging the terms in order, there will arise

 $x^4 - 2(a^2 + b^2)x^2 = b^2(2a^2 - b^2).$

Which equation being resolved after the manner of a quadratic, will give

 $x = \sqrt{a^2 + b^2 \pm a \sqrt{a^2 + 4b^2}}$

And, consequently, by adding b to, or subtracting it from this last expression, we shall have

BE =
$$\sqrt{\left\{a^2 + b^2 \pm a\sqrt{(a^2 + 4b^2)}\right\} + b}$$
, or BF = $\sqrt{\left\{a^2 + b^2 \pm a\sqrt{(a^2 + 4b^2)}\right\} - b}$.


Which values, by determining the point E, or F, will

satisfy the problem.

Where it may be observed, that the point c lies in the circumference of a circle, described from the centre p, with the radius rc, or half the given line.

PROBLEM XVI.

The perimeter of a right angled triangle ABC, and the radius of its inscribed circle being given, to determine the triangle.



Let the perimeter of the triangle =p, the radius on, or

OE, of the inscribed circle =r, $\Lambda E = x$, and ED = y,

Then, since in the right angled triangles AEO, AFO, OE is equal to OF, and OA is common, AF will also be equal AE, OF x.

And, in like manner, it may be shown, that BE is equal to

BD, or y.

But, by the question, and Euc. 1, 47, we have
$$(x+r)+(y+r)+(x+y)=p$$
, and $(x+r)^2+(y+r)^2=(x+y)^2$.

Or, by adding the terms of the first, and squaring those of the second.

 $x+y=\frac{1}{2}p-r$, and $r(x+y)=xy-r^2$.

Hence, since, in the first of these equations, $y = (\frac{1}{2}p - r) - x$, if this value be substituted for y in the second, there will arise

 $x^2 - (\frac{1}{2}p - r)x = -r(\frac{1}{2}p - r)$. Which equation, being resolved in the usual manner,

gives

$$x = \frac{1}{2} \left(\frac{1}{2} p - r \right) \pm \sqrt{\frac{1}{4} \left(\frac{1}{2} p - r \right)^2 - r \left(\frac{1}{2} p - r \right)},$$
and

$$y = \frac{1}{2}(\frac{1}{2}p - r) \mp \sqrt{\frac{1}{4}(\frac{1}{2}p - r)^2 - r(\frac{1}{2}p - r)}.$$

And, consequently, if r be added to each of these last expressions, we shall have

$$AC = \frac{1}{2} (\frac{1}{2}p + r) \pm \sqrt{\left\{ \frac{1}{4} (\frac{1}{2}p - r)^2 - r(\frac{1}{2}p - r) \right\}},$$

$$BC = \frac{1}{2} (\frac{1}{2}p + r) \mp \sqrt{\frac{1}{4} (\frac{1}{2}p - r)^2 - r(\frac{1}{2}p - r)},$$

for the values of the perpendicular and base of the triangle, as was required.

PROBLEM XVII.

From one of the extremities A, of the diameter of a given semicircle ADE, to draw a right line AE, so that the part DE, intercepted by the circumference and a perpendicular drawn from the other extremity, shall be of a given length.



Let the diameter AB=d, DE=a, and AE=x; and join BD. Then, because the angle ADB is a right angle, (Euc. 111, 31,) the triangles ABE ABD, are similar.

And consequently, by comparing their like sides, we shall have

x:d::d:x-a. Whence, multiplying the means and extremes of these proportionals, there will arise


 $x^2 - ax = d^2,$

Which equation, being resolved after the usual manner, gives

$$x = \frac{1}{2}a + \sqrt{(\frac{1}{4}a^2 + d^2)}$$
.

PROBLEM XVIII.

To describe a circle through two given points A, E, that shall touch a right line on given in position.



Join AB; and through o, the assumed centre of the required circle, draw FE perpendicular to AB; which will bisect it in E (Euc. 111, 3).

Also, join os; and draw en, oc, perpendicular to co; the latter of which will fall on the point of contact of

(Euc. 111. 18).

Hence, since A, E, B, H, F, are given points, put EB=a, EF=b, EH=c, and EO=x; which will give OF=b-x.

Then, because the triangle ore is right angled at r, we shall have

$$oB^2 = Eo^2 + EB^2$$
, or $oB = \sqrt{(x^2 + a^2)}$.

But, by similar triangles, FE : EH :: FO : OG OF OF; OF b:c:b-x:oB; whence, also,

$$OB = \frac{c}{b}(b-x).$$

And, consequently, if these two values of on be put equal to each other, there will arise

$$\checkmark(x^2+a)=\frac{c}{b}(b-x).$$

Or, by squaring each side of this equation, and simplifying the result,

 $(b^2-c^2)x^2+2bc^2x=b^2(c^2-a^2).$

Which last equation, when resolved in the usual manner, gives

$$x = -\frac{bc^2}{b^2 - c^2} + b\sqrt{\frac{c^4}{(b^2 - c^2)^2} + \frac{c^2 - a^2}{b^2 - c^2}},$$

for the distance of the centre o from the chord AB; where b must, evidently, be greater than c, and c greater than a.

PROBLEM XIX.

The three lines AO, BO, CO, drawn from the angular points of a plane triangle ABC, to the centre of its inscribed circle being given, to find the radius of the circle, and the sides of the triangle.

ALGEBRA TO GEOMETRY.



Let o be the centre of the circle, and, on so produced, let fall the perpendiculars co; and draw or, or, oc, to the

points of contact E, F, G.

Then, because the three angles of the triangle ABC are, together, equal to two right angles, (Euc. 1, 32.) the sum of their halves OAC+OCA+OBE will be equal to one right angle.

But the sum of the two former of these, oac+oca, is equal to the external angle Doc; whence the sum of Doc +obe, as also of Doc+ocn, is equal to a right angle; and,

consequently, obe=oco.

Let therefore, A0=a, B0=b, G0=c, and the radius OE, OF OF OG=x.

Then since the triangles BOE, COD are similar, BO OE :: CO : OD, or b : x :: c : OD; which gives

on =
$$\frac{cx}{b}$$
, and cn = $\sqrt{(c^2 - \frac{c^2 x^2}{b^2})}$ or $\frac{c}{b}\sqrt{(b^2 - x^2)}$.

Also, because the triangle Aoc is obtuse angled at o, we shall have (Euc. II, 12.)

$$Ac^{2} = Ac^{2} + co^{2} + 2Ao \times od; \text{ or } Ac = \sqrt{(a^{2} + c^{2} + \frac{2acx}{b})} \text{ or } \sqrt{(\frac{b(a^{3} + c^{2}) + 2acx}{b})}.$$

But the triangles ACD, AOF, being likewise similar, AC : CD :: AO : OF, OF

$$\sqrt{\frac{b(a^2+c^2)+2acx}{b}} : \frac{c}{b} \checkmark (b^2-x^2) :: a:x.$$

Whence, multiplying the means and extremes, and squaring the result, there will arise

 $bx^{2} \left\{ b(a^{2}+c^{2})+2acx \right\} = a^{2}c^{2}(b^{2}-x^{2}).$

Or, by collecting the terms together, and dividing by the coefficient of the highest power of x. D d 2.

$$x^3 + \left(\frac{ab}{2c} + \frac{ac}{2b} + \frac{bc}{2a}\right)x^2 = \frac{abc}{2}.$$

From which last equation x may be determined, and thence the sides of the triangle.* 3

PROBLEM XX.

Given the three sides AB, BC, CD, of a trapezium ABCD, inscribed in a semicircle, to find the diameter, or remaining side AD.



Let AB = a, BC = b, CD = c, and AD = x; then, by Euc. VI, D. AC \times BD = AD \times BC + AB \times CD = bx + ac.

But ABD, ACD, being right angles, (Euc. III, 31,) we shall have

$$AC = \sqrt{(AD^2 - DC^2)}$$
, or $\sqrt{(x^2 - c^2)}$, and $BD = \sqrt{(AD^2 - AB^2)}$, or $\sqrt{(x^2 - a^2)}$.

Whence, by substituting these two values in the former expression, there will arise.

 $\sqrt{(x^2-\varepsilon^2)} \times \sqrt{(x^2-a^2)} = bx + a\varepsilon.$

Or, by squaring each side, and reducing the result, $x^3 - (a^3 + b^2 + c^2)x = 2abc$.

From which last equation the value of x may be found, as in the last problem. †

† Newton, in his Universal Arithmetic, English edition, 1728, has resolved this problem in a variety of different ways, in order to show, that some methods of proceeding, in cases of this kind frequently lead to more elegant sobitions than others; and that a ready knowledge of these can only be obtain-

ed by practice.

^{*} This, and the following problem, cannot be constructed geometrically, or by means only of right lines and a circle, being what the ancients usually denominated solid problems, from the circumstance of their involving an equation of more than two dimensions; in which cases they generally employed the conic sections, or some of the higher orders of curves.

MISCELLANEOUS PROBLEMS.

PROBLEM 1.

To find the side of a square, inscribed in a given semicircle, whose diameter is d.

Ans. $\frac{1}{5}d\sqrt{5}$

PROBLEM II.

Having given the hypothenuse (13) of a right angled triangle, and the difference between the other two sides (7), to find these sides*

Ans. 5 and 12

PROBLEM III.

To find the side of an equilateral triangle, inscribed in a circle whose diameter is d; and that of another circumscribed about the same circle.

Ans. $\frac{1}{2}d\sqrt{3}$, and $d\sqrt{3}$

PROBLEM IV.

To find the side of a regular pentagon, inscribed in a circle, whose diameter is d.

Ans. $\frac{1}{4}d\sqrt{(10-2\sqrt{5})}$

PROBLEM V.

To find the sides of a rectangle, the perimeter of which shall be equal to that of a square, whose side is a, and its area half that of a square. Ans. $a+\frac{1}{2}a\sqrt{2}$ and $a-\frac{1}{2}a\sqrt{2}$

PROBLEM VI.

Having given the side (10) of an equilateral triangle, to find the radii of its inscribed and circumscribing circles.

Ans. 2.8868 and 5.7736

^{*} Such of these questions as are proposed in numbers, should first be resolved generally, by means of the usual symbols and then reduced to the answers above given, by substituting the numeral values of the letters in the results thus obtained.

PROBLEM VII.

Having given the perimeter (12) of a rhombus, and the sum (8) of its two diagonals, to find the diagonals.

Ans. 4+ 1/2 and 4-1/2

PROBLEM VIII.

Required the area of a right angled triangle, whose bypothenuse is x^{3x} , and the base and perpendicular x^{2x} and x^{2x} .

Ans. 1.029085

PROBLEM IX.

Having given the two contiguous sides (a, b) of a parallelogram, and one of its diagonals (d), to find the other diagonal.

Ans. $\sqrt{(2a^2+2b^2-d^2)}$

PROBLEM X.

Having given the perpendicular (300) of a plane triangle, the sum of the two sides (1156), and the difference of the segments of the base (495), to find the base and the sides.

Ans. 945, 375, and 780

PROBLEM XI.

The lengths of three lines drawn from the three angles of a plane triangle to the middle of the opposite sides, being 18, 24, and 30, respectively; it is required to find the sides.

Ans. 20, 28.844, and 34.176

PROBLEM XII.

In a plane triangle, there is given the base (50), the area (796), and the difference of the sides (10), to find the sides and the perpendicular.

Ans. 36, 46, and 33.261

PROBLEM XIII.

Given the base (194) of a plane triangle, the line that bisects the vertical angle (66), and the diameter (200) of the circumscribing circle, to find the other two sides.

Ans. 81.36587 and 157.43865

309

PROBLEM XIV.

The lengths of two lines that bisect the acute angles of a right angled plane triangle, being 40 and 50 respectively, it is required to determine the three sides of the triangle.

Ans. 35.80737, 47.40728, and 59.41143

PROBLEM XV.

Given the altitude (4), the base (8), and the sum of the sides (12), of a plane triangle, to find the sides.

Ans. $6 + \frac{4}{5} \sqrt{5}$ and $6 - \frac{4}{5} \sqrt{5}$

PROBLEM XVI.

Having given the base of a plane triangle (15), its area (45), and the ratio of its other two sides as 2 to 3, it is required to determine the lengths of these sides.

Ans. 7.7915 and 11.6872.

PROBLEM XVII.

Given the perpendicular (24), the line bisecting the base (40), and the line bisecting the vertical angle (25) to determine the triangle.

Ans. The base $\frac{250}{7}\sqrt{7}$

From which the other two sides may be readily found.

PROBLEM XVIII.

Given the hypothenuse (10) of a right angled triangle, and the difference of two lines drawn from its extremities to the centre of the inscribed circle (2), to determine the base and perpendicular.

Ans. 8.08004 and 5.87447

PROBLEM XIX.

Having given the lengths (a, b,) of two chords, cutting each other at right angles, in a circle, and the distance (c) of their point of intersection from the centre, to determine the diameter of a circle.

Ans. $\sqrt{\{8(a^2+b^2)+2c^2\}}$

PROBLEM XX.

Two trees, standing on a horizontal plane, are 120 feet asunder; the height of the highest of which is 100 feet, and that of the shortest 80; whereabouts in the plane must a person place himself, so that his distance from the top of each tree, and the distance of the tops themselves, shall be all equal to each other?

Ans. 20/21 feet from the bottom of the shortest, and 40/3 feet from the bottom of the other.

PROBLEM XXI.

Having given the sides of a trapezium, inscribed in a circle, equal to 6, 4, 5, and 3, respectively, to determine the diameter of the circle.

Ans. $\frac{1}{20}\sqrt{(130\times153)}$ or 7.051595

PROBLEM XXII.

Supposing the town A to be 30 miles from B, B 25 miles from c, and c 20 miles from A; whereabouts must a house be erected that it shall be at an equal distance from each of them?

Ans. 15.118556 miles from each

PROBLEM XXIII.

Given the area (100) of an equilateral triangle ABC, whose base BC falls on the diameter, and vertex A in the middle of the arc of a semicircle; required the diameter of the semicircle.

Ans. 201/3

PROBLEM XXIV.

In a plane triangle, having given the perpendicular (p), and the radii (r, n) of its inscribed and circumscribing circles, to determine the triangle.

Ans. The base $\frac{2r\sqrt{(2p_R-4r_R-r^2)}}{p-2r}$

PROBLEM XXV.

Having given the base of a plane triangle equal to 2a, the perpendicular equal to a, and the sum of the cubes of its other two sides equal to three times the cube of the base; to determine the sides.

Ans. $a(2+\frac{1}{3}\sqrt{6})$ and $a(2-\frac{1}{3}\sqrt{6})$

ERBATA.

N. B. All the lines in the second column are reckoned from the top.


PAGES. LINES.

```
хi
                For Simpson's, read Simpson.
                For Addition, read Subtraction.
12
                For -\frac{4c}{d}, read +\frac{4c^2}{d}.
       20
14
       7
                For 6a read -6a.
16
     8 9 10
                For 12ax, -2xy, and 6ab, read 12a2x,
                           -2x^2y, and 6a^4b.
17
       16
                For -y, read -y^2.
26
                For 8b, read 834
       25
                For \frac{5x^2}{2}, read \frac{5x^2}{2b}.
41
       17
                For -\frac{b^2}{a}, read -\frac{b^2}{a^2}.
42
         8
51
        16
                For 4a2x, read 4a3x.
                For x-1, read x^2-2x+1.
53
       22
                 For \sqrt{\frac{a}{a}}, read \sqrt{\frac{a}{h^2}}.
55
        11
59
        24
                 For x\sqrt{x-a^2}, read x\sqrt{(x-a^2)},
                 For 413/a and 223/ab, read 413/a and 25
66
        15
                            3/ab.
                 For 3/(6(1+/2+/4)), read 3/(6(1+
71
        18
                     3/2+3/4)).
                 For \sqrt{(a+\sqrt{-b})}, read \sqrt{(a-\sqrt{-b})}.
For \sqrt{(ax-x^2)}, read \sqrt{(a^2-x^2)}.
 72
        14
 74
        12
 76
                 For or, dele or.
         4
                 For -(x^2-y), read -(x^2-y^2).
        32
```


ERRATA.

PAGES.	LINES.	
80	4 8 23	For 3/9, -1/2/-5, and Cardaus', read 3/
		9, -1/5, and Cardan's.
36	12	For $\sqrt{10-2}\sqrt{2+(2-\sqrt{5})}\times \frac{4}{5}$, read 4
		$\{-\sqrt{10-2}\sqrt{2}+(2+\sqrt{5})\times\sqrt{5}\}$
103	23	For 291, read 9.
114	3 13	For $x^2 + y^2$, read $x^2 - y^2$, and for 13+,
		read +13.
135	37	For $x10-x$, read x , $10-x$.
155	14	For a+c, read d+c.
157	1	For 97h3 rend 97h2

A STATE OF THE STA







911366 Q 7

THE UNIVERSITY OF CALIFORNIA LIBRARY

JLLIMORE'S
OK STORE,
Park Street,
BALTIMORE.

